

Reg no: 54022-8

PM-15

18/17

<p>US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460</p> <p>NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)</p>	<p>EPA REGISTRATION NO. 54022-8</p> <p>DATE OF ISSUANCE JUL 26 1990</p> <p>TERM OF ISSUANCE Until Reregistration</p> <p>NAME OF PESTICIDE PRODUCT Protection 15 Plus Flea & Tick</p>
<p>NAME AND ADDRESS OF REGISTRANT (Include ZIP code)</p> <p style="text-align: right;">160/249264 18/2</p> <div style="border: 1px solid black; padding: 5px; text-align: center; margin: 10px auto; width: 200px;">BEST AVAILABLE COPY</div>	
<p>NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.</p>	
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.</p> <p>A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.</p> <p>Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p>	
<p><input type="checkbox"/> ATTACHMENT IS APPLICABLE</p> <p>SIGNATURE OF APPROVING OFFICIAL <i>[Signature]</i> DATE <i>[Date]</i></p>	

Richards

2 of 17

BLISTER PACKAGE

FRONT PANEL (CARDBOARD BACKING)

**PROTECTION™ Plus 150 Flea and Tick Collar for Dogs
with EFA***

*Slow release coat conditioner helps enhance texture, manageability and luster.

**Kills fleas for 5 months
Kills ticks for 5 months**

(Color of collar visible through affixed blister package)

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-	
methyl-4-pyrimidinyl)	
phosphorothioate]	11.0 %
INERT INGREDIENTS*	89.0 %
TOTAL	100.0%

*Includes 5% unsaturated fatty acids as a grooming aid.

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

For neck size up to 22".

FRANCODEX

**CONTAINS 1 FLEA AND TICK COLLAR
NET WT. 1.3 OZ. (36.0 g)**

54022-I

FRANCODEX
 54022-8
 1.3 OZ. (36.0 g)
 Contains 1 Flea and Tick Collar
 For neck size up to 22"
 Kills fleas for 5 months
 Kills ticks for 5 months
 *Includes 5% unsaturated fatty acids as a grooming aid.
 KEEP OUT OF REACH OF CHILDREN
 CAUTION: See back panel for additional precautionary statements and directions.
 For neck size up to 22".
 FRANCODEX
 CONTAINS 1 FLEA AND TICK COLLAR
 NET WT. 1.3 OZ. (36.0 g)

BLISTER PACKAGE**BACK PANEL (CARDBOARD BACKING)****PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: This product contains a cholinesterase inhibitor. Atropine sulfate and 2-PAM are antidotal only if signs of cholinesterase inhibition are present.

Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing dogs.

Adjustable - fits mature dogs, 22" neck or smaller

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labeling.

Place Protection™ Plus 150 collar around dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 150 days. For unusually large breeds, collar aids in tick control in the head and neck area for up to 2 months. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

This Protection™ Plus 150 collar contains Essential Fatty Acids necessary for healthy skin and hair. Unsaturated fatty acids released from the collar are rapidly absorbed to help maintain a lustrous, manageable coat.

DISPOSAL

Do not reuse empty blister. Wrap blister and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 54022 xx-xxx

EPA Est. 15577-FR-1

Manufactured for Francodex Laboratories, Inc.
Subsidiary of Virbac, Inc.
Fort Worth, Texas 76118
Made in France

FOIL POUCH/CARTON PACKAGE

FRONT PANEL (CARTON)

**PROTECTION™ Plus 150 Flea and Tick Collar for Dogs
with EFA*
(color)**

*Slow release coat conditioner helps enhance texture, manageability and luster.

**Kills fleas for 5 months
Kills ticks for 5 months**

For neck size up to 22".

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)

phosphorothioate] 11.0 %

INERT INGREDIENTS* 89.0 %

TOTAL 100.0%

*Includes 5% unsaturated fatty acids as a grooming aid.

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

FRANCODEX

**CONTAINS 1 FLEA AND TICK COLLAR
NET WT. 1.3 OZ. (36.0 g)**

FOIL POUCH/CARTON PACKAGE

BACK PANEL(CARTON)

**PROTECTION™ Plus 150 Flea and Tick Collar for Dogs with EFA (Color)
Adjustable - fits mature dogs, 22" neck or smaller.**

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: This product contains a cholinesterase inhibitor. Atropine sulfate and 2-PAM are antidotal only if signs of cholinesterase inhibition are present.

Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing dogs.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labeling.

Place Protection™ Plus 150 collar around the dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 150 days. For unusually large breeds, collar aids in tick control in the head and neck area for up to 2 months. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

This Protection™ Plus 150 collar contains Essential Fatty Acids necessary for healthy skin and hair. Unsaturated fatty acids released from the collar are rapidly absorbed to help maintain a lustrous, manageable coat.

DISPOSAL

Do not reuse empty pouch or box. Wrap pouch or box and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 54022-xx-xxxxx

EPA Est. 15577-FR-1

Manufactured for Francodex Laboratories, Inc.

Subsidiary of Virbac, Inc.

Fort Worth, Texas 76118

Made in France

FOIL POUCH/CARTON PACKAGE

POUCH LABEL

**PROTECTION™ Plus 150 Flea and Tick Collar for Dogs
with EFA**

Do not open until ready to use

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

**SEE BACK PANEL OF OUTER CARTON FOR
ADDITIONAL PRECAUTIONARY STATEMENTS**

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-	
methyl-4-pyrimidinyl)	
phosphorothioate]	11.0 %
INERT INGREDIENTS*	89.0 %
TOTAL	100.0%

*Includes 5% unsaturated fatty acids as a grooming aid

EPA Reg. No. 54022-xx-xxxxx

EPA Est. 15577-FR-1

**Manufactured for Francodex Laboratories, Inc.
Subsidiary of Virbac, Inc.
Fort Worth, Texas 76118
Made in France**

7 of 17

BLISTER PACKAGE

FRONT PANEL (CARDBOARD BACKING)

**PROTECTION™ Plus 150 Flea and Tick Collar for Large Dogs
with EFA***

*Slow release coat conditioner helps enhance texture, manageability and luster.

**Kills fleas for 5 months
Kills ticks for 5 months**

(Color of collar visible through affixed blister package)

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-	
methyl-4-pyrimidinyl)	
phosphorothioate]	11.0 %
INERT INGREDIENTS*	89.0 %
TOTAL	100.0%

*Includes 5% unsaturated fatty acids as a grooming aid.

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

For neck size up to 28".

FRANCODEX

**CONTAINS 1 FLEA AND TICK COLLAR
NET WT. 1.47 OZ. (41.5 g)**

BLISTER PACKAGE

BACK PANEL (CARDBOARD BACKING)

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: This product contains a cholinesterase inhibitor. Atropine sulfate and 2-PAM are antidotal only if signs of cholinesterase inhibition are present.

Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing dogs.

Adjustable - fits large dogs, 28" neck or smaller

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labeling.

Place Protection™ Plus 150 collar around dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 150 days. For unusually large breeds, collar aids in tick control in the head and neck area for up to 2 months. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

This Protection™ Plus 150 collar contains Essential Fatty Acids necessary for healthy skin and hair. Unsaturated fatty acids released from the collar are rapidly absorbed to help maintain a lustrous, manageable coat.

DISPOSAL

Do not reuse empty blister. Wrap blister and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 54022-xx-xxxxx

EPA Est. 15577-FR-1

Manufactured for Francodex Laboratories, Inc.
Subsidiary of Virbac, Inc
Fort Worth, Texas 76118
Made in France

FOIL POUCH/CARTON PACKAGE

FRONT PANEL (CARTON)

**PROTECTION™ Plus 150 Flea and Tick Collar for Large Dogs
with EFA*
(color)**

*Slow release coat conditioner helps enhance texture, manageability and luster.

**Kills fleas for 5 months
Kills ticks for 5 months**

For neck size up to 28".

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)

phosphorothioate] 11.0 %

INERT INGREDIENTS* 89.0 %

TOTAL 100.0%

*Includes 5% unsaturated fatty acids as a grooming aid.

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

FRANCODEX

**CONTAINS 1 FLEA AND TICK COLLAR
NET WT. 1.47 OZ. (41.5 g)**

FOIL POUCH/CARTON PACKAGE

BACK PANEL(CARTON)

**PROTECTION™ Plus 150 Flea and Tick Collar for Large Dogs with EFA (Color)
Adjustable - fits large dogs, 28" neck or smaller.**

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: This product contains a cholinesterase inhibitor. Atropine sulfate and 2-PAM are antidotal only if signs of cholinesterase inhibition are present.

Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing dogs.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labeling.

Place Protection™ Plus 150 collar around the dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 150 days. For unusually large breeds, collar aids in tick control in the head and neck area for up to 2 months. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

This Protection™ Plus 150 collar contains Essential Fatty Acids necessary for healthy skin and hair. Unsaturated fatty acids released from the collar are rapidly absorbed to help maintain a lustrous, manageable coat.

DISPOSAL

Do not reuse empty pouch or box. Wrap pouch or box and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 54022-7x-xxxxx

EPA Est. 15577-FR-1

**Manufactured for Francodex Laboratories, Inc.
Subsidiary of Virbac, Inc.
Fort Worth, Texas 76118
Made in France**

FOIL POUCH/CARTON PACKAGE

POUCH LABEL

**PROTECTION™ Plus 150 Flea and Tick Collar for Large Dogs
with EFA**

Do not open until ready to use

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

**SEE BACK PANEL OF OUTER CARTON FOR
ADDITIONAL PRECAUTIONARY STATEMENTS**

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-	
methyl-4-pyrimidinyl)	
phosphorothioate]	11.0 %
INERT INGREDIENTS*	89.0 %
TOTAL	100.0%

*Includes 5% unsaturated fatty acids as a grooming aid

EPA Reg. No. 54022-xx-xxxxx

EPA Est. 15577-FR-1

Manufactured for Franco-dex Laboratories, Inc.
Subsidiary of Virbac, Inc.
Fort Worth, Texas 75118
Made in France

**Protection™ Plus 150 Flea and Tick Collar for Dogs with EFA
(11% Diazinon)**

DRAFT LABELING

(5 Copies of Each Piece of Draft Labeling)

- Protection™ Plus 150 Flea and Tick Collar for Puppies/Virbac label
Blister Package
Pouch/Carton Package
- Protection™ Plus 150 Flea and Tick Collar for Dogs/Virbac label
Blister Package
Pouch/Carton Package
- Protection™ Plus 150 Flea and Tick Collar for Large Dogs/Virbac label
Blister Package
Pouch/Carton Package
- Protection™ Plus 150 Flea and Tick Collar for Puppies/Francodex (distributor) label
Blister Package
Pouch/Carton Package
- Protection™ Plus 150 Flea and Tick Collar for Dogs/Francodex (distributor) label
Blister Package
Pouch/Carton Package
- Protection™ Plus 150 Flea and Tick Collar for Large Dogs/Francodex (distributor) label
Blister Package
Pouch/Carton Package

**Virbac, Inc.
Fort Worth, Texas 76118**

BLISTER PACKAGE

FRONT PANEL (CARDBOARD BACKING)

**PROTECTION™ Plus 150 Flea and Tick Collar for Puppies
with EFA***

*Slow release coat conditioner helps enhance texture, manageability and luster.

**Kills fleas for 5 months
Kills ticks for 5 months**

(Color of collar visible through affixed blister package)

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-	
methyl-4-pyrimidinyl)	
phosphorothioate]	11.0 %
INERT INGREDIENTS*	89.0 %
TOTAL	100.0%

*Includes 5% unsaturated fatty acids as a grooming aid.

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

For puppies over 5 weeks - neck size up to 12".

VIRBAC

**CONTAINS 1 FLEA AND TICK COLLAR
NET WT. 0.71 OZ. (20.0 g)**

BLISTER PACKAGE

BACK PANEL (CARDBOARD BACKING)

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: This product contains a cholinesterase inhibitor. Atropine sulfate and 2-PAM are antidotal only if signs of cholinesterase inhibition are present.

Do not use other pesticides on puppy while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing puppies. Do not use on puppies less than 5 weeks of age.

Adjustable - will accommodate up to a 12" neck size.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labeling.

Place Protection™ Plus 150 collar around puppy's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 150 days. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

This Protection™ Plus 150 collar contains Essential Fatty Acids necessary for healthy skin and hair. Unsaturated fatty acids released from the collar are rapidly absorbed to help maintain a lustrous, manageable coat.

DISPOSAL

Do not reuse empty blister. Wrap blister and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 54022-

EPA Est. 15577-FR-1

Manufactured for Virbac, Inc.
Fort Worth, Texas 76118
Made in France

FOIL POUCH/CARTON PACKAGE

FRONT PANEL (CARTON)

**PROTECTION™ Plus 150 Flea and Tick Collar for Puppies
with EFA*
(color)**

*Slow release coat conditioner helps enhance texture, manageability and luster.

**Kills fleas for 5 months
Kills ticks for 5 months**

For neck size up to 12".

ACTIVE INGREDIENT

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)

phosphorothioate] 11.0 %

INERT INGREDIENTS* 89.0 %

TOTAL 100.0%

*Includes 5% unsaturated fatty acids as a grooming aid.

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

VIRBAC

CONTAINS 1 FLEA AND TICK COLLAR

NET WT. 0.71 OZ. (20.0 g)

FOIL POUCH/CARTON PACKAGE**BACK PANEL(CARTON)**

**PROTECTION™ Plus 150 Flea and Tick Collar for Puppies with EFA (Color)
Adjustable - will accomodate up to a 12" neck size.**

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: This product contains a cholinesterase inhibitor. Atropine sulfate and 2-PAM are antidotal only if signs of cholinesterase inhibition are present.

Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing puppies. Do not use on puppies less than 5 weeks of age.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labeling.

Place Protection™ Plus 150 collar around the puppy's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 150 days. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

This Protection™ Plus 150 collar contains Essential Fatty Acids necessary for healthy skin and hair. Unsaturated fatty acids released from the collar are rapidly absorbed to help maintain a lustrous, manageable coat.

DISPOSAL

Do not reuse empty pouch or box. Wrap pouch or box and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 54022-

EPA Est. 15577-FR-1

**Manufactured for Virbac, Inc.
Fort Worth, Texas 76118
Made in France**

FOIL POUCH/CARTON PACKAGE

POUCH LABEL

**PROTECTION™ Plus 150 Flea and Tick Collar for Puppies
with EFA**

Do not open until ready to use

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

**SEE BACK PANEL OF OUTER CARTON FOR
ADDITIONAL PRECAUTIONARY STATEMENTS**

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-	
methyl-4-pyrimidinyl)	
phosphorothioate]	11.0 %
INERT INGREDIENTS*	89.0 %
TOTAL	100.0%

*Includes 5% unsaturated fatty acids as a grooming aid

EPA Reg. No. 54022-

EPA Est. 15577-FR-1

Manufactured for Virbac, Inc.
Fort Worth, Texas 76118
Made in France