

Reg no: 54022-10

PM-15

1316

US ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDES PROGRAMS
REGISTRATION DIVISION (75-767)
WASHINGTON, DC 20480

EPA REGISTRATION NO.

DATE OF ISSUANCE

54022-10

JUL 26 1990

TERM OF ISSUANCE

Until Reregistration

NAME OF PESTICIDE PRODUCT

Protection 150 Flea and Tick
Collar for Dogs

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION

(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

┌

Wichita, KS
P.O. Box 1000
E. 10th St.

┐

160/249267
18/2

└

┘

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

The following conditions apply to the registration of this product:

1. The registrant shall submit to the Administrator a copy of the registration/reregistration of your product...

2. The registrant shall submit to the Administrator a copy of the labeling for the product for...

1. The registrant shall submit to the Administrator a copy of the following information: (a) A copy of the registration/reregistration of your product...

2. Subject to the conditions of registration, the registrant shall release the product for sale...

If these conditions are not met, the registration will be subject to cancellation in accordance with the Act. Your release for shipment of the product...

A stamp is required on the label of the product...

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

[Handwritten Signature]

DATE

JUL 26 1990

Richards

2916

BLISTER PACKAGE

FRONT PANEL (CARDBOARD BACKING)

PROTECTION™ 150 Flea and Tick Collar for Dogs

**Kills fleas for 5 months
Kills ticks for 4 months**

(Color of collar visible through affixed blister package)

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate]	15.0 %
INERT INGREDIENTS	85.0 %
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

For neck size up to 22".

FRANCODEX

**CONTAINS 1 FLEA AND TICK COLLAR
NET WT. 0.84 OZ. (23.7 g)**

541022-RN

ACCEPTED
with COMMENTS
in EPA Letter Dated:

Jul. 26 1990

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 541022-10

BLISTER PACKAGE**BACK PANEL(CARDBOARD BACKING)****PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: This product contains a cholinesterase inhibitor. Atropine sulfate and 2-PAM are antidotal only if signs of cholinesterase inhibition are present.

Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing animals.

Adjustable - fits dogs, 22" neck or smaller.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labeling.

Place Protection™ 150 collar around the dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 120 days. For unusually large breeds, aids in tick control in the head and neck area up to 2 months. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

DISPOSAL

Do not reuse empty blister. Wrap blister and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 54022-xx-xxxxx

EPA Est. 15577-FR-1

Manufactured for Francodex Laboratories, Inc.
Subsidiary of Virbac, Inc.
Fort Worth, Texas 76118
Made in France

FOIL POUCH/CARTON PACKAGE

FRONT PANEL (CARTON)

**PROTECTION™ 150 Flea and Tick Collar for Dogs
(color)**

**Kills fleas for 5 months
Kills ticks for 4 months**

For neck size up to 22"

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)

phosphorothioate]	15.0 %
INERT INGREDIENTS	85.0 %
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

FRANCODEX

**CONTAINS 1 FLEA AND TICK COLLAR
NET WT. 0.84 OZ. (23.7 g)**

FOIL POUCH/CARTON PACKAGE

BACK PANEL(CARTON)

**PROTECTION™ 150 Flea and Tick Collar for Dogs (Color)
Adjustable - fits dogs, 22" neck or smaller.**

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: This product contains a cholinesterase inhibitor. Atropine sulfate and 2-PAM are antidotal only if signs of cholinesterase inhibition are present.

Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing animals.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labeling.

Place Protection™ 150 collar around the dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 120 days. For unusually large breeds, aids in tick control in the head and neck area up to 2 months. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

DISPOSAL

Do not reuse empty pouch or box. Wrap pouch or box and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 54022-xx-xxxxx

EPA Est. 15577-FR-1

**Manufactured for Francodex Laboratories, Inc.
Subsidiary of Virbac, Inc.
Fort Worth, Texas 76118
Made in France**

FOIL POUCH/CARTON PACKAGE

POUCH LABEL

PROTECTION™ 150 Flea and Tick Collar for Dogs

Do not open until ready to use

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

**SEE BACK PANEL OF OUTER CARTON FOR
ADDITIONAL PRECAUTIONARY STATEMENTS**

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate]

.....	15.0 %
INERT INGREDIENTS	85.0 %
TOTAL	100.0%

EPA Reg. No. 54022-xx-xxxxx

EPA Est. 15577-FR-1

**Manufactured for Francodex Laboratories, Inc.
Subsidiary of Virbac, Inc.
Fort Worth, Texas 76118
Made in France**

7 7 16

BLISTER PACKAGE

FRONT PANEL (CARDBOARD BACKING)

PROTECTION™ 150 Flea and Tick Collar for Large Dogs

**Kills fleas for 5 months
Kills ticks for 2 months**

(Color of collar visible through affixed blister package)

ACTIVE INGREDIENT:	
Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate]	15.0 %
INERT INGREDIENTS	<u>85.0 %</u>
TOTAL	<u>100.0%</u>

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

For neck size up to 28".

FRANCODEX

**CONTAINS 1 FLEA AND TICK COLLAR
NET WT. 0.9 OZ. (25.5 g)**

BLISTER PACKAGE**BACK PANEL(CARDBOARD BACKING)****PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: This product contains a cholinesterase inhibitor. Atropine sulfate and 2-PAM are antidotal only if signs of cholinesterase inhibition are present.

Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing animals.

Adjustable - fits large dogs, 28" neck or smaller.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labeling.

Place Protection™ 150 collar around the dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue to aid in tick control in the head and neck area in unusually large breeds for up to 2 months. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

DISPOSAL

Do not reuse empty blister. Wrap blister and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 54022-xx-xxxxx

EPA Est. 15577-FR-1

Manufactured for Francodex Laboratories, Inc.
Subsidiary of Virbac, Inc.
Fort Worth, Texas 76118
Made in France

FOIL POUCH/CARTON PACKAGE

FRONT PANEL (CARTON)

**PROTECTION™ 150 Flea and Tick Collar for Large Dogs
(color)**

**Kills fleas for 5 months
Kills ticks for 2 months**

For neck size up to 28".

ACTIVE INGREDIENT:	
Diazinon [O,O-diethyl O-(2-isopropyl-6-	
methyl-4-pyrimidinyl)	
phosphorothioate]	15.0 %
INERT INGREDIENTS	85.0 %
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

FRANCODEX

**CONTAINS 1 FLEA AND TICK COLLAR
NET WT. 0.9 OZ. (25.5 g)**

FOIL POUCH/CARTON PACKAGE

BACK PANEL(CARTON)

**PROTECTION™ 150 Flea and Tick Collar for Large Dogs (Color)
Adjustable - fits large dogs, 28" neck or smaller.**

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: This product contains a cholinesterase inhibitor. Atropine sulfate and 2-PAM are antidotal only if signs of cholinesterase inhibition are present.

Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing animals.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labeling.

Place Protection™ 150 collar around the dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue to aid in tick control in the head and neck area in unusually large breeds for up to 2 months. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

DISPOSAL

Do not reuse empty pouch or box. Wrap pouch or box and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 54022-xx-xxxxx

EPA Est. 15577-FR-1

**Manufactured for Francodex Laboratories, Inc.
Subsidiary of Virbac, inc.
Fort Worth, Texas 76118
Made in France**

FOIL POUCH/CARTON PACKAGE

POUCH LABEL

PROTECTION™ 150 Flea and Tick Collar for Large Dogs

Do not open until ready to use

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

**SEE BACK PANEL OF OUTER CARTON FOR
ADDITIONAL PRECAUTIONARY STATEMENTS**

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate]	15.0 %
INERT INGREDIENTS	85.0 %
TOTAL	100.0%

EPA Reg. No. 54022-xx-xxxxx

EPA Est. 15577-FR-1

**Manufactured for Francodex Laboratories, Inc.
Subsidiary of Virbac, Inc.
Fort Worth, Texas 76118
Made in France**

BLISTER PACKAGE

FRONT PANEL (CARDBOARD BACKING)

PROTECTION™ 150 Flea and Tick Collar for Puppies

**Kills fleas for 5 months
Kills ticks for 4 months**

(Color of collar visible through affixed blister package)

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate]	
.....	15.0 %
INERT INGREDIENTS	85.0 %
.....	85.0 %
TOTAL	100.0%
.....	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

For puppies over 5 weeks - neck size up to 12".

VIRBAC

**CONTAINS 1 FLEA AND TICK COLLAR
NET WT. 0.46 OZ. (13.0 g)**

BLISTER PACKAGE

BACK PANEL(CARDBOARD BACKING)

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: This product contains a cholinesterase inhibitor. Atropine sulfate and 2-PAM are antidotal only if signs of cholinesterase inhibition are present.

Do not use other pesticides on puppy while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing puppies. Do not use on puppies less than 5 weeks of age.

Adjustable - will accomodate up to a 12" neck size.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labeling.

Place Protection™ 150 collar around the puppy's neck ,adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 120 days. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

DISPOSAL

Do not reuse empty blister. Wrap blister and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 54022-

EPA Est. 15577-FR-1

Manufactured for Virbac, Inc.
Fort Worth, Texas 76118
Made in France

FOIL POUCH/CARTON PACKAGE

FRONT PANEL (CARTON)

**PROTECTION™ 150 Flea and Tick Collar for Puppies
(color)**

**Kills fleas for 5 months
Kills ticks for 4 months**

For puppies over 5 weeks - neck size up to 12".

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)

phosphorothioate]	15.0 %
INERT INGREDIENTS	85.0 %
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

VIRBAC

**CONTAINS 1 FLEA AND TICK COLLAR
NET WT. 0.46 OZ. (13.0 g)**

FOIL POUCH/CARTON PACKAGE

BACK PANEL(CARTON)

**PROTECTION™ 150 Flea and Tick Collar for Puppies (Color)
Adjustable - will accomodate up to a 12" neck size.**

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: This product contains a cholinesterase inhibitor. Atropine sulfate and 2-PAM are anti-tal only if signs of cholinesterase inhibition are present.

Do not use other pesticides on puppy while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing puppies. Do not use on puppies less than 5 weeks of age.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labeling.

Place Protection™ 150 collar around the puppy's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 120 days. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

DISPOSAL

Do not reuse empty pouch or box. Wrap pouch or box and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 54022-

EPA Est. 15577-FR-1

**Manufactured for Virbac, Inc.
Fort Worth, Texas 76118
Made in France**

16 g 16

FOIL POUCH/CARTON PACKAGE

POUCH LABEL

PROTECTION™ 150 Flea and Tick Collar for Puppies

Do not open until ready to use

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

**SEE BACK PANEL OF OUTER CARTON FOR
ADDITIONAL PRECAUTIONARY STATEMENTS**

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate]	15.0 %
INERT INGREDIENTS	85.0 %
TOTAL	100.0%

EPA Reg. No. 54022-

EPA Est. 15577-FR-1

**Manufactured for Virbac, Inc.
Fort Worth, Texas 76118
Made in France**

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION
(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

NAME OF PESTICIDE PRODUCT
Protection 150 Flea and Tick
Collar for Dogs

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Virbac, Inc.
P.O. Box 16598
Ft. Worth, TX 76161

160-249267
10/2

BEST AVAILABLE COPY

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is hereby registered/re-registered in accordance with the Act.

1. Submit and/or amend your labeling to comply with the requirements for registration of your product under FIFRA, 40 CFR, and the Act. The labeling must be submitted to the Administrator for approval.

2. The labeling must be submitted to the Administrator for approval. The labeling must be submitted to the Administrator for approval.

3. The Note to Registrant should read as follows:
Note to Registrant: Diazinon is an organophosphorus insecticide. Its symptoms of cholinesterase inhibition are present, but are sufficiently mild to be considered. 2-PAM is also present and may be used for only in certain conditions.

3. Submit five (5) copies of your final labeling for you release the product for shipment. Refer to the Act for a further description of final print labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of this label is enclosed for your records.

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL
[Signature]

DATE
July 26, 1990

Richards

PROTECTION™ 150 Flea and Tick Collar for Dogs

**Kills fleas for 5 months
Kills ticks for 4 months**

(Color of collar visible through affixed blister package)

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)

phosphorothioate] 15.0 %

INERT INGREDIENTS 85.0 %

TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

For neck size up to 22".

FRANCODEX

**CONTAINS 1 FLEA AND TICK COLLAR
NET WT. 0.84 OZ. (23.7 g)**

521022-02N

ACCEPTED
with COMMENTS
in EPA Letter Dated:

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
amended, for the purpose of
enforcing the EPA Act, the

521022-10

BLISTER PACKAGE

BACK PANEL(CARDBOARD BACKING)

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: This product contains a cholinesterase inhibitor. Atropine sulfate and 2-PAM are antidotal only if signs of cholinesterase inhibition are present.

Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing animals.

Adjustable - fits dogs, 22" neck or smaller.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labeling.

Place Protection™ 150 collar around the dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 120 days. For unusually large breeds, aids in tick control in the head and neck area up to 2 months. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

DISPOSAL

Do not reuse empty blister. Wrap blister and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 54022-xx-xxxxx

EPA Est. 15577-FR-1

Manufactured for Francodex Laboratories, Inc.
Subsidiary of Virbac, Inc.
Fort Worth, Texas 76118
Made in France

FOIL POUCH/CARTON PACKAGE

FRONT PANEL (CARTON)

**PROTECTION™ 150 Flea and Tick Collar for Dogs
(color)**

**Kills fleas for 5 months
Kills ticks for 4 months**

For neck size up to 22".

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-
methyl-4-pyrimidinyl)

phosphorothioate]	15.0 %
INERT INGREDIENTS	85.0 %
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

FRANCODEX

**CONTAINS 1 FLEA AND TICK COLLAR
NET WT. 0.84 OZ. (23.7 g)**

FOIL POUCH/CARTON PACKAGE

BACK PANEL(CARTON)

**PROTECTION™ 150 Flea and Tick Collar for Dogs (Color)
Adjustable - fits dogs, 22" neck or smaller.**

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: This product contains a cholinesterase inhibitor. Atropine sulfate and 2-PAM are antidotal only if signs of cholinesterase inhibition are present.

Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing animals.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labeling.

Place Protection™ 150 collar around the dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 120 days. For unusually large breeds, aids in tick control in the head and neck area up to 2 months. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

DISPOSAL

Do not reuse empty pouch or box. Wrap pouch or box and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 54022-xx-xxxxx

EPA Est. 15577-FR-1

**Manufactured for Francodex Laboratories, Inc.
Subsidiary of Virbac, Inc.
Fort Worth, Texas 76118
Made in France**

FOIL POUCH/CARTON PACKAGE

POUCH LABEL

PROTECTION™ 150 Flea and Tick Collar for Dogs

Do not open until ready to use

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

**SEE BACK PANEL OF OUTER CARTON FOR
ADDITIONAL PRECAUTIONARY STATEMENTS**

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate]	15.0 %
INERT INGREDIENTS	85.0 %
TOTAL	100.0%

EPA Reg. No. 54022-xx-xxxxx

EPA Est. 15577-FR-1

**Manufactured for Francodex Laboratories, Inc.
Subsidiary of Virbac, Inc.
Fort Worth, Texas 76118
Made in France**