

PM 33

54289-3

9/30/98

1/8



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (H7505C)  
401 "M" St., S.W.  
Washington, D.C. 20460

EPA Reg. Number:

54289-3

Date of Issuance:

SEP 30 1998

**NOTICE OF PESTICIDE:**

Registration  
 Reregistration

(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Peraclean 5%  
(Peroxyacetic Acid Solution)

Name and Address of Registrant (include ZIP Code):

Degussa Corporation  
65 Challenger Road  
Ridgefield Park, N.J. 07660

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

In the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label changes:

a. Revise the EPA Registration Number to read, "EPA Reg. No. "54289-3".

Signature of Approving Official:

Marshall Swindell  
Product Manager 33  
Regulatory Management Branch 1  
Antimicrobial Division (7510W)

Date:

SEP 30 1998

Page 2  
EPA Registration No. 54289-3

b. On the second page of your label, under the heading "Sanitizing Food Contact Surfaces", change the sentence "All surfaces should be exposed to the sanitizing solution for a period of at least 60 seconds unless otherwise specified by governing sanitary code" to read "All surfaces should be exposed to the sanitizing solution for a period of at least 60 seconds or more if specified by governing sanitary code".

3. Submit two copies of the revised final printed label for the records.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

If you have any questions concerning this letter, please contact Martha Terry at 703-305-6217.

Sincerely,

*Martha S Terry*  
Marshall Swindell  
Product Manager 33  
Regulatory Management Branch 1  
Antimicrobial Division (7510W)

Enclosure

## Peraclean™ 5% (Peroxyacetic Acid Solution)

Peraclean™ 5% is a peroxyacetic acid-based sanitizer/disinfectant developed for the following uses:

Institutional/Industrial Sanitizing  
of Previously Cleaned Non-  
Porous Food Contact Surfaces in:

- ◆ Dairies
- ◆ Wineries
- ◆ Breweries
- ◆ Food and Beverage Plants
- ◆ Disinfecting Poultry Premises
- ◆ Poultry Hatcheries
- ◆ Animal Housing Facilities

Hard Surface Disinfection in:

- ◆ Hospitals
- ◆ Health Care Facilities
- ◆ Schools
- ◆ Colleges
- ◆ Veterinary Clinics
- ◆ Animal Life Science  
Laboratories
- ◆ Industrial Facilities
- ◆ Office Buildings
- ◆ Recreational Facilities
- ◆ Retail and Wholesale  
Establishments

Bacteria, Fungi, and Slime  
Control in

- ◆ Pulp and Paper Mill Systems
- ◆ Dispersed Pigments
- ◆ Cooling Water Systems
- ◆ Coatings Preservation

Active Ingredients:

Hydrogen Peroxide	26.5%
Peroxyacetic Acid	2.5%

Inert Ingredients:	68.5%
--------------------	-------

Total:	100%
--------	------

EPA Reg. No. 54289-  
EPA Estab. No.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

SEP 30 1998

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended for the pesticide,  
registered under EPA Reg. No.

54289-3

Before Using This Product, Please Read This Entire Label Carefully.

**KEEP OUT OF REACH OF CHILDREN**

**DANGER**

### STATEMENT OF PRACTICAL TREATMENT

If in eyes: Hold eyelids open and flush with a steady, gentle stream of water for 5 minutes. Get medical attention.

If on skin: Wash with plenty of soap and water. Get medical attention.

If swallowed: Call a doctor or get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

Manufactured by:

Degussa Corporation  
65 Challenger Road  
Ridgefield, NJ 07660

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER. CORROSIVE.** Causes irreversible eye damage and skin burns. May be fatal if inhaled or absorbed through the skin. Harmful if swallowed. Do not breathe vapors or spray mist. Do not get in eyes on skin or on clothing. Wear goggles and/or face shield and rubber gloves when handling. Do not enter an enclosed area without proper respiratory protection. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse

**PHYSICAL AND CHEMICAL HAZARDS**

**STRONG OXIDIZING AGENT. Corrosive.** Mix only with water. Product must be diluted in accordance with label directions prior to use. Peraclean™ 5% is not combustible; however, at temperatures exceeding 156°F, decomposition occurs releasing oxygen. The oxygen released could initiate combustion.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to birds, fish, and aquatic invertebrates. Caution should be used when applying indoors because pets may be at risk. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the US Environmental Protection Agency.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**SANITIZATION**

Peraclean™ 5% peroxyacetic acid sanitizer is recommended for use on precleaned surfaces such as equipment, pipelines, tanks, vats, fillers, evaporators, pasteurizers and aseptic equipment in dairies, breweries, wineries, beverage and food processing/packing plants, egg processing/packing equipment surfaces, and eating establishments. This product is effective as a sanitizer when solution is prepared in water of up to 400 ppm hardness as CaCO<sub>3</sub>. This product has demonstrated greater than a 99.999% reduction of survivors after a 30 second exposure period in the AOAC Germicidal and Detergent Sanitizing Action of Disinfectants study.

**Sanitizing Food Contact Surfaces**

Effective against *Staphylococcus aureus* and *Escherichia coli*.

Prior to sanitizing, remove gross food particles, then wash with a detergent solution, followed by a potable water rinse. Sanitize with a concentration of 1.0 ounce Peraclean™ 5% dissolved in 5 gallons of water (0.16% v/v concentration). At this dilution Peraclean™ 5% is effective against *Staphylococcus aureus* and *Escherichia coli*. Use immersion, coarse spray or circulation techniques as appropriate to the equipment. All surfaces should be exposed to the sanitizing solution for a period of at least 60 seconds unless otherwise specified by governing sanitary code. Drain thoroughly and allow to air dry. Do not rinse.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

SEP 30 1998

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide,  
registered under EPA Reg. No.  
5-4289-3

Sanitizing, Eating, Drinking, And Food Prep Utensils

Remove gross food particles by a prescrape, a preflush and, when necessary, a presoak treatment.

Wash with a recommended detergent.

Rinse with clean water.

Sanitize in a solution of 1.0 ounce Peraclean™ 5% dissolved in 5 gallons of water. Immerse all utensils for at least 60 seconds or contact time specified by governing sanitary code.

Drain and air dry.

Sanitizing Tableware

For sanitizing tableware in low temperature warewashing machines, inject Peraclean™ 5% into the final rinse water at a concentration of 1.0 ounce Peraclean™ 5% dissolved in 5 gallons of water. Do not exceed 0.16 % v/v. Air dry.

To insure that the Peraclean™ 5% sanitizer concentration does not fall below 0.1%, periodically test the rinse solution with a suitable test kit and adjust the dispensing rate accordingly. Consult your technical service representative for assistance and further information on sanitizing tableware in warewashing machines.

Final Sanitizing Bottle Rinse

Peraclean™ 5% may be used as a final sanitizing rinse for returnable and non-returnable bottles at a 0.16% dilution (1.0 ounce Peraclean™ 5% dissolved in 5 gallons of water).

**NOTE: FOR MECHANICAL OPERATIONS** prepared use solution may not be reused for sanitizing but may be reused for other purposes such as cleaning.

**FOR MANUAL OPERATIONS** fresh sanitizing solutions should be prepared at least daily or more often if the solution becomes diluted or soiled.

HARD SURFACE DISINFECTION

Peraclean™ 5% disinfects as it cleans in one operation. Peraclean™ 5% can be used to disinfect floors, walls and other hard nonporous surfaces such as tables, chairs, countertops, bathroom fixtures, sinks, bed frames, shelves, racks, carts, refrigerators, coolers, tile, linoleum, vinyl, asphalt, porcelain, plastic (such as polypropylene and polyethylene), stainless steel, or glass.

Areas of use in hospitals, use Peraclean™ 5% for surgical and obstetrical suites; housekeeping services; physical therapy departments; nursing services; autopsy facilities. Also, use Peraclean™ 5% in nursing homes, other health-care facilities, schools, colleges, veterinary clinics, animal life science laboratories, industrial facilities, dietary areas, office buildings, recreational facilities, retail and wholesale establishments.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

SEP 30 1998

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide,  
registered under EPA Reg. No.

Combination Disinfection and Cleaning

Peraclean™ 5% is effective against *Staphylococcus aureus*, *Salmonella choleraesuis*, *Pseudomonas aeruginosa*, *Trichophyton mentagrophytes*, and *Escherichia coli* 0157:H7 at 0.23% (1.5 oz./5 gal.) in hard water (400 ppm as CaCO<sub>3</sub>) and 5% fetal bovine serum on hard nonporous surfaces. For heavily soiled areas a pre-cleaning step is required. Apply solution with mop, cloth, sponge, brush, scrubber, or coarse spray device, or by soaking so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes, then remove solution and entrapped soil with a clean wet mop, cloth, or wet vacuum pickup. Prepare a fresh solution daily or when it becomes soiled or diluted.

Peraclean™ 5% is designed for use in animal hospitals, animal laboratories, kennels, pet shops, zoos, pet animal quarters, poultry premises, poultry hatcheries, and livestock quarters. When used as directed, Peraclean™ 5% is specifically designed to disinfect, deodorize and clean inanimate, hard, surfaces such as walls, floors, sink tops, furniture, operating tables, kennel runs, cages and feeding and watering equipment. In addition Peraclean™ 5% will deodorize those areas which are generally hard to keep smelling fresh such as, garbage storage areas, empty garbage bins and cans, and any other areas which are prone to odors caused by microorganisms.

All treated equipment that will contact food, feed, or drinking water must be rinsed with potable water before reuse.

For heavily soiled areas, a pre-cleaning step is required. Prepare a fresh solution for each use.

BIOFOULING CONTROL IN PULP AND PAPER MILL SYSTEMS

For use in the manufacture of paper and paperboard intended for non-food contact.

Peraclean™ 5% can be used to control bacteria, fungi, and fresh water organisms in paper, paperboard, or nonwoven process water and influent water systems. Suitable dosing points include but are not limited to: stock chests, pulpers, the white water loop and white water storage systems and influent water streams.

Influent Water Systems:

Peraclean™ 5% should be fed continuously to incoming fresh water streams (nonpotable use only) at dosages ranging from 5-500 ppm.

Mill Process Waters:

- **Continuous Feed** – Peraclean™ 5% should be fed continuously at dosages ranging from 5-500 ppm. This range may be equivalent to 0.4 – 40 lbs. Peraclean™ 5% per ton (dry basis) of pulp or paper produced.
- **Intermittent Feed** – Peraclean™ 5% should be fed intermittently (6-8 times per day) at dosages ranging from 5-500 ppm. This dosage may be equivalent to 0.4-40 lbs. Peraclean™ 5% per ton (dry basis) of pulp or paper produced during the feed period.
- **Shock Dose** – Peraclean™ 5% should be shock dosed at dosages ranging from 50-1000 ppm. This dosage may be equivalent to 4-82 lbs. Peraclean™ 5% per ton (dry basis) of pulp or paper produced during the feed period.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

SEP 30 1998  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide,  
registered under EPA Reg. No.  
5-4289-3

CONTROL OF BACTERIA AND FUNGI IN DISPERSED PIGMENTS

Peraclean™ 5% can be used in the control of bacteria and fungi in the manufacture and storage of dispersed pigments such as kaolin clay, titanium dioxide, calcium carbonate, calcium sulfate, barium sulfate, magnesium silicate and diatomaceous earth used in paint and paper production.

Add 0.3 to 1.5 lbs. (5 to 24 fl. oz.) of Peraclean™ 5% to each 1000 lbs. of pigment slurry. This will provide 300 to 1500 ppm of product.

CONTROL OF SLIME FORMING BACTERIA TO COOLING WATER SYSTEMS (COOLING TOWERS, EVAPORATIVE CONDENSERS)

- Severely fouled systems should be cleaned before adding the Peraclean™ 5% solution. Peraclean™ 5% should be added in the system directly and not mixed with any other chemicals or additives. Contamination with other chemicals could result in product decomposition.
- Add Peraclean™ 5% at a point in the system where uniform mixing and even distribution will occur.
- Use 0.3 to 1.5 lb (5 to 24 fl. oz) of Peraclean™ 5% per 1000 gallons of solution as a continuous or intermittent slug treatment. This will provide 300 – 1500 ppm of product. Repeat treatment as required to maintain control.

CONTROL OF BACTERIA AND FUNGI IN COATING PRESERVATION

Not for use in manufacture of material intended for food contact.

Peraclean™ 5% can be used as an in-container preservative for the control of bacteria and fungi in water-based coatings such as paper coatings.

Add 0.2 to 1.5 lbs. (5 to 24 fl oz) of Peraclean™ 5% to each 1000 lbs. of water. This will provide 200 to 1400 ppm of product.

DISINFECTION OF POULTRY PREMISES, TRUCKS, COOPS AND CRATES

Remove all poultry and feeds from premises, trucks, coops and crates.

Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by poultry.

Empty all troughs, racks and other feeding and watering appliances.

Thoroughly clean all surfaces with a detergent and rinse with water.

Saturate surfaces with a 0.23% (1.5 oz./5 gal.) solution of Peraclean™ 5% for a period of 10 minutes.

Ventilate buildings, coops and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set or dried.

Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with a detergent and rinse with potable water before reuse.

See your technical representative for specific recommendations for all cleaning and rinsing requirements.

Poultry Hatchery Disinfection

Clean out any remaining eggs and chicks. Remove gross soils, such as litter, down, shell fragments or other hatching related debris. Empty all racks and other equipment. Thoroughly wash all surfaces, including floors, walls, conveyors, trays and water systems with a recommended detergent. Rinse thoroughly with water. Apply a 0.23% (1.5 oz./5 gal.) solution of Peraclean™ 5% with a mop, cloth, brush or coarse spray. Wet all surfaces and allow to remain wet for 10 minutes. Ventilate buildings and other closed spaces. Allow to dry before reintroducing eggs.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

SEP 30 1998

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide,  
registered under EPA Reg. No.  
54289-3

Disinfection And Deodorizing Of Animal Housing Facilities (Barns, Kennels, Hutches, Etc.)

Remove animals and feed from facilities. Remove litter, waste matter, and gross soils. Empty all troughs, rack and other feeding and watering equipment. Wash surfaces with a recommended alkaline detergent, by manual, foam, or spray application. Rinse with water. Apply a 0.23% (1.5oz/5 gal.) solution of Peraclean™ 5% with a mop, cloth, brush or coarse spray. Wet all surfaces and allow to remain wet for 10 minutes. Ventilate buildings and other closed spaces. Allow to air dry before reintroducing animals.

For Sanitizing Of Hatching Eggs

Prepare a solution of Peraclean™ 5% by diluting 1.0 ounce Peraclean™ 5% with 5 gallons of water. As eggs are gathered or prior to setting, apply solution as a coarse spray so as to lightly wet all shell surfaces.

NOTE: This product at its use dilution is compatible with stainless steel and aluminum surfaces. If product is intended to be used on any other surface, it is recommended that you apply product to a smaller test area to determine compatibility before proceeding with its use.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Never return Peraclean™ 5% to the original container after it has been removed. Avoid all contaminants, especially dirt, caustic, reducing agents and metals. Contamination and impurities will reduce shelf life and can induce decomposition. In case of decomposition, isolate container, douse container with cool water and dilute Peraclean™ 5% with large volumes of water. Avoid damage to containers. Keep container closed at all times when not in use. Keep container out of direct sunlight. To maintain product quality, store at temperatures below 86°F. Do not store on wooden pallets.

Procedures for Leak or Spill: Stop leak if this can be done without risk. Shut off ignition sources; no flames, smoking, flares or spark-producing tools. Keep combustible and organic materials away. Flush spilled material with large quantities of water. Undiluted material should not enter confined spaces.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. If material has been spilled, an acceptable method of disposal is to dilute with at least 20 volumes of water followed by discharge into suitable treatment system in accordance with all local, state, and Federal environmental laws, rules, regulations, standards, and other requirements. Because acceptable methods of disposal may vary by location, regulatory agencies should be contacted by prior to disposal. Peraclean™ 5%, which is to be discarded, should be disposed of as hazardous waste after contacting the appropriate local, state, or Federal agency to determine proper procedures.

Container Disposal: > 5 gallon plastic drums: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

>5 gallons plastic tote bin liners: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

SEP 30 1998

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide  
registered under EPA Reg. No.

54289-3