

PM21 54401-1 10/4
UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

Ms. Linda C. Watson
Consultant/Regulatory Affairs
Lykes AGRI Sales, Inc.
P.O. Box 1758
Dade City, FL 33526-1758

NOV 5 1992

Dear Ms. Watson:

Subject: Extra Value Dusting Sulfur
EPA Registration No. 54401-1
Your Submission Dated April 30, 1992

The amendment referred to above, submitted in connection with registration under section 3(c) (7) (A) of the Federal Insecticide, Fungicide, and Rodenticide Act, (FIFRA) is acceptable provided you:

1. Make the following label change before you release the product for shipment bearing the amended labeling:

At the next label printing, change the first sentence under Environmental Hazards to read "Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark".

2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Susan T. Lewis
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

H7505C/CDAVIS/cd/11-5-92/Room 265/305-6700

CONCURRENCES							
SYMBOL	H7525C						
SURNAME	DAVIS						
DATE	11/5/92						

PH 21

54401-1

2014



LYKES

AGRI SALES, INC.



S U N G R O

DUSTING SULFUR

50 Lbs. Net Weight

ACTIVE INGREDIENTS:

Sulfur 93.0%

INERT INGREDIENTS: 7.0%

100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENTS OF PRACTICAL TREATMENT

If on skin: Wash with plenty of soap and water.
Get medical attention.

If in Eyes: Flush with water for at least 15 minutes and get
medical attention.

See side panel for additional precautionary statements.

Manufactured By

LYKES AGRI SALES, INC.

P.O. Box 1685

Eaton Park, Florida 33840

EPA Reg. No. 54401-1

EPA Est No. 54401-FL-002

A-2507

ACCEPTED
with COMMENTS
in EPA Letter Dated

NOV 05 1992

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
amended, for the pesticide
registered under EPA Reg. No.

54401-1

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

CAUTION

Harmful if inhaled or absorbed through skin. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before re-use.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from treated areas.

PHYSICAL AND CHEMICAL HAZARDS

Sulfur dust suspended in air ignites easily. Keep away from heat, sparks, and flame. Do not smoke while applying this product.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT: Do not re-enter treated crop area prior to 24 hours after application.

GENERAL WORKER PROTECTION STATEMENTS: Do not apply this product in such a manner as to directly, or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warning must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. (Indicate specific oral warnings which inform workers of areas or fields that may not be entered without specific protective clothings, period of time field must be vacated and appropriate actions to take in case of accidental exposure.) When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written warnings must include the following information: "CAUTION" "Area treated with SULFUR on (date of application). Do not enter without appropriate protective clothing for 24 hours. In case of accidental exposure wash with plenty of soap and water, flush eyes with water for at least 15 minutes, get medical attention."

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

STORAGE: Store in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site, or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration; or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

4074

**OBSERVE THE FOLLOWING
USE PRECAUTIONS**

Sulfur may cause severe fruit and foliage injury to certain crops. Use only if local use has proved that sulfur does not damage crops in your area. Sulfur may burn foliage when temperature is high. Do not apply if temperature during, or within 3 days after application is expected to exceed 90° F.

When growing crops for processing, consult the processor before applying sulfur.

Do not use sulfur within 3 to 4 weeks of an oil spray.

Atomization Prohibition

Do not apply this product through any type of irrigation system.

APPLES: Scab, Powdery mildew. Apply 30 to 40 lbs. per acre. Make first application in Green Tip Stage. Repeat application in Pink Bud Stage. Use low rate on smaller trees, increasing rate according to size of tree. Use high rate on larger trees. Use on sulfur tolerant varieties only.

CITRUS: Six Spotted and citrus rust mites. Apply 80 lbs. per acre; (½ to 1½ lbs. per tree, depending on size, for citrus rust mite in Florida). Citrus flat and Yuma spider mites. Apply 100 lbs. per acre; Citrus thrips - Apply 100 lbs. per acre in three applications; mid-March, Petal fall and 2-4 weeks later. Do not apply within 3 to 6 weeks of an oil spray.

PEACH, NECTARINE: Brown rot blossom blight, Fruit brown rot, Coryneum blight, Rust, Peach silver mite, Powdery mildew. Apply 50 lbs. per acre at Blossom, Petal-fall, Shuck-fall, and four weeks before harvest.

BEANS: Powdery mildew, Rust, Spider mites. Apply 15-30 lbs. per acre. For Rust control, start applications as soon as the first true leaves form and continue at weekly intervals until blossom time.

CAUTION: May cause injury to some varieties of beans. Use low rate when plants are small; increase rate as plants increase in size. Use high rate on larger plants. Use high rate when infection is severe.

COTTON: Atlantic mite, Cotton rust. Apply 25 to 30 lbs. per acre. For Rust control, start applications as soon as the first true leaves form and continue at weekly intervals until blossom time. Use low rate when plants are small; increase rate as plants increase in size. Use high rate on larger plants. Use high rate when infection is severe.

PEAS: Powdery mildew. Apply 15 to 20 lbs. per acre. Begin application at first bloom or first disease appearance. Continue applications at weekly intervals until 4 days prior to harvest. Use the low rate when plants are small; increasing to the high rate as plants grow larger. Use the high rate during cool, humid weather or following application of irrigation water. Use the high rate if infection is severe.

PEANUTS: Rust, Leaf spot. Apply 15 to 20 lbs. per acre. Begin applications at first appearance of disease. Continue applications at 14 day intervals, until 4 days prior to harvest. Use the low rate when plants are small, increasing to the high rate when plants are 35 days old. Use the high rate if infection is severe.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.