54705-5

11-29-2002

	1)	5
	2-21	
1	Z: Z:	_ 11

Hose reed instructions on	, reverse <b>kal</b> are comoleti	ina form.		Form Apr	Nove	. OMB No.	2070-0	OSO. Appro	val sopires 2	-23-95
	Un	nited States	<u> </u>			Registra	ation	OPP Ide	entifier Numb	
<b>⊕EPA</b>	Environmental Washin	Protection Angton, DC 20460	/gency	l		Amenda Other	ment	;		
		Application for	or Pastic	ida - Saci	للللا	,				
1. Company/Product Number		thousand.	- T	A Product Mane			3.	, Proposed C	lessification	
54705-5			1	ne I. Miller			l r	None	Restri	inted
4. Company/Product (Name WEED STOPPER	)		PM# PM-2	23	~					
5. Name and Address of Ap	xplicant (Include ZIP Cod	je)		pedited Rev						-
Lawn and Garden Pr	•	4.°	(b)(i), to:	my product is	is sim	illar or ident	tical in	compositio	n and labeli	ing
c/o Lynne Zahigian F 6650 Wade Lane S	•	~	1	Reg. No		And the latest the lat				<del></del>
	is is a new address	The C. C.	Prod	luct Name						
			Section -							
Amendment - Explei	in below.	K ar man marro	1			ls in repsons	e to			
	ponse to Agency letter d	dated		Agency letter "Me Too" A	ter dat	ted		NOTIFIC/	TION	
Notification - Explain		-	,	Other - Expl	••		i	NOV 2 9	2002	
	**************************************	- 1		<u> </u>			<del></del>			
Explanation: Use addition					<del></del>	Idanna ji	- 00 Nr	···· 2001_R	29 40 and th	
Notification of label changes requirements of EPA regular	itions at 40 CFR 156.10 ar	ind 40 CFR 152.46,	and no other	changes have l	been i	made to the k	labeling d	or the confide	ential stateme	ent of
formula of this product. I und notification is not consistent	derstand that it is a violation	ion of 18 U.S.C. Sec	c. 1001 to will	lfully make any l	false s	statement to E	EPA. I fu	urther underst	tand that if this	is
violation of FIFRA and I may							<b>U</b>	Action and	Address	
		\$	Section -	111						
1. Material This Product Wi	A Be Packaged In:									
Child-Resistant Peckaging	Unit Packaging	w	/ater Solubie	Packaging		2. Type of				
Yes	Yes		Yes		_		Mete Plest			i
No	No	L	No			1	Glass	18	•	
* Certification must	if "Yes" Unit Packaging wgt.	No. per lf container Pa	Yes Wat	No. per container	r		Pepel Other	or or (Specify)		
be submitted	1									
3. Location of Net Contents		4. Size(s) Retail Co	ontainer	ſ	5; Lo	cation of Lat	pel Direc	ctions		
	Container  Affixed to Product	1,104,							<del></del>	
6. Manner in Whiten Label is	Affixed to Product	Lithograph Paper glued Stendied	1	Other	′ —					
			ection -	IV				ର ଓ ନମ୍ୟର୍ଶର ମ	·	
1. Contact Point   (Complete	items directly below fo				if nec	essery, to pi	rocess t	this epilostic	on.)	
Name		Title					Teleph	hone No. (Inc	lude Area Co	ode)
Lynne C. Zahigian		Age	int			Å.	775/62	29-1408		
	_	Certification				_		1 0.000	Application	
	ements i have made on t ny knowlingliy false or m i law.							-	itamped)	
2. Signature		3. Tit	<del></del>				<del></del>	22 4 0 p		
	Zahigian	Age								
4. Typed Name		5. Da	ite , j				-			
Lynne C. Zahigian		1	1/13/	02						



### **NOTIFICATION**

#### **APPLICATION INSTRUCTIONS**

SOIL PREPARATION

NOV 2 9 2002 WEED STOPPER controls weeds growing from seed. WEED STOPPER will not control control established weeds, weeds growing from stolens, rhizomes, or root pieces. Therefore, areas to be treated should be free of emerged weeds. Weed residues, pruning and trash should be thoroughly mixed into the soil or removed prior to treatment.

#### Handheld or Backpack Sprayer Application

The amount of water used to apply WEED STOPPER herbicide is not critical, but should be sufficient for uniform coverage of the target area. Calibrate by determining the volume of water required to treat 1,000 square feet. Use this calibration volume to determine the amount of water and WEED STOPPER herbicide needed to treat the target area (see the following calibration example). Note: Sprayer calibration (volume of spray needed to treat 1,000 square feet) will vary with each individual operator.

#### Steps in Calibration:

- 1. Mark an area of 1,000 square feet (i.e. 20 by 50 feet or 25 by 40 feet).
- 2. Place the sprayer on a level surface and add water noting the final level of water in the spray tank.
- 3. Spray the marked area with a sufficient volume of water to provide uniform coverage. Refill the sprayer to the same level as before measuring the amount of water added. The measured water added to the sprayer is the volume needed to cover 1,000 square feet.
- 4. Determine the application rate (fl. oz./1,000 sq. ft.) for WEED STOPPER from the Recommendations section of this label.
- 5. To each volume of water used as measured in step 3, add the amount of WEED STOPPER as determined in step 4.

Example: If the sprayer used 2 gallons of water to cover 1,000 square feet and the desired application rate of WEED STOPPER is 3 fluid oz./1,000 square feet, then you would add 3 fluid ounces of WEED STOPPER to every 2 gallons of water to be used.

PRECAUTION: Avoid spray drift to non-target areas when applying WEED STOPPER. Spray drift may result in reduced emergence of non-target plants adjacent to the treated area. Poor weed control may result if directions are not followed. Overapplication may result in crop injury or excessive soil residue.

#### **MIXING DIRECTIONS**

SHAKE WELL BEFORE USING.

Make sure spray tank is clean and use only clean water. Fill spray tank 1/2 to 3/4 full. Start agitation and add the required amount of WEED STOPPER. Continue agitation and finish filling the spray tank. Maintain continuous agitation until application is completed.

#### **EQUIPMENT CLEANING**

If a buildup of material occurs on the walls of the spray tank, it should be removed between fillings by washing with soap and water and rinsing thoroughly. Tanks, lines, screens and nozzles should be cleaned thoroughly after each use.

#### **ACTIVATION AND CULTIVATION**

WEED STOPPER will remain stable on the soil surface up to 21 days following application. In the absence of timely rainfall, irrigation can be used to activate WEED STOPPER. A minimum of one-half (1/2) inch of rain or its equivalent in sprinkler irrigation is necessary to activate WEED STOPPER. If weeds begin to emerge due to lack of rainfall or irrigation, shallow cultivate 1 to 2 inches deep to destroy existing weeds or remove them by hand. Shallow cultivation to a depth of 1 to 2 inches will enhance herbicidal effectiveness. If WEED STOPPER is not activated by rainfall, irrigation or cultivation within 21 days of application or existing weeds have not been removed, erratic weed control may result.

#### WEEDS CONTROLLED BY WEED STOPPER

## Annual Grasses: Barley, little Barnyardgrass (Wetergrass) Bluegrass, armuel Crabgrass, large Crabgrass, amooth

Crowloctoress Cupgrass, southwes Foxtail, bristlegrass Foxtail, giant Foxtail, green (Pigeongrass)

Fordail robust

Foxteil, yellov Goosegrass (Silver crabgrass)

**Broadleaf Weeds:** Bittercress

Carpet weed Chickweed, common Fiddleneck, coast Filaree Groundsel, common Henbit

Knotweed, prostrate

Signalgrass (Brach aria) Sprangletop, red **Puncturevine** 

Lovegrass, orcult Oat, wild

Panicum, browntop

Johnsongrass (Seedling only) Junglerice Lovegrass, mexican

Pursiane, common Pusiey, Florida (Florida pursiane, Mexican clover, pusiey) Rocket, London

remount, prowntop
Panicum, fall (Spreading paniograss)
Panicum, Falas (Buffelograss, Coloradograss)
Ryagrass, Italian
Sendbur, field

Rockpursians, desert
Shapardapurse; c o
Spurga, praetrate: c
Woodsorrel; yellow (axalis)

#### 40000 WEEDS SUPRESSED BY WEED STOFFER

Control of the following weeds may be erratic, ranging from pron to excellent, depending upon soil temperature, time of germination, depth of seed in the soil, and amount and timing of soil moisture:

Horseweed Ladysthumb Lettuce, prickly Mallow, common Milkweed, climbing Morninggiory

Nightshadu, black <sup>6</sup> Ragweed, common Sowthistie, annual Spurge, spotted Teaweed (Prickly side)

# **WEED STOPPER**

#### FOR HOMEOWNER USE

ACTIVE INGREDI	IENT:	
Oryzalin: 3,	5-dinitro-N*,N*-dipropylsulfanilamide	40.40%
OTHER INGREDI	ENTS:	59.60%
TOTAL		00.00%

Contains 4.0 pounds of active ingredient per gallon. EPA Reg. No. 54705-5

EPA Est. No. 48498-CA-1

#### Manufactured for: LAWN AND GARDEN PRODUCTS. INC.

P. O. Box 35000 • Fresno, CA 93745 • (559) 499-2100 www.montereylawngarden.com

#### KEEP OUT OF REACH OF CHILDREN CAUTION

#### **FIRST AID**

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15 20 minutes
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye
- Call a poison control center or doctor for treatment advice.

#### **HOT LINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information

#### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

CAUTION: Causes eye irritation. Prolonged or frequently repeated contact may cause allergic reactions in some individuals. Avoid contact with eyes or clothing. Applicators should wear long-sleeved shirt and long pants, chemical-resistant gloves and shoes plus socks. Wash thoroughly with soap and water after handling. Remove conteminated clothing and wash before reuse. Keep all persons, children and pets out of treated area until sprays have dried.

#### **Environmental Hazards**

This pesticide is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its

READ ENTIRE LABEL BEFORE EACH USE.

#### **GENERAL INFORMATION**

WEED STOPPER herbicide is a preemergence surface-applied product for the control of many annual grasses and broadleaf weeds in ornamental plantings, bedding plants, bulbs, ground covers/perennials and established warm-season turfgrass.

WEED STOPPER is orange in color and may cause temporary discoloration of sprayed surfaces. If this discoloration is undesirable, it may be altered by using a commercially available colorant such as Biazon® or removed by spraying surface with water or washing with an industrial cleaner immediately after application. WEED STOPPER may also be applied with mulch colorants, such as Mulch Magic or Nu-Mulch.

Treatment of Plant Species Not Listed on the Label for WEED STOPPER.

Users who wish to use WEED STOPPER on plant species not recommended on this label may determine the suitability for use by treating a small number of such plants at a recommended rate. Prior to treatment of larger areas, the treated plants should be observed for any sign of herbicidal injury during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious to the target plant species. The user assumes responsibility for any plant damage or other liability resulting from use of WEED STOPPER on plant species not recommended on this label.

#### RECOMMENDATIONS

#### **ORNAMENTAL PLANTINGS**

WEED STOPPER is recommended for use on certain established landscape ornamental plants, including trees, shoubs, ground covers/perennials, and flowers, non-bearing fruit and nut trees, non-bearing vineyards; and omamental bulbs (See "Omamental Bulbs" section for special use directions).

**Broadcast Application Rates** 

Length of Control	WEED STOPPER Fl. Oz./1,000 Sq. Ft.	Minimum Time Between Applications	Total WEED STOPPER Allowed Per Year (Fl. Oz./1,000 Sq. Ft.)
2 to 4 months	1.5	2 Months	6
4 to 8 months	3	4 Months	9

Note: 1 to 2 gallons of water will usually treat 1,000 square feet. Therefore, mix 1.5 to 3 ounces in 1 to 2 gallons of water and apply to 1,000 square feet.

SPECIAL USE PRECAUTIONS: Apply only to established plants that have been transplanted into their growing location for a sufficient period of time to allow the soil to be firmly settled around the roots from packing and rainfall or irrigation.

To avoid possible injury, do not apply WEED STOPPER to:
Ground covers until they are established and well rooted.

- Omamental plantings where there is tikelihood of runoff onto lawn areas.
- Areas containing dichondra or cool season turfgrass species.

Applications of WEED STOPPER over the top of plants with newly forming buds may cause injury. In this situation a directed spray is recommended or imigate immediately after application to wash WEED STOPPER into the soil.

For soils treated with WEED STOPPER during the previous season, plant only the ornamental species listed on this label or injury may occur.

Iceplant: When establishing unrooted ice plant on coarse textured soils in (andscape planting, do not exceed the 1.5 fl. oz. per 1,000 sq. ft. of WEED STOPPER or crop injury may occur.

Note: Injury on the following plant species has been observed following applications of WEED STOPPER and use is not recommended.

Arborvitae, techny (Thuja occidentalis 'Techny')

Begonia (Begonia spp.) Coleus (Coleus hybridus) Deutzia, siender (Deutzia gracilis) Douglas Fir (Pseudotsuga menziesii) Eastern Hernlock (Tsuga canadensis)

#### TANK MIX COMBINATIONS

Tank mix combinations of WEED STOPPER plus Roundup®, and many other labeled herbicides may be used to control undesirable vegetation in omamental areas. WEED STOPPER may also be tank mixed with Gallery\* herbicide and applied preemergence to broaden the spectrum of broadleaf weed control in ornamental areas. Applied as directed, these tank mixes of WEED STOPPER will provide control of susceptible weed species listed on the respective labels. Refer to tank mix product labels for specific use directions, precautions and limitations before use.

WEED STOPPER Plus Roundup®: Tank mix combinations of WEED STOPPER plus Roundup are recommended to control existing undestrable vegetation. Applied as directed, WEED STOPPER plus Roundup will provide postemergence control of susceptible weed species listed on the label for Roundup and residual preemergence control of susceptible weed species listed on the label for WEED STOPPER. Refer to the label for Roundup for specific use directions, precautions and limitations before use. Precautions: Do not apply sprays containing Roundup over the top of ornamental plants. Extreme care must be exercised to prevent sprays containing Roundup from coming in contact with foliage and stems of turfgrasses, trees, shrubs, or other desirable vegetation since severe damage or death may result. If spraying with Roundup in areas adjacent to desirable plants, use a shield to prevent spray from contacting foliage and stems of desirable plants.

WEED STOPPER may be used on the following established plant species:

Abelia, Glossy Acacia Aceve Almond (non-bearing)

Andromede Apple (non-beening) Apricot (non-bearing)
\*Arborvitee Ash Aster

Avocado (non-bearing) Azalas

Baby's Breath Barberry Bellflower Rird of Paradisa Blazing S Bleeding Heart Bottlebrush Bougainvillea Boxwood Вгоот **Brush Cherry** 

Caladium California Laurel Canabarries (non-bearing) Cape Weed Carob Carpet Sugle

Carrot wood

Jaconnese Source Jerseylea Junioir Kiwi (non-bearing) Laurelcherry Leucothoe Lilec Lily Turf Lily-of-the-Nile Lilv-of-the-Valley

Linden Macadamia Nut (non-bearing)

Macnolia Mehogany Manzanita Marigold Mockorange Mondo Grass Moss Rose Mountain Laurel Мирепу Myoporum

Nectarine (non-bearing)

Olive (non-bearing) Oregon Grape Osmanthus Palo Verde

Cassia Pampas Grass Century Plant Panev Peach (non-bearing) Cherry (non-bearing) Chrys Peer (non-bearing) enthemum Cinquefoil Pecan (non-bearing) Citrus (non-bearing) Citrus, Ornamental Pepper Tree Pariwinkle Cleyers, Japanese Petunia Photinia Coreopsis Pine Pistachio (non-bearing) Cotoneer Pittosporum Cottonwood Coyotebush Plum (non-bearing)

Covolomeria Japanese Pomecranate (non-bearing) Current (non-bearing) Prive Protee Cypress

Plumbago

Podocerous

Crabappie

Fir

Crape Myrtle

Daisy Prune (non-bearing) Daylily Pyracantha (Firethorn) 'Deutzie Ramanas Ross Dewberry (non-bearing) Ranunculus Raphioteosis Dogwood Eastercactus Elderberry (non-bearing) Red Ceder Redbud Elm Redwood Escallonia Rhododendron

Eucalyptus Rose Rose of Sheron Eyonymus Rosemary False Spirea Falsecypress Sece Sequoia Ficus Shrimo Plant Fig (non-bearing) Snapdragon Filbert (non-be Spruce Star Jasmina Stonecrop Forsythia Gardenia Sumec Sweet William Gazania Sweetgum Geranium Geum Sycamore Ginigo . Gladiolus Tovan Trumpet Vine

Viburnum Goldennain tree Gooseberry (non-bearing) Grape (non-bearing) Wainut (non-bearing) Heavenly Barriboo (Nandina) Hibiscus Wild Lilac Willow Holly Honey Locust Worken Honeysuckie Hopseedbusi Xylosma

Hosta Yellow Tab ice Plant Yow Impetiens Yucca Zinnia

See list of plant species under Special Use Precautions for which use is not recommended. Note: Non-bearing plants are defined as those which will not bear fruit for at least one year after

#### ORNAMENTAL BULBS

WEED STOPPER may be applied for control of susceptible annual weeds in ornamental-bulbs, e.g. bulbous iris, defiodil (narcissus), hyacinth and tulip. Apply WEED STOPPER to the soil surface 2 to 4 weeks after planting, but prior to the emergence of annual weeds. For fall planted bulbs, apply WEED STOPPER again in late winter or early spring to weed-free soil surfaces.

#### **Broadcast Application Rates**

Time of Application	Soli Texture	WEED STOPPER Fl. Oz./1,000 Sq. Pt.	Minimum Time Between Applications	Total WEED STOPPER Allowed Per Year (Pl. Oz./1,000 Sq. Pt.)
Fall	Coarse	0.5	3 Months 🗈	1
Fall	Medium and Fine	1,0	3 Months	2
Feb March	All Soil Textures	0.5	3 Months _	2

Special Use Precautions

DO NOT apply to tulip plants that have emerged to a height greater than 3/4 inch. DO NOT apply to gladioti corms prior to entergence or less than one inch in diameter.

#### WARM SEASON TUMFGRASSES

WEED STOPPER may be applied as a preamergence treatment for control of annual grasses and certain broadlest weeds in established warm season furf including behingrass. bermudagrasa, buffalograsa, centipeulugrasa, St. Augustiniegrasa, zoysiagrasa and established tall feacue growing in warm season areas. Established turf is defined as a dense turf having a well-anchored root system and healthy, vigorous top growth. Use WEED STOPPER only as a part of a total turf management program that includes good fertilization practices.

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of WEED STOPPER.

WEED STOPPER will not control emerged weeds. Successful preemergence control of weeds listed on this label requires that WEED STOPPER be applied prior to weed germination and be activated by at least one-half (1/2) inch of rainfall or irrigation within 21 days of application.



WEED STOPFER may diffure turf that is not well-established or is stressed or weakened due to unfavorable winter climatic conditions, drought, nematodes, or other factors which damage or weaken turf root systems. Apply WEED STOPPER only to healthy, well-established turf that has a well-anchored root system

DO NOT apply WEED STOPPER in the spring or early summer to tall fescue turfgrass reseeded the previous fall. DO NOT use on other fescue species.

In bermudagrass areas that have been reseeded with winter grasses, a spring application of WEED STOPPER will thin the overseeded grasses.

#### **Annual Grasses Controlled by WEED STOPPER**

lummer Annuals

Bernyardgrass (Watergrass) Crabgrass, large Crabgrass, amooth Crowfootgrass Foxiail, bristlegrass

Foxtail, robust Fontail, yellow Goosegrass (silver crabgrass) Johnsongrass (seedling only) Ryegrass, Italian andbur field

Foxtail, giant Foxtail, green (pigeongrass)

Winter Annuals: Bluegrass, Annual

#### Annual Broadleaf Weeds Controlled by WEED STOPPER

Summer Annuals: Carpetweed Knotweed, Prostrate Pursiane, Common

Winter Annuals: Chickweed, Common

#### Broadleaf Weeds Suppressed by WEED STOPPER:

Groundsel, Common Spurge, Prostrate Woodsorral, Yellow (Oxelis)

#### APPLICATION RATES, FREQUENCY AND TIMING OF APPLICATION

WEED STOPPER can be applied in the spring for summer annual grass and broadleaf weed control, and in the fall for annual bluegrass and winter annual broadleaf weed control.

Broadcast Application Rates:

Use Area	WEED STOPPER Fi. Oz./1,000 Sq. Ft.	Minimum Time Between Applications	Total WEED STOPPER Allowed Per Year (FL Oz./1,000 Sq. FL)
All, except Florida	1 - 1.5	3 Months	4
Florida	1	3 Months	3

#### APPLICATION EQUIPMENT

Apply WEED STOPPER evenly over the turigrass area. Avoid spray pattern skips and overlaps that may result in incomplete coverage or over-application. For best results use application equipment designed to uniformly broadcast liquid herbicides. Check equipment frequently to make sure it is workling properly and distributing spray uniformly.

#### RESEEDING

Herbicides that control annual weeds may also affect establishment of desireable turfgrass seedlings. Reseeding should be delayed for at least 90 - 120 days following application of WEED STOPPER. With reseeding, it is essential that proper cultural practices such as soil cultivation and seedbed preparation, irrigation and fertilization be followed. For satisfactory reseeding results following use of WEED STOPPER, the seeding rate should be increased and equipment designed to play seed in full contact with soil (such as the Rogers Aero Seeder) should be employed.

#### SPECIAL USE PRECAUTIONS

To avoid possible injury, DO NOT apply WEED STOPPER to:

- Cool season turfgrass species (bluegrass, fescues, bentgrasses, etc.).
- Newly sprigged or sodded areas of bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until these turfgrasses are well-established and have well-anchored root
- Newly hydromulched areas of bermudagrass until such areas are well-established.
- · Bermudagrass variety "Sun Turf" when WEED STOPPER is tank mixed with atrazine.

#### NONPLANTED AREAS

Tank mix combinations of WEED STOPPER plus Roundup and many other labeled herbicides may be used to control undesirable vegetation in nonplanted areas such as roadsides, rights-of-ways, etc. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions, and limitaions before use

#### Broadcast Application Rates:

Length of Control	WEED STOPPER FI, Oz_/1,000 Sq. Ft.	Minimum Time Between Applications	Total WEED STOPPER Allowed Per Year (Fi. Oz./1,000 Sq. Ft.)
2 to 4 months	1.5	2 Months	6
8 to 12 months	4.5	8 Months	12

#### STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL STORAGE: Store in a cool dry place, preferably a locked storage area out of reach of

children and pets. Store above 40°F. CONTAINER DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recyling if evallable. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or

#### **CONDITIONS OF SALE AND WARRANTY**

Lawn and Garden Products, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated when used in accordance with the use directions under normal conditions. Lawn and Garden Products, Inc. neither makes, nor authorizes any agent or representative to make, any other warranties, express or implied, including fitness or merchantability.

The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Plant injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and plant conditions, presence of other materials, or other influencing factors, all of which are beyond the control of Lawn and Garden Products, Inc. and Seller. Buyer and user acknowledge and assume all risks and liability resulting from the handling, storage and use of the material not in strict accordance with directions given herewith. In no case shall Lawn and Garden Products, Inc. or the Seller be liable for consequential, special, indirect, or incidental damages or losses resulting from the handling or use of this product. The foregoing is a condition of sale by Lawn and Garden Products, Inc. and is accepted as such by the Buyer.

(Blazon® - Milliken Chemical) (Roundup® - Monsanto) (Gallery\* - DowAgroSciences LLC)

0996/1102(03)

6 & 6038608

£1. . (06

1.0000





November 13, 2002

Document Processing Desk (NOTIF)
Office of Pesticide Programs (H7504-C)
U.S. Environmental Protection Agency
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, DC 20460-0001

ATTENTION:

Ms. Joanne I. Miller, Product Manager (23)

Fungicide-Herbicide Branch

RE:

**WEED STOPPER** 

EPA Reg. No. 54705-5

Dear Ms. Miller:

On behalf of Lawn and Garden Products, Inc., I am submitting the following Notification in response to PR Notice 2001-6 and have made two (2) minor Non-Notification changes per PR Notice 98-10 which are as follows:

1) Revision to the Storage and Disposal statements.

2) Revised the term "Inert" in the ingredient Statement to "Other".

The change of the Hotline Number in the First Aid section. Initially we used CHEMTREC's phone number as our emergency number. We have now revised it to the National Pesticide Information Center emergency number.

I am enclosing the following in support of this Notification:

1) A completed Application for Pesticide, EPA Form 8570-1.

2) Two (2) copies of the revised labeling.

3) A self-addressed return postcard acknowledging USEPA's acceptance of our Notification.

If you require any questions regarding this submission, please contact me at (775) 629-1408.

Regards,

Lynne C. Zahigian

Agent for Lawn and Garden Products, Inc.

CC:

**LGPI-EPAWeed Stopper** 

LCZ-Chron