<b>5</b> 4705-5, 2	6 2003 115					
riffese read instructions on reverse before completing form.	Form Approved. OMB No. 2070-006 . Approvel expires 2-28-95					
United States Environmental Protection Washington, DC 20-						
Application	ion for Pesticide - Section I					
1. Company/Product Number 54705-5	2. EPA Product Manager     3. Proposed Classification       Joanne I. Miller        ✓ None					
4. Company/Product (Name) WEED STOPPER	PM# []					
<ul> <li>5. Name and Address of Applicant (Include ZIP Code)</li> <li>6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3)</li> <li>(b)(i), my product is similar or identical in composition and labeling to:</li> <li>6650 Wade Lane Stagecoach, NV 89429-8443</li> <li>Check If this is a new eddress</li> </ul>						
	Product Name					
	٢					
Amendment - Explain below.           Resubmission in response to Agency letter deted	Agency letter dated NOTIFICATION					
Notification - Explain below.	Other - Explain below. FEB 0 6 2003					
Explanation: Use additional page(s) if necessary. (For section I and Section II.) Notification of minor Storage statement revision relative to PR Notice 98-10. This notification is consistent with the guidance in PR Notice 98-10 and the requirements of EPA regulations at 40 CFR 156.10 and 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that It is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and the requirements of 40 CFR 156.10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under						
sections 12 and 14 of FIFRA.	Section - III					
1. Material This Product Will Be Peckaged In:	Section - III					
Child-Resistant Packaging Unit Packaging	Water Soluble Packaging 2. Type of Container					
Yos	Yes Mistal					
No	No Plastic Glass					
• Cartification must If "Yes" No. per Unit Packaging wgt. No. per Container	If "Yes"         No. per         Paper           Package wgt         container         Other (Specify)					
3. Location of Net Contents Information 4. Size(s) Ret	stail Container 5. bosation of Label Directions					
Label						
6. Manner in Which Label is Affixed to Product Lithog Paper Stend	graph Other					
Section - IV						
1. Contact Point (Complete items directly below for identification	an of individual to be contected, if necessary, to process this enplication.)					
Name Lynne C. Zahigian	Title     Telephone No. (Include Aree Code)       Agent     75/829-1408					
Certifica I certify that the statements I have made on this form and I acknowledge that any knowlingity false or misleading sta	d all attachments thereto are true, acourate and complete.					
both under applicable law.	atement may be purishable by fine or imprisonment of a Stamped)					
both under applicable law.	atement may be punishable by fine or imprisonment of (Stamped) 3. Title					
both under applicable law. 2. Signature Xythe C. Zahigier						

EPA Form 8570-1 (Rev. 3-34) Previous editions are obsolete.

.



# WEED STOPPER

FOR HOMEOWNER USE

ACTIVE INGREDIENT

Oryzalin: 3,5-dinitro-M,M-dipropylsulfanilamide	40.40%
OTHER INGREDIENTS:	59,60%
TOTAL	100.00%

Contains 4.0 pounds of active ingredient per gallon. EPA Reg. No. 54705-5

### Manufactured for: LAWN AND GARDEN PRODUCTS, INC.

EPA Est. No. 48498-CA-1

P. O. Box 35000 • Fresno, CA 93745 • (559) 499-2100

www.montereylawngarden.com

-	KEEP OUT OF REACH OF CHILDREN CAUTION
	FIRST AID
IF IN EYES	<ul> <li>Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
	HOT LINE NUMBER
Have the pro	duct container or label with you when calling a poison control center or

doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.

# PRECAUTIONARY STATEMENTS

# Hazards to Humans and Domestic Animals

CAUTION: Causes eye irritation. Prolonged or frequently repeated contact may cause allergic reactions in some individuals. Avoid contact with eyes or clothing. Applicators should wear long-sleeved shirt and long pants, chemical-resistant gloves and shoes plus socks. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Keep all persons, children and pets out of treated area until sprays have dried.

#### Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

# **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE EACH USE.

#### **GENERAL INFORMATION**

WEED STOPPER herbicide is a preemergence surface-applied product for the control of many annual grasses and broadleaf weeds in omamental plantings, bedding plants, bulbs, ground covers/perennials and established warm-season turfgrass.

WEED STOPPER is orange in color and may cause temporary discoloration of sprayed surfaces. If this discoloration is undesirable, it may be altered by using a commercially available colorant such as Blazone or removed by spraying surface with water or washing with an industrial cleaner immediately after application. WEED STOPPER may also be applied with mulch colorants, such as Mulch Magic or Nu-Mulch.

#### Treatment of Plant Species Not Listed on the Label for WEED STOPPER.

Users who wish to use WEED STOPPER on plant species not recommended on this label may determine the suitability for use by treating a small number of such plants at a recommended rate. Prior to treatment of larger areas, the treated plants should be observed for any sign of herbicidal injury during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious to the target plant species. The user assumes responsibility for any plant damage or other liability resulting from use of WEED STOPPER on plant species not recommended on this label.

# **APPLICATION INSTRUCTIONS** SOIL PREPARATION

WEED STOPPER controls weeds growing from seed. WEED STOPPER will not control emerged weeds. WEED STOPPER does not control established weeds, weeds growing from stolens, rhizomes, or root pieces. Therefore, areas to be treated should be free of emerged weeds. Weed residues, pruning and trash should be thoroughly mixed into the soil or removed prior to treatment.

#### Handheid or Backpack Sprayer Application

The amount of water used to apply WEED STOPPER herbicide is not critical, but should be sufficient for uniform coverage of the target area. Calibrate by determining the volume of water required to treat 1,000 square feet. Use this calibration volume to determine the amount of water and WEED STOPPER herbicide needed to treat the target area (see the following calibration example). Note: Sprayer calibration (volume of spray needed to treat 1,000 square feet) will vary with each individual operator.

Steps in Calibration:

- 1. Mark an area of 1,000 square feet (i.e. 20 by 50 feet or 25 by 40 feet).
- 2. Place the sprayer on a level surface and add water noting the final level of water in the spray tank.
- 3. Spray the marked area with a sufficient volume of water to provide uniform coverage. Refill the sprayer to the same level as before measuring the amount of water added. The measured water added to the sprayer is the volume needed to cover 1.000 square feet.
- 4. Determine the application rate (R. oz./1,000 sq. ft.) for WEED STOPPER from the Recommendations section of this label.
- 5. To each volume of water used as measured in step 3, add the amount of WEED STOPPER as determined in step 4.

Example: If the spraver used 2 gallons of water to cover 1,000 square feet and the desired application rate of WEED STOPPER is 3 fluid oz./1,000 square feet, then you would add 3 fluid ounces of WEED STOPPER to every 2 gallons of water to be used.

PRECAUTION: Avoid spray drift to non-target areas when applying WEED STOPPER. Spray drift may result in reduced emergence of non-target plants adjacent to the treated area. Poor weed control may result if directions are not followed. Overapplication may result in crop injury or excessive soil residue.

#### MIXING DIRECTIONS

# SHAKE WELL BEFORE USING.

Make sure spray tank is clean and use only clean water. Fill spray tank 1/2 to 3/4 full. Start agitation and add the required amount of WEED STOPPER. Continue agitation and finish filling the spray tank. Maintain continuous agitation until application is completed.

# EQUIPMENT CLEANING

If a buildup of material occurs on the walls of the spray tank, it should be removed between fillings by washing with soap and water and rinsing thoroughly. Tanks, lines, screens and nozzles should be cleaned thoroughly after each use.

#### **ACTIVATION AND CULTIVATION**

WEED STOPPER will remain stable on the soil surface up to 21 days following application. In the absence of timely reinfall, irrigation can be used to activate WEED STOPPER. A minimum of one-half (1/2) inch of rain or its equivalent in sprinkler irrigation is necessary to activate WEED STOPPER. If weeds begin to emerge due to lack of rainfail or irrigation, shallow cultivate 1 to 2 inches deep to destroy existing weeds or remove them by hand. Shallow cultivation to a depth of 1 to 2 inches will enhance herbicidal effectiveness. If WEED STOPPER is not activated by rainfall, irrigation or cultivation within 21 days of application or existing weeds have not been removed, erratic weed control may result.

#### WEEDS CONTROLLED BY WEED STOPPER

Annual Grasses: Barley, Ittle Barnyradgrass (Watergrass) Bluegrass, arruai Crabgrass, large Crabgrass, smooth Crowfootgrass Cupgrass, scuthwestern Foxtail, brietlegrass Foxtail, green (Pigeongrass) Foxtail, green (Pigeongrass) Foxtail, choust Foxtail, choust Foxtail, choust Foxtail, choust Foxtail, choust Foxtail, choust Foxtail, choust Foxtail, choust		Rvermes Hellen
Broadiaaf Weeds: Bittercreas Carpet weed Chickweed, common Fiddleneck, coast Fideree Groundsel, common Henbit Knotweed, prostrate Lambequarters Pigweed	с Колас Д н. — В П. — В	Purstere A Purster, Forida (Florida purstane, Mexican clover, pualey) Rockout, London Rockpurshee, desert Shaperdimurse e Spurge, postrate Woodsorre/; seitow (oxalia)

# WEEDS SUPRESSED BY WEED STOPPER

Control of the following weeds may be erratic, ranging from poor to excellent, depending upon soil temperature, time of germination, depth of seed in the soil, and amount and timing of soil moisture:

Horseweed Ladysthumb Lettuce, prickly Mellow, common Milkweed, climbing Morningglory Musterd, blac Musterd, wild

Nichtahade, black Ragweed, common Smartweed Sowthistle, annual Spurge, spotted d (Prickly side) Wheel, volunteer

216

# RECOMMENDATIONS ORNAMENTAL PLANTINGS

WEED STOPPER is recommended for use on certain established landscape ornamental plants, including trees, shrubs, ground covers/perennials, and flowers, non-bearing fruit and nut trees, non-bearing vineyards; and omamental bulbs (See "Omamental Bulbs" section for special use directions).

#### **Broadcast Application Rates**

Length of Control	WEED STOPPER Fl. Oz/1,000 Sq. Fl.	Minimum Time Between Applications	Total WEED STOPPER Allowed Per Year (FI. Oz./1,000 Sq. Ft.)
2 to 4 months	1.5	2 Months	6
4 to 8 months	3	4 Months	9

Note: 1 to 2 galions of water will usually treat 1,000 square feet. Therefore, mix 1.5 to 3 ounces in 1 to 2 gallons of water and apply to 1,000 square feet.

SPECIAL USE PRECAUTIONS: Apply only to established plants that have been transplanted into their growing location for a sufficient period of time to allow the soil to be firmly settled around the roots from packing and rainfall or irrigation. To avoid possible injury, do not apply WEED STOPPER to:

- Ground covers until they are established and well rooted.
- Ornamental plantings where there is likelihood of runoff onto lawn areas.
- Areas containing dichondra or cool season turfgrass species.

Applications of WEED STOPPER over the top of plants with newly forming buds may cause injury. In this situation a directed spray is recommended or imigate immediately after application to wash WEED STOPPER into the soil.

For soils treated with WEED STOPPER during the previous season, plant only the ornamental species listed on this label or injury may occur.

Iceplant: When establishing unrooted ice plant on coarse textured solls in landscape planting, do not exceed the 1.5 fl. oz. per 1,000 sq. ft. of WEED STOPPER or crop iniury may occur.

Note: Injury on the following plant species has been observed following applications of WEED STOPPER and use is not recommended.

Arborvitae, techny (Thuja occidentalis 'Techny') Begonia (Begonia spp.) Coleus (Coleus hybridus) Deutzia, siender (Deutzia gracilis) Douglas Fir (Pseudotsuga menziesii) Eastern Hemlock (Tsuga canadensis)

#### TANK MIX COMBINATIONS

Tank mix combinations of WEED STOPPER plus Roundupe, and many other labeled herbicides may be used to control undesirable vegetation in ornamental areas. WEED STOPPER may also be tank mixed with Gallery" herbicide and applied preemergence to broaden the spectrum of broadleaf weed control in ornamental areas. Applied as directed, these tank mixes of WEED STOPPER will provide control of susceptible weed species listed on the respective labels. Refer to tank mix product labels for specific use directions, precautions and limitations before use.

WEED STOPPER Plus Roundup\*: Tank mix combinations of WEED STOPPER plus Roundup are recommended to control existing undesirable vegetation. Applied as directed, WEED STOPPER plus Roundup will provide postemergence control of susceptible weed species listed on the label for Roundup and residual preemergence control of susceptible weed species listed on the label for WEED STOPPER. Refer to the label for Roundup for specific use directions, precautions and limitations before use. Precautions: Do not apply sprays containing Roundup over the top of omamental plants. Extreme care must be exercised to prevent sprays containing Roundup from coming in contact with foliage and stems of turfgrasses, trees, shrubs, or other desirable vegetation since severe damage or death may result. If spraying with Roundup in areas adjacent to desirable plants, use a shield to prevent spray from contacting foliage and stems of desirable plants.

WEED STOPPER may be used on the following established plant species:

WEED STOPPER may be used on t	ne ronoward esternished heart shecies.
Abelia, Glossy	Jepanese Spurge
Acacia	Jerseytea
Ageve	Juniper
Almond (non-bearing)	Kiwi (non-beening)
Andromeda	Laureicheny
Apple (non-bearing)	Leucothoe
Apricot (non-bearing)	Lilac
*Arborvitae	Lily Turf
Ash	Lily-of-the-Nile
Aster	Lily-of-the-Valley
Avocado (non-bearing)	Linden
Azalea	Macadamia Nut (non-bearing)
Baby's Breath	Magnolia
Barberry	Mahogany
Bellfiower	Manzanita
Birch	Maple
Bird of Paradise	Marigold
Blazing Star	Mockorange
Bleeding Heart	Mondo Grass
Bottlebrush	Moss Rose
Bougainvillea	Mountain Laurel
Boxwood	Mulberry
Bridal Wreath	Myoporum
Broom	Myrtle
Brush Cherry	Nectarine (non-bearing)
Caladium	Oak
California Laurel	Oleander
Caneberries (non-bearing)	Olive (non-bearing)
Cape Weed	Oregon Grape
Carob	Osmanthus
Carpet Bugle	Paim
Carrot wood	Palo Varda

Cessis Century Plant Charry (non-be Chrysanthemum Cinqueloil Citrue (non-beering) Citrus, Omement Clevera, Japane Conellowe Coreopeis Colon Cottonwood Coyotebuah Crabanole Спаре Мутве Cryptomeria, Japan Current (non-beering) Cypress Daisy Daylily Deutzie Dewberry (non-bearing) Dogwood Eastercectus Elderberry (non-bearing) Elm Escalionia Eucelyptus Eyonymus False Spire Falsecypre Fatanecera Ficus Fig (non-bearing) Filbert (non-beering) Forsythia Gardania Gezenia Geranium Gaum Ginigo , Gladiolus **Coldensein tree** Gooseberry (non-bearing) Grape (non-bearing) Heavenly Bamboo (Nandina) Hibiscus Holly Honey Locust Honeysucki Hoosedbush Hosta

Parnoas Grass Paney Peach (non-bearing) Pear (non-bearing) Pecan (non-bearing) Pepper Tree Periwinkle Petunia Photinia Pine Pistachio (non-bearing) Pittosporum Plum (non-bearing) Plumbago Podocarpus Pomegranate (non-baaring) Privet Protect Prune (non-bearing) Pyracantha (Firethorn) Remenes Rose Ranunculus Raphiolep Red Cedar Redbud Redwood Rhododendron Ross of Sharon Rosemary Seco Sequoia Shrimp Plant Snepáragon Spruce Star Jeamine Stonecrop Sumac Sweet William muglee Sycamore Toyon Trumpet Vine Viburnum Walnut (non-bearing) Vilainai Wild Lilac Willow Wintercreece Woedwee Xviosma Yarrow Yellow Tab Yew Ymenine Yucca

\* See list of plant species under Special Use Precautions for which use is not recommended. Note: Non-bearing plants are defined as those which will not bear fruit for at least one year after

Zinnie

#### ORNAMENTAL BULBS

WEED STOPPER may be applied for control of susceptible annual weeds in ornamental builte, e.g. builtous inte, defindil (narcissus), hyscinth and tulip. Apply WEED STOPPER to the soil surface 2 to 4 weeks after planting, but prior to the emergence of annual weeds. For fall planted builds, apply WEED STOPPER again in late winter or early spring to weed-free soil surfaces.

#### **Broadcast Application Rates**

ice Plani

Impatiens

Iris

lvy

Time of Application	Soll Texture	WEED STOPPER Fl. Oz./1,000 Sq. Pt.	Minimum Time Between Applications	Total WEED STOPPER Allowed Per Year (FL. Oz./1,000 Sq. FL)
Fall	Coarse	Q.5	3 Months	1
Fall	Medium and Fine	1.0	3 Months	2
Feb March	All Soil Textures	0.5	3 Months	2

**Special Use Precautions** 

....... DO NOT apply to tulip plants that have emerged to a height greater than 3/4 Inch.

DO NOT apply to gladioli come prior to emergence or less than one inch in diameter.

WARM SEARON TURFGRAMSES

WEED STOPPER may be applied as a presence transitient for control of annual grasses and certain broadleaf weeds in catabilished warm sonesh turf including bahiagrass, bermudagrass, buffalograss, centipedegrass, St. Augustinegrass, zoyslagrass and established tall fescue growing in warm season areas. Established turf is defined as a dense turf having a well-anchored root system and healthy, Vigorous top growth. Use WEED STOPPER only as a part of a total turf management program that includes good fertilization practices.

Any cultural practices that disturb the soil, such as avrification or verticutting, should be done prior to application of WEED STOPPER.

WEED STOPPER will not control emerged weeds. Successful preemergence control of weeds listed on this label requires that WEED STOPPER be applied prior to weed germination and be activated by at least one-half (1/2) inch of rainfall or irrigation within 21 days of application.

WEED STOPPER may injure turf that is not well-established or is stressed or weakened due to unfavorable winter climatic conditions, drought, nematodes, or other factors which damage or weaken turf root systems. Apply WEED STOPPER only to healthy, well-established turf that has a well-anchored root system.

DO NOT apply WEED STOPPER in the spring or early summer to tall fescue turgrass reseeded the previous fall. DO NOT use on other fescue species.

in bermudagrass areas that have been reseaded with winter grasses, a spring application of WEED STOPPER will thin the overseeded grasses.

Annual Grasses Controlled by WEED STOPPER

Summer Annuals Barnyardgrass (Watergrass) Crabgrass, large Crabgrass, smooth Crowfootgrass Foxtail, briatlegrass oxtail, giant Foxtail, green (pigeongrass)

todail, robust Foxtail, yellow Foxtail, yellow Goosegrass (silver crabgrass) Johnsongrass (seedling only) Rypornes Itali dur, field

Winter Annuals: Bluegrass, Annual

Annual Broadleaf Weeds Controlled by WEED STOPPER Summer Annuals: Carpetweed Kontweed Prostrate Purslane, Commo

Winter Annuals: Chickweed, Common Henhit

# Broadleaf Weeds Suppressed by WEED STOPPER:

Groundsel, Common Source, Prostrate

Woodsorrel, Yellow (Oxalis)

# APPLICATION RATES, FREQUENCY AND TIMING OF APPLICATION

WEED STOPPER can be applied in the spring for summer annual grass and broadleaf weed control, and in the fall for annual bluegrass and winter annual broadleaf weed control.

#### **Broadcast Application Rates:**

Use Area	WEED STOPPER Fl. Oz./1,000 Sq. Ft.	Minimum Time Between Applications	Total WEED STOPPER Allowed Per Year (Fl. Oz./1,000 Sq. Fl.)
All, except Florida	1-1.5	3 Months	4
Florida	1	3 Months	3

#### APPLICATION EQUIPMENT

Apply WEED STOPPER evenly over the turigress area. Avoid spray pattern skips and overlaps that may result in incomplete coverage or over-application. For best results use application equipment designed to uniformly broadcast liquid herbicides. Check equipment frequently to make sure it is workling property and distributing spray uniformly.

# RESEEDING

Herbicides that control annual weeds may also affect establishment of desireable turgrass seedlings. Reseeding should be delayed for at least 90 - 120 days following application of WEED STOPPER. With reseeding, it is essential that proper cultural practices such as soil cultivation and seedbed preparation, irrigation and fertilization be followed. For satisfactory reseeding results following use of WEED STOPPER, the seeding rate should be increased and equipment designed to play seed in full contact with soil (such as the Rogers Aero Seeder) should be employed.

#### SPECIAL USE PRECAUTIONS

To avoid possible injury, DO NOT apply WEED STOPPER to:

- Cool season turigrass species (bluegrass, fescues, bentgrasses, etc.).
- Newly sprigged or sodded areas of bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until these turigrasses are well-established and have well-anchored root systems
- Newly hydromulched areas of bermudagrass until such areas are well-established. Bermudagrass variety "Sun Turt" when WEED STOPPER is tank mixed with atrazine.

#### NONPLANTED AREAS

Tank mix combinations of WEED STOPPER plus Roundup and many other labeled herbicides may be used to control undesirable vegetation in nonplanted areas such as roadsides, rights-of-ways, etc. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions, and limitations before use.

#### **Broadcast Application Rates:**

Length of Control	WEED STOPPER Fl. Oz./1,000 Sq. Ft.	Minimum Time Between Applications	Total WEED STOPPER Allowed Per Year (Fl. Oz./1,000 Sq. Fl.)
2 to 4 months	1.5	2 Months	6
6 to 12 months	45	8 Months	12

# STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL STORAGE: To be stored in original container in a cool, well-ventilated area inaccessible to children and pets. Store above 40°F.

CONTAINER DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain

# CONDITIONS OF SALE AND WARRANTY

Lawn and Garden Products, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated when used in accordance with the use directions under normal conditions. Lawn and Garden Products, Inc. neither makes. nor authorizes any agent or representative to make, any other warranties, express or implied, including fitness or merchantability

The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Plant injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and plant conditions, presence of other materials, or other influencing factors, all of which are beyond the control of Lawn and Garden Products, Inc. and Seller. Buyer and user acknowledge and assume all risks and liability resulting from the handling, storage and use of the material not in strict accordance with directions given herewith. In no case shall Lawn and Garden Products, Inc. or the Seller be liable for consequential, special, indirect, or incidental damages or losses resulting from the handling or use of this product. The foregoing is a condition of sale by Lawn and Garden Products, Inc. and is accepted as such by the Buver.

(Blazon® - Milliken Chemical) (Roundup® - Monsanto) (Gallery\* - DowAgroSciences LLC)

.

0996/0103(03)

. ....

8 25 C

1.10

Real Sector

2010/01/2

. . . . .

1. 1. 1. 1.

÷ .

۰. v 2 6 6



January 16, 2003

Document Processing Desk (NOTIF) Office of Pesticide Programs (H7504-C) U.S. Environmental Protection Agency Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, DC 20460-0001

ATTENTION: Ms. Joanne I. Miller, Product Manager (23) Fungicide-Herbicide Branch

RE: WEED STOPPER EPA Reg. No. 54705-5

Dear Ms. Miller:

On behalf of Lawn and Garden Products, Inc., I am submitting the following Notification regarding a minor Storage statement revision per PR Notice 98-10. The revision consists of:

1) Changing the Storage statement back to the last EPA approved amendment. When submitting an amendment to California, I realized that the latest Storage statement was really not correct or accurate. Therefore, I am notifying the Agency of this revision back to a more accurate statement.

I am enclosing the following in support of this Notification:

- 1) A completed Application for Pesticide, EPA Form 8570-1.
- 2) Two (2) copies of the revised labeling.
- 3) A self-addressed return postcard acknowledging USEPA's acceptance of our Notification.

00000

ခ် င ဂိ ဒေဇ ဂန

0.000

5002

.. <del>-</del> 6

If you require any questions regarding this submission, please contact me at (775) 629-1408.

Recards.

CC:

Lyrine C. Zahigian/

Agent for Lawn and Garden Products, Inc.

¥

LGPI-EPA/Weed Stopper LCZ-Chron