

54705-7

11/03/2005

1/20

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

 Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 54705-7	2. EPA Product Manager Tony Kish - Acting Product Mgr.	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) LIQUI-COP	PM# 22	
5. Name and Address of Applicant (Include ZIP Code) Lawn and Garden Products, Inc. c/o Lynne Zahigian Regulatory Consulting P.O. Box 1566 Fallon, NV 89407-1566 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION NOV - 3 2005
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Submission of revised Storage and Disposal Statement per Agency letter dated August 19, 2005. This notification is consistent with the guidance in PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
* Certification must be submitted	If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	Other (Specify) _____	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Lynne C. Zahigian	Title Agent	Telephone No. (Include Area Code) (775) 423-7122
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Agent	
4. Typed Name Lynne C. Zahigian	5. Date 11/4/05	

LIQUI-COP

Copper Fungicidal Garden Spray



FOR HOMEOWNER USE ONLY

Active Ingredient	
CopperAmmonium	
Complex*	31.4%
Other Ingredients	68.6%
Total	100.0%


*Metallic Copper Equivalent: 8.0%
Contains 0.784 lbs. Copper per gallon.

EPA Reg. No. 14706-7
EPA Est. No. 45989-CA-1

**KEEP OUT OF
REACH OF CHILDREN
CAUTION**

(See Always/Read/and/Precautionary
Statements and Directions for Use)

**NET CONTENTS:
1 QUART / 946 mL**



3/20

LIQUI-COP

Copper Fungicidal Garden Spray



Controls Many Diseases
Attacking Citrus, Fruits, Nuts,
Vegetables and Ornamentals


Manufactured For:
LAWN AND GARDEN PRODUCTS, INC.
P. O. Box 35000 • Fresno, CA 93745
(559) 499-2100
www.montereylawnngarden.com



NET CONTENTS: 1 QUART/946 mL

0100/0101(02)

Job #20212



4/20

KEEP OUT OF REACH OF CHILDREN CAUTION	
FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye • Call a poison control center or doctor for treatment advice
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15-20 minutes • Call a poison control center or doctor for treatment advice
IF SWALLOWED	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by the poison control center or doctor • Do not give anything by mouth to an unconscious person
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible • Call a poison control center or doctor for further treatment advice
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.	

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Causes skin and eye irritation. Harmful if swallowed, absorbed through the skin or inhaled. May cause skin sensitization reactions in certain individuals. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Wear long-sleeved shirt and long pants, chemical resistant gloves and shoes plus socks when applying this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Keep unprotected persons, children and pets out of treated areas until sprays have dried.

5/20

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. **SHAKE WELL BEFORE USING.**

The control of fungal disease is based on prevention, including keeping garden and border areas clean, destroying leaves in the fall, and applying fungicides either before or at the onset of disease. Plant surfaces must be completely covered with fungicide to successfully prevent infection. Repeat applications may be needed for adequate control. Use the highest rate indicated during conditions of severe disease pressure, such as wet or humid weather. The lower rate is suitable for general preventive applications.

MIXING AND SPRAYING DIRECTIONS

Fill spray tank half full of water. Add recommended amount of LIQUI-COP. Add balance of water.

Using either a hose-end or pressurized hand sprayer, apply at the rate of one gallon mixed spray solution to 200 sq. ft. of garden area or up to two gallons mixed spray solution to a large tree. Thoroughly wet foliage to the point of runoff, taking care to cover all plant surfaces. Agitate the container of mixed spray solution regularly during application to maintain an adequate suspension.

NOTE: This product may be reactive on metal and masonry surfaces such as galvanized roofing. AVOID contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.

FRUITS AND NUTS

Crop	Disease Controlled	Rate/1 Gal Water	
		Rate/1 Gal Water	Instructions
ALMONDS	Brown Rot	3 - 4 Tb.p.	Apply at delayed dormant* bud swell stage.
	Shot Hole	3 - 4 Tb.p.	Apply at leaf fall to protect buds and shoots from infection during rainy periods. Reapply every 3 to 4 weeks up until late bud swell (the period in Spring before buds begin to open). Do not apply after full bloom.

*Delayed Dormant — The period just before blossom buds break open.

FRUITS AND NUTS (continued)

Crop	Disease Controlled	Rate/ 1 Gal Water	Instructions
APPLES	Anthraxnose	3 - 4 tsp	Apply before fall rains. Use on yellow varieties may cause discoloration. To avoid, pick before spraying.
	Apple Scab (Black Spot) Bacterial Canker Blossom and Shoot Blast	3 - 4 Tbsp	Apply post-harvest before fall rains.
	Crown or Collar Rot	1 Tbsp	Apply as a drench on lower trunk either in spring or in late fall after harvest. Do not use if soil pH is below 5.5 since copper toxicity may result.
	Fireblight	2 - 4 tsp	Make application when buds are silvery but before most have turned green. Phytotoxicity may occur from late applications. Discontinue use when green-tip is 1/2 inch.
APRICOTS	Brown Rot Blossom Blight	3 - 4 Tbsp	Apply when buds show red color until brown bud jackets have fallen.
	Shot Hole	3 - 4 Tbsp	Apply in late fall or early winter before winter rains or snow. Repeat in late winter or early spring before buds break. Do not apply after full bloom.
AVOCADOS	Anthraxnose	3 Tbsp	Begin application when the flower buds begin to swell and continue at monthly intervals until August.
BANANAS	Sigatoka	2 tsp	Apply every 3-4 weeks.
	Black Pitting	1-3/4 - 3 Tbsp	Apply directly to the fruit stem and include the basal portion of the leaf crown. Apply during the first and second weeks after fruit emergence.

7/100

FRUITS AND NUTS (continued)

Crop	Disease Controlled	Rate/ 1 Gal Water	Instructions	
BLUEBERRIES	Bacterial Canker	3 - 4	Apply before fall rains and again 4 weeks later. In the spring during wet weather, apply at 10-14 day intervals beginning at leaf emergence.	
	Cane Canker	tsp		
CANEberries	Anthraxnose	2 tsp	Apply when leaf buds open. Repeat when flower buds show white and continue at 10-14 day intervals.	
	Leaf and Cane Spot			
	Purple Blotch			
	Yellow Rust			
	Anthraxnose	4 tsp	Apply in the fall after harvest.	
	Bacterial Blight			
	Leaf and Cane Spot			
	Purple Blotch			
CHERRIES	Deadbud	1-1/2	Apply in October (before heavy fall rains) and again in January. Where disease is severe, another application should be applied in August.	
	Coryneum Blight	tsp		
	Brown Rot	2 - 3	Apply as a full cover spray at popcorn stage and at full bloom.	
	Blossom Blight			tsp
CITRUS	Greasy Spot	3/4 - 1	Apply as a pre-bloom and post bloom sprays.	
	Melanose			tsp
	Pink Pitting Scab			
	Brown Rot	2 - 6	Begin applications in the fall before or just after heavy rains.	
		tsp		
COCOA	Black Pod Rot	2 - 3	Apply on a 14-21 day schedule in high rainfall areas.	
COFFEE	Iron Spot	1 - 3	Apply 3 applications at monthly intervals at the beginning of the wet season.	
	Pink Disease	Tbsp		
	Bacterial Blight	1-1/4	Apply as locally recommended, usually at 3-4 week intervals depending upon disease severity and rainfall conditions.	
	Berry Spot	3 Tbsp		
	Leaf Rust			
	Leaf Spot			

8/20

FRUITS AND NUTS (continued)			
Crop	Disease Controlled	Rate/1 Gal Water	Instructions
CRANBERRIES	Fruit Rot	3 Tbsp	Apply at late bloom. One or two additional applications made at 10-14 day intervals may be required.
CURRENTS GOOSEBERRIES	Anthraxnose Leaf Spot (Cane Blight)	1-1/2 2 Tbsp	Make 3 applications starting after harvest, before bloom and after petal fall.
FILBERTS	Bacterial Blight	3 - 4 Tbsp	Apply after harvest in late August or early September in seasons of heavy rainfall, apply again when 3/4 of the leaves have dropped.
	Eastern Filbert Blight	3 - 4 Tbsp	Apply after harvest before heavy rains begin. The next application should be made in late February or early March, followed by another application 1 month later.
GRAPES	Anthraxnose Black Rot Downy Mildew Powdery Mildew	2 tsp	Make first application when new shoots are 6-8 inches long and repeat at 4 to 10 day intervals. NOTICE: Slight to severe foliage injury may occur on copper sensitive varieties such as Concord, Delaware, Niagara and Rosetti.
HOPS	Downy Mildew	2 tsp	Apply as needed at 10 day intervals. Begin with crown treatment (after pruning but before training) and continue until 2 weeks before harvest.
KIWI	<i>Pseudomonas syringae</i> <i>Erwinia herbicola</i> <i>Pseudomonas fluorescens</i>	3 Tbsp	Make applications on a monthly basis. A maximum of 3 applications may be made.
LIMES	Greasy Spot	3 Tbsp	Apply in June and continue at monthly intervals through August.

9/20

FRUITS AND NUTS (continued)

Crop	Disease Controlled	Rate/ 1 Gal Water	Instructions
MANGOS	Anthracnose	3 Tbsp	Apply weekly from the time the flowers are 2 inches in length until all fruits are set and monthly thereafter until August
OLIVES	Peacock Spot	3 - 4 Tbsp	Make first application before winter rains fall. A second application should be made in early spring if disease is severe
PEACHES NECTARINES	Bacterial Spot	2 tsp	Apply as dormant spray and/or use 1/4 tsp as a post bloom spray. Apply twice during the season, but do not use later than 3 weeks prior to harvest. Do not use at rates above those recommended. NOTICE: Slight loss of leaves may occur from post bloom use
	Blossom Brown Rot	3 - 4 Tbsp	Apply as a dormant or delayed dormant* spray. Do not apply at or after full bloom
	Leaf Curl	3 - 4 Tbsp	Apply at leaf fall. Clean up and destroy fallen leaves
	Shot Hole	3 - 4 Tbsp	Apply at leaf fall to protect buds and shoots from infection during rainy periods. Reapply every 3 to 4 weeks up until late bud swell. Do not apply after full bloom
PEARS QUINCE	Fire Blight	1 tsp	Apply at 5 to 7 day intervals throughout bloom period. Excessive dosages may discolor fruit surface. DO NOT use on d'Anjou, Comice or Seckel varieties
	Blossom Blast	3 - 4 Tbsp	Apply as a dormant spray. Apply only at bud break to control primary infection

*Delayed Dormant --- The period just before blossom buds break open

10/20

FRUITS AND NUTS (continued)

Crop	Disease Controlled	Rate/ 1 Gal Water	Instructions
PECANS	Shuck and Kernel Rot	1 - 3 Tbsp	For suppression, apply in sufficient water to ensure complete spray coverage at 2-4 week intervals starting at kernel growth and continuing until shucks open. Use the higher rate and shorter interval if frequent rainfall occurs.
	Zonate Leaf Spot		
PISTACHIOS	Alternaria Late Blight	3 - 4 Tbsp	Apply at 50% and full bloom followed by up to 3 applications at 30 day intervals.
	Botrytis Blight	3 - 4 Tbsp	Make initial application at bud swell and repeat on a 14-28 day schedule as dictated by disease conditions. If disease conditions are severe, use the higher rate and shorter interval.
	Botryosphaeria Panicle		
	Shoot Blight Septoria Leaf Blight		
PLUMS PRUNES	Brown Rot Blossom Blight Shot Hole	3 - 4 Tbsp	Apply as a dormant spray before heavy rains begin. For brown rot, apply at early green bud to full popcorn stages.
	Bacterial Blast Bacterial Canker	3 - 4 Tbsp	Apply at dormant to early pink stage. Where disease is severe, apply 1 tsp. per gallon of water at 2 week intervals post-bloom. Slight leaf injury may occur.
STRAWBERRIES	Leaf Spot Scorch	1 - 2 tsp	Spray at 7 to 10 day intervals from the time new growth starts until harvest.
	WALNUTS	Walnut Blight	3 - 4 Tbsp

11/20

VEGETABLES

Crop	Disease Controlled	Rate/ 1 Gal Water	Instructions
BEANS, PEAS LENTILS (Succulent and Dry)	Bacterial Blight (Halo and Common)	1 - 3 tsp	Apply when plants are 3 to 5 inches high and before diseases appear Repeat at 7 to 10 day intervals or at 5 to 7 day intervals under conditions of severe disease pressure
BEETS	Cercospora Leaf Spot	2 - 3 tsp	Apply when disease appears making 3 to 6 sprays at 10 to 14 day intervals Apply more frequently under severe disease pressure
BRUSSELS SPROUTS CABBAGE CAULIFLOWER GREENS (Collard Mustard and Turnip)	Blackleaf Spot Black Rot	2 tsp	Begin applications after transplants are set in the garden or shortly after the emergence of seedlings, or when conditions favor disease development Continue at 7 to 10 day intervals NOTICE: A slight reddening of older leaves may occur on broccoli and a slight flecking of outer leaves may occur on cabbage at this rate
	Downy Mildew (Cabbage only)	1 - 2 tsp	Begin applications after transplants are set in the garden, or shortly after the emergence of seedlings, or when conditions favor disease development Continue at 7 day intervals
CANTALOUPE MELONS CUCUMBERS SQUASH	Alternaria Leaf Spot Angular Leaf Spot Anthracnose Downy Mildew Gummy Stem Blight Powdery Mildew Watermelon Bacterial Fruit Blotch	2 - 3 tsp	Apply when disease appears and repeat at 7 to 10 day intervals

12/20

VEGETABLES (continued)			
Crop	Disease Controlled	Rate/ 1 Gal Water	Instructions
CARROTS	Early Blight	2 - 3	Apply when plants are 6" high. Make 3 to 5 applications at 7 to 10 day intervals.
	Late Blight	tsp	
CELERY	Bacterial Blight	2 - 3	Apply on first appearance of disease. Make 4 to 7 applications at 7 day intervals.
	Early Blight	tsp	
	Late Blight	tsp	
CORN (Sweet)	Bacterial Rot	2 tsp	Apply when disease first appears and repeat as necessary.
	Bacterial Stripe		
	Bacterial Wilt		
	Leaf Blight		
	Stalk Rot		
EGGPLANT	Alternaria Blight	2 tsp	Apply before disease appears and repeat at 7 to 10 day intervals.
	Anthracoise		
	Phomopsis		
ORIONS	Downy Mildew	2 tsp	Apply when plants are 4 to 6 inches high and repeat at 7 to 10 day intervals.
	Purple Blotch		
PEPPERS	Bacterial Spot	1 - 1/2	Make first application upon emergence of seedlings or immediately after transplanting. Apply at 7 to 10 day intervals. When disease is severe, apply at 4 to 5 day intervals. Disease control is critical during fruiting.
	Cercospora Leaf Spot	3 tsp	
	Spot	tsp	
POTATOES	Early Blight	1 - 1/2	Apply at first appearance of disease at 7 to 10 day intervals.
	Late Blight	2 tsp	
SPINACH	Anthracoise	1 - 1/2	Begin when disease first appears. Repeat at 7 to 10 day intervals.
	Cercospora		
	Leaf Spot		
	Downy Mildew		
TOMATOES	Bacterial Speck	1 - 3	Begin applications upon emergence of seedlings or immediately after transplanting. Apply at 7 to 10 day intervals. When disease is severe, apply at 4 to 5 day intervals. Complete coverage is essential for disease control.
	Bacterial Spot		
	Early Blight		
	Late Blight		

13/20

MISCELLANEOUS			
Crop	Disease Controlled	Rate/ 1 Gal Water	Instructions
ATAMOYA	Anthracoese	1 Tbsp	Apply just before flowering and repeat on a weekly schedule until just before harvest.
CARAMBOLA	Anthracoese	1 Tbsp	Apply just before flowering and repeat on a weekly schedule until just before harvest.
CHIVES	Downy Mildew	2 tsp	Apply when plants are first established and repeat at 7 to 10 day intervals depending upon disease severity.
DILL	Phyoma Leaf Spot Rhizoctonia Foliage Blight	1 Tbsp	Apply when plants are first established and repeat at 7 to 10 day intervals depending upon disease severity.
DOUGLAS FIR	Rhabdocline Needlecast	2 tsp	Apply at bud break and repeat at 3 to 4 week intervals.
GINSENG	Alternaria Leaf Stem Blight	1 Tbsp	Apply as soon as plants have emerged in spring making sure to thoroughly cover the stems with spray. Repeat every 7 days until plants become dormant in fall.
GUAVA	Anthracoese Red Algae	1 Tbsp	Apply just before flowering and repeat on a weekly schedule until just before harvest.
LITCHI	Anthracoese	1 Tbsp	Apply just before flowering and repeat on a weekly schedule until just before harvest.
LIVE OAK	Ball moss	2 Tbsp	Apply in spring when ball moss is actively growing. Make sure to wet ball moss tufts thoroughly. A second application may be necessary after 12 months. NOTE: LIQUI-COP may be injurious to ornamentals grown under Live Oaks.

14/20

MISCELLANEOUS (continued)

Crop	Disease Controlled	Rate/1 Gal Water	Instructions
MACADAMIA	Anthracnose	2 Tbsp	Apply at first sign of flowering and repeat on a weekly schedule until just before harvest
	Phytophthora Blight Raceme Blight	1 - 3 Tbsp	Apply during raceme development and bloom periods
MAMEY SAPOTE	Anthracnose Algal Leaf Spot	2 Tbsp	Apply when conditions favor disease Repeat on a 14 to 30 day schedule as necessary
PAPAYA	Anthracnose	1 1/2 3 Tbsp	Apply before disease appears and repeat on a 10 to 14 day interval Apply at 5 to 7 day intervals during periods of heavy rainfall Use higher rates when conditions favor disease
PARSLEY	Bacterial Blight	1 Tbsp	Apply when plants are first established and repeat at 5 to 7 day intervals
PASSION FRUIT	Anthracnose	3 Tbsp	Apply just before flowering and repeat on a weekly schedule until just before harvest
SUGAR APPLE (Annona)	Anthracnose	3 - 4 Tbsp	Apply just before flowering and repeat on a weekly schedule until just before harvest
SYCAMORE	Anthracnose	2 - 3 tsp	Apply as a full coverage spray at bud crack with a second application 7 to 10 days later

15/20

ORNAMENTALS

Apply as a thorough coverage spray using 3 tablespoons per 1 gallon of water. Begin application at first sign of disease and repeat at 7-14 day intervals as needed. Use shorter interval during periods of frequent rains or when severe disease conditions persist.

Notice to User: Plant sensitivities to LIQUI-COP have been found acceptable in specific genera and species listed on this label, however it is impossible to know sensitivities under all conditions and phytotoxicity may occur. Due to the large numbers of species and varieties of ornamentals, it is impossible to test every one for sensitivity to LIQUI-COP. Neither the manufacturer or seller recommends use upon species not listed on the label nor has it been determined that LIQUI-COP can safely be used on ornamental plants not listed on this label.

Crop	Disease Controlled
Althaea (Rose of Sharon)	Bacterial leaf spot
Aralia	Xanthomonas leaf spot, Cercospora leaf spot, Alternaria
Arborvitae	Alternaria twig blight, Cercospora leaf blight
Azalea ¹⁾	Cercospora leaf spot, Botrytis blight, Phytophthora dieback, Powdery mildew
Begonia	Bacterial leaf spot (Xanthomonas sp, Erwinia sp, Pseudomonas sp)
Bougainvillea	Anthrachnose, Bacterial leaf spot
Bulbs (Tulip, Gladiolus)	Anthrachnose, Botrytis blight
Camellia	Anthrachnose, Bacterial leaf spot
Camphor tree	Pseudomonas leaf spot
Canna	Pseudomonas leaf spot
Carnation ¹⁾	Alternaria blight, Pseudomonas leaf spot, Botrytis blight
Chinese tallow tree	Bacterial leaf spot (Xanthomonas sp, Pseudomonas sp)
Chrysanthemum ¹⁾	Septoria leaf spot, Botrytis blight
Cotoneaster	Botrytis blight
Dahlia	Alternaria leaf spot, Botrytis gray mold, Cercospora leaf spot
Date palm	Pestalotia leaf spot
Dianthus	Bacterial spot, Bacterial soft rot

¹⁾Discoloration of foliage and/or blooms has been noted on some varieties

16/20

ORNAMENTALS (continued)	
Crop	Diseases Controlled
Dogwood	Anthracoese
Dusty miller	Bacterial leaf spot (Pseudomonas cichorii)
Easter lily	Botrytis blight
Echinaceae	Bacterial leaf spot (Pseudomonas cichorii)
Elm - Drake	Xanthomonas leaf spot
Euonymus	Botrytis blight, Anthracnose
European fan palm	Pestalotia leaf spot
Gardenia	Alternaria leaf spot, Botrytis bud rot, Cercospora leaf spot
Geranium	Alternaria leaf spot, Botrytis gray mold, Cercospora leaf spot
Gladiolus	Alternaria leaf spot, Botrytis gray mold, Bacterial leaf blight
Goldenrain tree	Bacterial leaf spot
Hibiscus	Bacterial leaf spot
Holly fern	Pseudomonas leaf spot
Impatiens	Bacterial leaf spot
India hawthorn	Anthracoese, Entomosporium leaf spot
Ivy (English, Algerian) ¹	Xanthomonas leaf spot
Ixora	Xanthomonas leaf spot
Juniper (Eastern red cedar)	Anthracoese
Lantana	Bacterial leaf spot
Lilac	Cercospora leaf spot
Loblolly bay	Anthracoese
Loquat	Entomosporium maculata, Colletotrichum sp
Magnolia (Saucer)	Bacterial leaf spot
Magnolia (Southern)	Algal leaf spot, Anthracnose, Bacterial leaf spot
Magnolia (Sweet bay)	Anthracoese
Mandevilla	Anthracoese
Marigold	Alternaria leaf spot, Botrytis leaf and flower rot, Cercospora leaf spot
Mulberry weeping	Bacterial leaf spot

¹Discoloration of foliage and/or blooms has been noted on some varieties

17/20

ORNAMENTALS (continued)

Crop	Disease Controlled
Oak, laurel	Algal leaf spot (<i>Cephalosporium virescens</i>)
Oleander	Bacterial leaf spot, Fungal leaf spot
Pachysandra	Voluella leaf blight
Pansy	Downy mildew
Pear (Flowering)	Fireblight, Leaf spot
Pentas (Egyptian star)	Bacterial leaf spot (<i>Xanthomonas</i> sp.)
Peony	Botrytis blight
Periwinkle	Phomopsis stem blight
Philodendron	Bacterial leaf spot
Phlox	Alternaria leaf spot
Photinia	Anthracoze, Entomosporium
Pistachio	Anthracoze
Plantain lily	Bacterial leaf spot
Powder puff plant	Bacterial leaf spot
Pyracantha	Fireblight, scab
Queen palm	Exosporium leaf spot, Phytophthora bud rot
Rhododendron	Alternaria flower spot
Rose ⁽¹⁾	Powdery mildew, Black spot
Verbena	Xanthomonas leaf spot
Viburnum	Anthracoze
Washingtonia palm	Pestalotia leaf spot
Weeping willow	Anthracoze
Yucca (Adam's needle)	Cercospora leaf spot, Septoria leaf spot

⁽¹⁾Discoloration of foliage and/or blooms has been noted on some varieties

ALGAE CONTROL FOR ORNAMENTAL LAWNS AND TURF

To control algae in ornamental lawns and turf, apply 1 pint of LIQUI-COP in 5 gallons of water per 1 000 square feet. LIQUI-COP may be used alone or in combination with other registered fungicides as a maintenance spray. Observe all precautions and limitations on the label of each product used in tank mixes.

NOTE: Phytotoxicity may occur depending upon varietal differences. Apply the recommended rate to a small area and observe for 7 to 10 days for signs of injury. If phytotoxicity occurs, discontinue use. Do not apply in a spray solution with a pH of less than 6.5.

18/20

FROST PROTECTION

Bacterial Ice Nucleation Inhibitor --- Application of LIQUI-COP to all Use Sites listed on this label at rates indicated, just prior to anticipated frost conditions, will sustain control of ice nucleating bacteria (*Pseudomonas syringae*, *Erwinia herbicola* and *Pseudomonas fluorescens*) and may therefore provide protection against light frost.

NOTE: Not recommended for those geographic areas where weather conditions favor severe frost.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal

PESTICIDE STORAGE: Store unused pesticide in original container only, in cool, dry area out of reach of children and animals, preferably in a locked storage area. **KEEP FROM FREEZING** Do not store in areas where temperatures frequently exceed 100°F. After use, replace cap on pesticide container. Rinse hose and unit outdoors in clear water.

PESTICIDE DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONDITIONS OF SALE AND WARRANTY

Lawn and Garden Products, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated when used in accordance with the use directions under normal conditions. Lawn and Garden Products, Inc. neither makes nor authorizes any agent or representative to make any other warranties, express or implied including fitness or merchantability.

The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Plant injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and plant conditions, presence of other materials, or other influencing factors, all of which are beyond the control of Lawn and Garden Products, Inc. and Seller. Buyer and user acknowledge and assume all risks and liability resulting from the handling, storage and use of this material not in strict accordance with directions given herewith. In no case shall Lawn and Garden Products, Inc. or the Seller be liable for consequential, special, indirect, or incidental damages or losses resulting from the handling or use of this product. The foregoing is a condition of sale by Lawn and Garden Products, Inc. and is accepted as such by the Buyer.

19/20

LIQUI-COP

Copper Fungicidal Garden Spray



Controls Many Diseases Attacking Citrus,
Fruits, Nuts, Vegetables and Ornamentals

FOR HOMEOWNER USE ONLY

Active Ingredient.

Copper Ammonium Complex*	31.4%
Other Ingredients:	68.6%
Total	100.0%

*Metallic Copper Equivalent, 8.0%
Contains 0.784 lbs. Copper per gallon.

EPA Reg. No. 54705-7

EPA Est. No. 48498-CA-1

Manufactured For:

LAWN AND GARDEN PRODUCTS, INC.

P. O. Box 35000 • Fresno, CA 93745

(559) 499-2100

www.montereylawngarden.com



NET CONTENTS: 1 QUART / 946 mL

PEEL BACK BOOK HERE AND RESEAL AFTER OPENING

20/20

Lynne
LZ **Zahigian**
Regulatory Consulting

October 4, 2005

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
Room 266A, Crystal Mall #2
1801 South Bell Street
Arlington, VA 22202-4501

ATTN: Tony Kish, Acting PM 22
Fungicide Branch


RE: Notification of Revision to the Storage and Disposal Statement in Response to Agency
Letter dated August 19, 2005
LIQUI-COP, EPA Reg. No. 54705-7

On behalf of Lawn and Garden Products, Inc., I am responding to your August 19, 2005 letter regarding our label amendment dated July 25, 2005. On September 6, 2005, I sent e-mails to you and to Rose Kearns stating that we decided to drop that label amendment, but that we were going to address the Storage and Disposal revision you requested. I am now submitting the following in support of our Notification regarding the Storage and Disposal revision:

- 1) A completed Application for Pesticide with the certification statement per PR Notice 98-10.
- 2) Two (2) copies of our market label (one of which is highlighted to show the revision).
- 3) A self-addressed stamped envelope and Notification Form with the product name, EPA Reg. No. and Notification Action, so that I may be notified of this acceptance.

If you have any questions regarding this submission, please contact me at (775) 423-9122.

Regards,


Lynne C. Zahigian
Agent — Lawn and Garden Products, Inc.

cc: LGPI-EPA/Liqui-Cop
LCZ-Chron

