

54705-7

12/6/2012

11/18



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D C 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Ms Lynne C Zahigian
Agent for Lawn and Garden Products, Inc
C/O Lynne Zahigian Regulatory Consulting
PO Box 1566
Fallon, NV 89407-1566

NOTIFICATION

DEC 06 2012

Subject Product Name Liqui-Cop
 EPA Reg No 54705-7
 Submission date 11/15/2012
 Notification for Label Change per PRN 98-10 Add Mangoes Use Restriction in
 California
 Decision Number 472450

Dear Registrant

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 11/15/2012 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have questions concerning this letter, please contact Banza Djapao at 703-305-7269, or via email at djapao_banza@epa.gov or you may call me at 703-308-9443.

Sincerely,

A handwritten signature in black ink that reads "Tony Kish".

Tony Kish
Product Manager 22
Fungicide Branch
Registration Division (7504P)

2/18



United States
Environmental Protection Agency
Washington DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1 Company/Product Number 54705-7	2 EPA Product Manager Tony Kish	3 Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4 Company/Product (Name) LIQUI-COP	PM# 22	
5 Name and Address of Applicant (Include ZIP Code) Lawn and Garden Products Inc c/o Lynne Zahigian Regulatory Consulting P O Box 1566 Fallon NV 89407-1566 <input type="checkbox"/> Check if this is a new address	6 Expedited Review In accordance with FIFRA Section 3(c)(3) (b)(i) my product is similar or identical in composition and labeling to EPA Reg No 10465-3 Product Name COPPER-COUNT-N	

Section - II

<input type="checkbox"/> Amendment Explain below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application
<input checked="" type="checkbox"/> Notification Explain below	<input type="checkbox"/> Other Explain below

Explanation Use additional page(s) if necessary (For section I and Section II)


Notification to add the statement Not for use in California to Mangoes See cover letter for explanation
This notification is consistent with the guidance in PR Notice 98 10 and the requirements of EPA regulations at 40 CFR 156 10 and 40 CFR 152 46 and no other changes have been made to the labeling or the confidential statement of formula of this product I understand that it is a violation of 18 U S C Sec 1001 to willfully make any false statement to EPA I further understand that if this notification is not consistent with the terms of PR Notice 98 10 and the requirements of 40 CFR 156 10 and 40 CFR 152 46 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA

Section - III

1 Material This Product Will Be Packaged In				2 Type of Container	
Child Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
* Certification must be submitted		If Yes* Unit Packaging wgt	No per container	If Yes* Package wgt	No per container
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
				<input type="checkbox"/> Other (Specify) _____	
3 Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4 Size(s) Retail Container 1 pint 1 quart (32 fl oz)		5 Location of Label Directions <input checked="" type="checkbox"/> on label affixed to plastic container	
6 Manner in Which Label is Affixed to Product		<input checked="" type="checkbox"/> Lithograph Paper glued Stenciled		<input type="checkbox"/> Other _____	

Section - IV

1 Contact Point (Complete items directly below for identification of individual to be contacted if necessary to process this application)		
Name Lynne C Zahigian	Title Agent	Telephone No (Include Area Code) (775) 423 9122
2 Signature 		3 Title Agent
4 Typed Name Lynne C Zahigian		5 Date 11/15/12
Certification I certify that the statements I have made on this form and all attachments thereto are true accurate and complete I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law		6 Date Application Received (Stamped)



3/18

11/26

November 15 2012

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
Registration Division
U S Environmental Protection Agency
1200 Pennsylvania Ave NW
Washington DC 20460

ATTENTION Tony Kish
Program Manager Team 22

SUBJECT Lawn and Garden Products Inc s
Liqui Cop (EPA Reg No 54705 7)

Dear Mr Kish

On behalf of Lawn and Garden Products Inc I am submitting the enclosed Notification to add statements to the label that are allowed under PR Notice 98 10

The proposed revisions or additions to the Master Label are as follows

- 1 On Pages 5 and 13 of the label we are adding the statement (Not for use in California) There is not currently any data on file with California for this active ingredient to be used on Mangoes

Please find enclosed the following in support of this Notification

- 1) A completed Application for Pesticide (Form 8570-1)
2) A copy of the revised Master Label with the proposed additions highlighted A copy of the revised Master Label without highlights

If you have any questions regarding this submission please contact me at (775) 423 9122

Regards

Handwritten signature of Lynne Zahigian

Lynne Zahigian
Agent for Lawn and Garden Products Inc

cc LGPI EPA/Liqui Cop Master File
LCZ Chron

Handwritten marks consisting of several rows of 'c' characters, possibly a signature or initials.

4/18

MASTER LABEL

NOTIFICATION

DEC 06 2012

LIQUI-COP

Alternate Brand Name LIQUI-COP RTS

COPPER FUNGICIDE GARDEN SPRAY

**Controls Many Diseases Attacking
Citrus, Fruits, Nuts, Vegetables and Ornamentals**

FOR RESIDENTIAL USE ONLY

ACTIVE INGREDIENT	
Copper diammonia diacetate complex*	27.15%
[Bis (acetate O) Diamminecopper]	
CAS No. 13822-80-5	
OTHER INGREDIENTS	72.85%
TOTAL	100.00%

* Metallic Copper Equivalent 8.0%
Contains 0.193 lbs Copper per quart or Contains 0.965 lbs Copper per pint

Sublabel A – Non-Commercial/Residential Uses (Not Subject to the Worker Protection Standard)

Sublabel B – Hose-End Sprayer Label for Non-Commercial/Residential Uses (Not Subject to the Worker Protection Standard)

EPA Reg. No. 54705-7

EPA Act No. 48198 CA.1



Manufactured for
LAWN AND GARDEN PRODUCTS, INC
P O Box 35000 Fresno CA 93745 (559) 499 2100
info@montereylawngarden.com



SUBLABEL A – NONCOMMERCIAL/RESIDENTIAL USES

LIQUI-COP

COPPER FUNGICIDE GARDEN SPRAY

**Controls Many Diseases Attacking
Citrus, Fruits, Nuts, Vegetables and Ornamentals**

FOR RESIDENTIAL USE ONLY

ACTIVE INGREDIENT	
Copper diammonia diacetate complex*	27 15%
[Bis (acetate O) Diamminecopper]	
CAS No 13822 80 5	
OTHER INGREDIENTS	<u>72 85%</u>
TOTAL	100 00%

* Metallic Copper Equivalent 8 0%
Contains 0 193 lbs Copper per quart or Contains 0 0965 lbs Copper per pint

EPA Reg No 54705-7

EPA Est No 48498 CA 1

NET CONTENTS 1 Pint or 1 Quart



Manufactured for
LAWN AND GARDEN PRODUCTS, INC
P O Box 35000 Fresno CA 93745 (559) 499 2100
info@montereylawngarden.com

KEEP OUT OF REACH OF CHILDREN		
CAUTION		c c c c c c
FIRST AID		c c c c c c
IF IN EYES	Hold eye open and rinse slowly and gently with water for 15 20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing eye Call a poison control center or doctor for treatment advice	c c c c c c c c c c c c
IF ON SKIN OR CLOTHING	Take off contaminated clothing Rinse skin immediately with plenty of water for 15 20 minutes Call a poison control center or doctor for treatment advice	c c c c c c c c c c c c
IF SWALLOWED	Call poison control center or doctor immediately for treatment advice Have person sip a glass of water if able to swallow Do not induce vomiting unless told to do so by the poison control center or doctor Do not give anything by mouth to an unconscious person	c c c c c c c c c c c c
HOT LINE NUMBER		
Have the product container or label with you when calling a poison control center or doctor or going for treatment You may also contact the National Pesticide Information Center at 1 800 858 7378 for emergency medical treatment information		

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION Causes eye and skin irritation Harmful if swallowed absorbed through the skin or inhaled Avoid contact with skin eyes or clothing Avoid breathing vapor or spray mist Wear protective eyewear (goggles face shield or safety glasses) long sleeved shirt and long pants shoes and socks and chemical resistant gloves when handling this product Wash thoroughly with soap and water after handling and before eating drinking or using tobacco Remove contaminated clothing and wash clothing before reuse Do not use food utensils such as teaspoons or tablespoons for food purposes after use with pesticides

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may contaminate water through runoff To protect the environment do not allow pesticide to enter or run off into storm drains drainage ditches gutters or surface waters Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

Do not apply this product in a way that will contact adults children or pets either directly or through drift Do not allow adults children or pets to enter the treated area until sprays have dried

Shake well before using

INFORMATION

The control of diseases with fungicides is based on PREVENTION Plant surfaces must be completely covered with the fungicide to successfully prevent infection Copper pesticides are considered preventative not curative of plant diseases Use the highest indicated rate when disease incidence is high or expected to be depending on rainfall and temperature The lower rate is suitable for general preventive sprays under normal conditions Since weather conditions and disease incidence vary consult your Agricultural Extension Service for timing and initial application

PRODUCT INSTRUCTIONS

Seek advice on identifying specific disease problems and solutions from your local Agricultural Extension Agent or other knowledgeable persons

Partially fill the spray tank/container with water Add the desired amount of LIQUI COP and continue filling the tank/container If applied with other products add LIQUI COP last Agitate the tank during mixing and application until tank is empty A plastic or metal stirring stick should provide adequate agitation Observe all cautions and limitations on labeling of all products used in mixtures Start with clean equipment Equipment should be flushed well with water after use

APPLICATION

Fruits and Nuts Mix specified rate in 1 gallon of water and apply 3 gallons of mixed solution to a small tree or bush 6 gallons of mixed solution to a medium size tree or 9 gallons of mixed solution to a large tree Thoroughly spray tree to point of runoff including upper and lower surfaces of foliage Do not overspray Do not mix more spray solution than needed

Vegetables Mix specified rate in 1 gallon of water and apply 2 gallons of mixed solution per 1000 sq ft (1 gallon of mixed solution per 500 sq ft)

Miscellaneous For herbs Mix specified rate in 1 gallon of water and apply 2 gallons of mixed solution per 1000 sq ft (1/2 gallon of mixed solution per 250 sq ft) For trees Mix specified rate in 1 gallon of water and apply 3 gallons of mixed solution to a small tree 6 gallons of mixed solution to a medium size tree or 9 gallons of mixed solution to a large tree Thoroughly spray tree to point of runoff including upper and lower surfaces of foliage Do not overspray Do not mix more spray solution than needed

NOTE This product may be reactive on metal and masonry surfaces such as galvanized roofing AVOID CONTACT with metal surfaces Do not spray on cars houses lawn furniture etc

FRUITS AND NUTS

FRUITS AND NUTS	DISEASE CONTROLLED	RATE / GALLON OF WATER	INSTRUCTIONS
ALMONDS	Brown Rot	4 – 6 tsp	Apply at delayed dormant bud swell stage Dormant oil may be used If additional applications are needed apply at no less than 7 day intervals
	Shot Hole	4 – 6 tsp	Apply as a dormant spray at leaf fall to protect buds and shoots from infection during rainy periods Reapply every 3 to 4 weeks up to late bud swell Do not apply at less than 7 day intervals Do not apply after full bloom
	NOTE Do not apply more than 410 tsp of LIQUI COP (0 41 lbs metallic copper) to Almonds per 1 000 sq ft per year		
APPLES	Anthraco nose	4 – 5 tsp	Apply as a dormant spray once to foliage after harvest
	Apple Scab (Black Spot) Bacterial Canker Blossom and Shoot Blast	4 – 6 tsp	Apply as a dormant spray once post harvest before fall rains
	Fire Blight	1/2 – 1 tsp	Apply at 10% bloom and repeat at no less than 5 – 7 day intervals during the bloom period Do not use on copper sensitive varieties
		4 – 6 tsp	Apply once as a full cover spray between silver tip and green tip Do not apply when green tip reaches 1/2 inch as injuries may occur
NOTE Do not apply more than 370 tsp of LIQUI COP (0 37 lbs metallic copper) to Apples per 1 000 sq ft per year			
APRICOTS	Brown Rot Blossom Blight Shot Hole	4 – 6 tsp	Apply as a dormant spray at leaf fall to protect buds and shoots from infection during rainy periods Reapply up to late bud swell at no less than 7 day intervals Do not apply after full bloom
	NOTE Do not apply more than 410 tsp of LIQUI COP (0 41 lbs metallic copper) to Apricots per 1 000 sq ft per year		
AVOCADOS	Anthraco nose	4 tsp	Apply when the flower buds begin to swell and continue at monthly intervals until August
	NOTE Do not apply more than 430 tsp of LIQUI COP (0 43 lbs metallic copper) to Avocados per 1 000 sq ft per year		
BANANAS	Sigatoka	1 – 2 tsp	Apply every 3 – 4 weeks
	Black Pitting	2 5 fl oz	Mix in 1 gallon of water Apply directly to the fruit stem and include the basal portion of the leaf crown Apply during the first and second weeks after fruit emergence Do not apply at less than 7 days
	NOTE Do not apply more than 71 6 fl oz of LIQUI COP (430 tsp / 0 43 lbs metallic copper) to Bananas per 1 000 sq ft per year		
BLUEBERRIES	Bacterial Canker	2 – 5 tsp	Apply before fall rains and again 4 weeks later
	Cane Canker	2 – 5 tsp	Apply before fall rains and again 4 weeks later In the spring during wet weather apply at 10 14 day intervals beginning at leaf emergence
	NOTE Do not apply more than 190 tsp of LIQUI COP (0 19 lbs metallic copper) to Blueberries per 1 000 sq ft per year		
CANE BERRIES	Anthraco nose Leaf and Cane Spot Purple Blotch Yellow Rust	1 – 3 tsp	Apply when leaf buds open Repeat when flower buds show white and continue at 10 14 day intervals Do not apply at less than 7 day intervals
	Anthraco nose Bacterial Blight Leaf and Cane Spot Purple Blotch Yellow Rust	4 – 5 tsp	Apply in the fall after harvest
	NOTE Do not apply more than 230 tsp of LIQUI COP (0 23 lbs metallic copper) to Caneberries per 1 000 sq ft per year		
CHERRIES	Deadbud Coryneum Blight	2 fl oz	Apply as a dormant spray before heavy fall rains and in January Do not apply at less than 7 day intervals
	Brown Rot Blossom Blight	4 – 6 tsp	Apply as a full cover spray at popcorn stage and at full bloom If additional applications are needed Do not apply at less than 7 day intervals
	NOTE Do not apply more than 68 3 fl oz of LIQUI COP (399 36 tsp / 0 41 lbs metallic copper) to Cherries per 1 000 sq ft per year		
CITRUS	Greasy Spot Melanose Pink Pitting Scab	2 – 4 tsp	Apply as pre bloom and post bloom spray Do not apply at less than 7 day intervals
	Brown Rot	1 – 3 tsp	Apply in the fall before or just after heavy rains In areas of skirt sprays apply to a height of at least 4 feet (Skirt sprays are applications up to 4 ft high on the tree The whole tree is not sprayed) Do not apply at less than 7 day intervals
	NOTE Do not apply more than 290 tsp of LIQUI COP (0 29 lbs metallic copper) to Citrus per 1 000 sq ft per year		
CURRANTS GOOSEBERRIES	Anthraco nose Leaf Spot (Cane Blight)	3 – 5 tsp	Make 3 applications starting after harvest before bloom and after petal fall Do not make applications at less than 10 day intervals
	NOTE Do not apply more than 230 tsp of LIQUI COP (0 23 lbs metallic copper) to Currants/Gooseberries per 1 000 sq ft per year		

FILBERTS (Permitted only in Washington State and Oregon)	Bacterial Blight	4 – 6 tsp	Apply after harvest Under severe conditions apply again when / of the leaves have dropped Do not apply at less than 14 day intervals
	Eastern Filbert Blight	4 – 6 tsp	Make initial application after harvest in October before heavy rains begin The next application should be made in late February to early March followed by another application 1 month later Use higher rate when rainfall is heavy and disease pressure is high
	NOTE Do not apply more than 550 tsp of LIQUI COP (0 55 lbs metallic copper) to Filberts per 1 000 sq ft per year		
GRAPES	Anthraxnose Black Rot Downy Mildew Powdery Mildew	1 tsp	Apply just before bud break when the shoots are 6 – 8 inches long just after bloom and every 4 – 10 days throughout the season as needed Foliar injury may occur on copper sensitive varieties
	NOTE Do not apply more than 460 tsp of LIQUI COP (0 46 lbs metallic copper) to Grapes per 1 000 sq ft per year		
KIWI	<i>Pseudomonas syringae</i> <i>Erwinia herbicola</i> <i>Pseudomonas fluorescens</i>	4 tsp	Apply in 4 gallons of water per 1 000 sq ft Make applications on a monthly basis A maximum of 3 applications may be made
	NOTE Do not apply more than 140 tsp of LIQUI COP (0 14 lbs metallic copper) to Kiwi per 1 000 sq ft per year		
LIMES	Greasy Spot	4 tsp	Apply in June and continue at monthly intervals through August
	NOTE Do not apply more than 290 tsp of LIQUI COP (0 29 lbs metallic copper) to Limes per 1 000 sq ft per year		
MANGOS (Not for use in California)	Anthraxnose	4 tsp	Apply at 30 day intervals from the time the panicles are 2 inches in length until all fruits are set until August
	NOTE Do not apply more than 145 tsp of LIQUI COP (0 145 lbs metallic copper) to Mangos per 1 000 sq ft per year		
OLIVES	Peacock Spot	2 – 4 tsp	Make first application before winter rains fall A second application should be made in early spring if disease is severe If additional applications are necessary apply at no less than 30 day intervals
	NOTE Do not apply more than 145 tsp of LIQUI COP (0 145 lbs metallic copper) to Olives per 1 000 sq ft per year		
PEACHES NECTARINES	Bacterial Spot	4 tsp	Apply as a dormant spray at no less than 7 day intervals
	Blossom Brown Rot Leaf Curl Shot Hole	4 – 6 tsp	Apply as a dormant or delayed dormant spray to protect buds and shoots from infection during rainy period Reapply up until late bud swell Do not apply at less than 7 day intervals Do not apply after full bloom
	NOTE Do not apply more than 410 tsp of LIQUI COP (0 41 lbs metallic copper) to Peaches/Nectarines per 1 000 sq ft per year		
PEARS QUINCE	Fire Blight	/ 1 tsp	Apply at 10% bloom and repeat at no less than 5 7 day intervals throughout the bloom period Do not use on copper sensitive varieties
	Blossom Blast	4 – 6 tsp	On/ one application permitted in fall late dormant season
	NOTE Do not apply more than 370 tsp of LIQUI COP (0 37 lbs metallic copper) to Pears/Quince per 1 000 sq ft per year		
PECANS	Shuck and Kernel Rot Zonate Leaf Spot	2 – 5 tsp	For suppression apply in sufficient water to ensure complete spray coverage at 2 – 4 week intervals (no less than 14 days) starting at kernel growth and continuing until shucks open Use the higher rate and shorter interval if frequent rainfall occurs
	NOTE Do not apply more than 190 tsp of LIQUI COP (0 19 lbs metallic copper) to Pecans per 1 000 sq ft per year		
PISTACHIOS	Alternaria Late Blight	4 – 5 tsp	Apply at 50% and full bloom followed by up to 3 applications at 30 day intervals
	Botrytis Blight Botryosphaeria Panicle Shoot Blight Septoria Leaf Blight	4 – 5 tsp	Make initial application at bud swell and repeat on a 14 – 28 day schedule as dictated by disease conditions If disease conditions are severe use the higher rate and shorter interval
	NOTE Do not apply more than 190 tsp of LIQUI COP (0 19 lbs metallic copper) to Pistachios per 1 000 sq ft per year		
PLUMS PRUNES	Bacterial Blast Bacterial Canker Brown Rot Blossom Blight Coryneum Blight (Shot Hole)	4 – 6 tsp	Apply as a dormant spray at early green bud to full popcorn states Do not apply at less than 7 day intervals
	NOTE Do not apply more than 410 tsp of LIQUI COP (0 41 lbs metallic copper) to Plums/Prunes per 1 000 sq ft per year		
STRAWBERRIES	Leaf Spot Scorch	3 – 4 tsp	Apply at 7 – 10 day intervals from the time new growth starts until harvest
	NOTE Do not apply more than 190 tsp of LIQUI COP (0 19 lbs metallic copper) to Strawberries per 1 000 sq ft per year		
WALNUTS	Walnut Blight	4 – 6 tsp	Make first application at early pre bloom Make second application at late pre bloom Make additional applications at no less than 7 day intervals if disease conditions persist
	NOTE Do not apply more than 580 tsp of LIQUI COP (0 58 lbs metallic copper) to Walnuts per 1 000 sq ft per year		

* Delayed Dormant – The period just before blossom buds break open

9/18

VEGETABLES

VEGETABLES	DISEASE CONTROLLED	RATE / GALLON OF WATER	INSTRUCTIONS
BEANS PEAS LENTILS (Succulent and Dry)	Bacterial Blight (Halo and Common)	2 – 6 tsp	Apply when plants are 3 – 5 inches high and before diseases appear Repeat at 7 – 10 day intervals Apply at 7 day intervals if disease pressure is severe
	NOTE Do not apply more than 30 tsp of LIQUI COP (0 03 lbs metallic copper) to Beans/Peas/Lentils per 1 000 sq ft per year		
BEETS	Cercospora Leaf Spot	3 – 6 tsp	Apply when disease appears making 3 to 6 sprays at 10 14 day intervals Apply at 10 day intervals if disease pressure is severe
	NOTE Do not apply more than 110 tsp of LIQUI COP (0 11 lbs metallic copper) to Beets per 1 000 sq ft per year		
CARROTS	Early Blight Late Blight	4 – 6 tsp	Apply when plants are 6 high Make 3 to 5 applications at 7 – 10 day intervals
	NOTE Do not apply more than 120 tsp of LIQUI COP (0 12 lbs metallic copper) to Carrots per 1 000 sq ft per year		
CELERY	Bacterial Blight Early Blight Late Blight	4 – 6 tsp	Apply as soon as plants are established and repeat 7 day intervals
	NOTE Do not apply more than 120 tsp of LIQUI COP (0 12 lbs metallic copper) to Celery per 1 000 sq ft per year		
CORN (Pop Sweet)	Bacterial Rot Bacterial Stripe Bacterial Wilt Leaf Blight Stalk Rot	4 tsp	Apply when disease appears and repeat as necessary at 7 day intervals
	NOTE Do not apply more than 100 tsp of LIQUI COP (0 1 lbs metallic copper) to Corn per 1 000 sq ft per year		
CRUCIFERS [Broccoli Brussels Sprouts Cabbage Cauliflower Greens (Collard Mustard and Turnip)]	Blackleaf Spot Black Rot	2 – 6 tsp	Apply when disease appears and repeat at 7 10 day intervals
	Downy Mildew	1 – 2 tsp	Apply when disease appears and repeat at 7 10 day intervals
	NOTE Do not apply more than 60 tsp of LIQUI COP (0 06 lbs metallic copper) to Crucifers per 1 000 sq ft per year		
CUCURBITS (Cantaloupe Cucumber Honeydew Muskmelon Pumpkin Squash Watermelon)	Alternaria Leaf Spot Angular Leaf Spot Anthracnose Downy Mildew Gummy Stem Blight Powdery Mildew Watermelon Bacterial Fruit Blotch	3 – 4 tsp	Apply when disease appears and repeat 7 10 day intervals
	NOTE Do not apply more than 115 2 tsp (0 12 lbs metallic copper) to Cucurbits per 1 000 sq ft per year		
EGGPLANT	Alternaria Blight Anthracnose Phomopsis	4 tsp	Apply before disease appears and repeat at 7 10 day intervals
	NOTE Do not apply more than 180 tsp of LIQUI COP (0 18 lbs metallic copper) to Eggplant per 1 000 sq ft per year		
ONIONS	Downy Mildew Purple Blotch	4 tsp	Apply when plants are 4 6 inches high and repeat at 7 10 day intervals
	NOTE Do not apply more than 140 tsp of LIQUI COP (0 14 lbs metallic copper) to Onions per 1 000 sq ft per year		
PEANUTS	Cercospora Leaf Spot	3 – 6 tsp	Apply on first appearance of disease and repeat at no less than 10 14 day intervals
	Pod Rot Complex (Pythium mynotylum Rhizoctonia solani and Sclerotium rolfsii)	2 5 fl oz	Apply at pegging in a 12 15 inch band over the row If repeat application is necessary do not apply at less than 7 day intervals
	NOTE Do not apply more than 18 3 fl oz of LIQUI COP (110 tsp / 0 11 lbs metallic copper) to Peanuts per 1 000 sq ft per year		
PEPPERS	Bacterial Spot Cercospora Leaf Spot	3 6 tsp	Make first application upon emergence of seedlings or immediately after transplanting and repeat at 7–10 day intervals When disease is severe apply at 7 day intervals NOTE Disease control is crucial during fruiting
	NOTE Do not apply more than 270 tsp of LIQUI COP (0 27 lbs metallic copper) to Peppers per 1 000 sq ft per year		
POTATOES	Early Blight Late Blight	3 6 tsp	Apply on first appearance of disease and repeat at 7 10 day intervals
	NOTE Do not apply more than 570 tsp of LIQUI COP (0 57 lbs metallic copper) to Potatoes per 1 000 sq ft per year		
SPINACH	Anthracnose Cercospora Leaf Spot Downy Mildew	3 tsp	Apply on first appearance of disease and repeat at 7 10 day intervals
	NOTE Do not apply more than 90 tsp of LIQUI COP (0 09 lbs metallic copper) to Spinach per 1 000 sq ft per year		
TOMATOES	Bacterial Speck Bacterial Spot Early Blight Late Blight	3 – 5 tsp	Make first application upon emergence of seedlings or immediately after transplanting and repeat at 7–10 day intervals When disease pressure is severe apply at the shorter spray interval Complete coverage is essential for disease control NOTE While the labeled rate is particularly effective against Bacterial Spot a tank mix of

10/18

			Maneb or Mancozeb used at the labeled rates controls a broad range of diseases
NOTE Do not apply more than 400 tsp of LIQUI COP (0.4 lbs metallic copper) to Tomatoes per 1 000 sq ft per year			

MISCELLANEOUS

MISCELLANEOUS	DISEASE CONTROLLED	RATE / GALLON OF WATER	INSTRUCTIONS
CARAMBOLA	Anthracnose	6 tsp	Make initial application just before flowering and repeat on a weekly schedule until just before harvest
	Do not apply more than 240 tsp of LIQUI COP (0.24 lbs metallic copper) to Carambola per 1 000 sq ft per year		
CHIVES	Downy Mildew	4 tsp	Begin applications when plants are first established Repeat applications every 7 - 10 days as dictated by disease conditions If disease pressure is high use the shorter spray interval
	NOTE Do not apply more than 60 tsp of LIQUI COP (0.06 lbs metallic copper) to Chives per 1 000 sq ft per year		
DILL	Phoma Leaf Spot Rhizoctonia Foliage Blight	6 tsp	Begin applications when plants are first established and repeat at 7 - 10 day intervals depending upon disease severity and environmental conditions If disease pressure is high use the shorter spray interval
	NOTE Do not apply more than 90 tsp of LIQUI COP (0.09 lbs metallic copper) to Dill per 1 000 sq ft per year		
DOUGLAS FIR	Rhabdocline Needlecast	4 tsp	Begin applications at bud break and repeat at 3 - 4 week intervals Apply in a tank mix with another registered pesticide if moderate to severe disease pressure is present
	NOTE Do not apply more than 460 tsp of LIQUI COP (0.46 lbs metallic copper) to Douglas Fir per 1 000 sq ft per year		
GUAVA	Anthracnose Red Algae	6 tsp	Make initial application just before flowering and repeat on a weekly schedule until just before harvest
	NOTE Do not apply more than 110 tsp of LIQUI COP (0.11 lbs metallic copper) to Guava per 1 000 sq ft per year		
LITCHI	Anthracnose	6 tsp	Make initial application just before flowering and repeat on a weekly schedule until just before harvest
	NOTE Do not apply more than 110 tsp of LIQUI COP (0.11 lbs metallic copper) to Litchi per 1 000 sq ft per year		
LIVE OAK	Ball Moss	2 fl oz	Apply in the spring when ball moss is actively growing using 1.5 gallons of spray per foot of tree height Make sure to wet ball moss tufts thoroughly A second application may be required after 12 months NOTE LIQUI COP may be injurious to ornamentals grown under Live Oaks This product may be reactive on metal and masonry surfaces such as galvanized roofing Avoid contact with metal surfaces Do not spray on cars houses lawn furniture etc
	NOTE Do not apply more than 76.7 fl oz of LIQUI COP (0.46 lbs metallic copper) to Live Oak per 1 000 sq ft per year		
MACADAMIA	Anthracnose	3 tsp	Initiate sprays at first sign of flowering and repeat on a weekly schedule until just before harvest
	Phytophthora Blight Raceme Blight	2 - 3 tsp	Apply during raceme development and bloom periods Do not apply at less than 7 day intervals
	NOTE Do not apply more than 220 tsp of LIQUI COP (0.22 lbs metallic copper) to Macadamia per 1 000 sq ft per year		
PAPAYA	Anthracnose	2 - 5 tsp	Begin applications before disease appears and repeat at 10 - 14 day intervals Use higher rates when conditions favor disease
	NOTE Do not apply more than 490 tsp of LIQUI COP (0.49 lbs metallic copper) to Papaya per 1 000 sq ft per year		
PARSLEY	Bacterial Blight	6 tsp	Begin applications when plants are first established and repeat at 10 day intervals
	NOTE Do not apply more than 50 tsp of LIQUI COP (0.05 lbs metallic copper) to Parsley per 1 000 sq ft per year		
PASSION FRUIT	Anthracnose	3 tsp	Make initial application just before flowering and repeat on a weekly schedule until just before harvest
	NOTE Do not apply more than 220 tsp of LIQUI COP (0.22 lbs metallic copper) to Passion Fruit per 1 000 sq ft per year		
SUGAR APPLE (Annona)	Anthracnose	4 - 6 tsp	Make initial application just before flowering and repeat on a weekly schedule until just before harvest
	NOTE Do not apply more than 290 tsp of LIQUI COP (0.29 lbs metallic copper) to Sugar Apple per 1 000 sq ft per year		
SYCAMORE	Anthracnose	4 - 6 tsp	Apply as a full coverage spray Apply in 1-gallon of water or sufficient volume for thorough coverage Make first application at bud break and second application 7 - 10 days later after leaf expansion
	NOTE Do not apply more than 460 tsp of LIQUI COP (0.46 lbs metallic copper) to Sycamore per 1 000 sq ft per year		

ORNAMENTALS

For use in container landscape and outdoor ornamental plantings LIQUI COP can also be used on ornamentals in home greenhouses and shadehouses

Apply as a thorough coverage spray using 2 teaspoons of LIQUI COP per gallon of water Begin application at first sign of disease and repeat at 7 - 14 day intervals as needed Use shorter interval during periods of frequent rains or

when severe disease conditions persist Do not apply more than 460 tsp of LIQUI COP (0.46 lbs metallic copper) per 1 000 sq ft per year

Notice to User Plant sensitivities to LIQUI COP have been found to be acceptable in specific genera and species listed on this label however it is impossible to know sensitivities under all conditions and phytotoxicity may occur Due to the large numbers of species and varieties of ornamentals it is impossible to test every one for sensitivity to LIQUI COP Neither the manufacturer or seller recommends use upon species not listed on the label nor has it been determined that LIQUI COP can safely be used on ornamental plants not listed on this label The user should determine if LIQUI COP can be used safely prior to use

PLANT	DISEASE CONTROLLED
Althea (Rose of Sharon)	Bacterial leaf spot
Aralia	Alternaria Cercospora leaf spot Xanthomonas leaf spot
Arborvitae	Alternaria twig blight Cercospora leaf blight
Azalea ⁽¹⁾	Botrytis blight Cercospora leaf spot Phytophthora dieback Powdery mildew
Begonia	Bacterial leaf spot (Erwinia sp Pseudomonas sp Xanthomonas sp)
Bougainvillea	Anthraxnose Bacterial leaf spot
Bulbs (Tulip Gladiolus)	Anthraxnose Botrytis blight
Camellia	Anthraxnose Bacterial leaf spot
Camphor tree	Pseudomonas leaf spot
Canna	Pseudomonas leaf spot
Carnation ⁽¹⁾	Alternaria blight Botrytis blight Pseudomonas leaf spot
Chinese tallow tree	Bacterial leaf spot (Pseudomonas sp Xanthomonas sp)
Chrysanthemum ⁽¹⁾	Botrytis blight Septoria leaf spot
Cotoneaster	Botrytis blight
Dahlia	Alternaria leaf spot Botrytis gray mold Cercospora leaf spot
Date palm	Pestalotia leaf spot
Dianthus	Bacterial soft rot Bacterial spot
Dogwood	Anthraxnose
Dusty miller	Bacterial leaf spot (Pseudomonas cichorii)
Echinacea	Bacterial leaf spot (Pseudomonas chiconi)
Elm Drake	Xanthomonas leaf spot
Euonymus	Anthraxnose Botrytis blight
European fan palm	Pestalotia leaf spot
Gardenia	Alternaria leaf spot Botrytis bud rot Cercospora leaf spot
Geranium	Alternaria leaf spot Botrytis gray mold Cercospora leaf spot
Gladiolus	Alternaria leaf spot Bacterial leaf blight Botrytis gray mold
Goldenrain tree	Bacterial leaf spot
Hibiscus	Bacterial leaf spot
Holly fern	Pseudomonas leaf spot
Impatiens	Bacterial leaf spot
Ivy (English Algenan) ⁽¹⁾	Xanthomonas leaf spot
Ixora	Xanthomonas leaf spot
Juniper (Eastern red cedar)	Anthraxnose
Lantana	Bacterial leaf spot
Lilac	Cercospora leaf spot
Loblolly bay	Anthraxnose
Loquat	Colletotrichum sp Entomosporium maculate
Magnolia (Saucer)	Bacterial leaf spot
Magnolia (Southern)	Algal leaf spot Anthracnose Bacterial leaf spot
Magnolia (Sweet bay)	Anthraxnose
Mandevillas	Anthraxnose
Mangold	Alternaria leaf spot Botrytis leaf and flower rot Cercospora leaf spot
Mulberry weeping	Bacterial leaf spot
Oak laurel	Algal leaf spot (Cephaleuros virescens)
Oleander	Bacterial leaf spot Fungal leaf spot
Pachysandra	Volutella leaf blight
Pansy	Downy mildew
Pear (Flowering)	Fireblight Leaf spot

12/18

Pentas (Egyptian star)	Bacterial leaf spot (Xanthomonas sp)
Peony	Botrytis blight
Periwinkle	Phomopsis stem blight
Philodendron	Bacterial leaf spot
Phlox	Alternaria leaf spot
Photinia	Anthracoise Entomosporium
Pistachio	Anthracoise
Plantain lily	Bacteria leaf spot
Powder puff plant	Bacterial leaf spot
Pyracantha	Fireblight scab
Queen palm	Exosporium leaf spot Phytophthora bud rot
Rhododendron	Alternaria flower spot
Rose ⁽¹⁾	Black spot Powdery mildew
Verbena	Xanthomonas leaf spot
Viburnum	Anthracoise
Washingtonia palm	Pestalotia leaf spot
Weeping willow	Anthracoise
Yucca (Adam s needle)	Cercospora leaf spot Septoria leaf spot

⁽¹⁾ Discoloration of foliage and/or blooms has been noted on some varieties

TURFGRASS

To control algae in ornamental lawns and turf apply 10 fluid ounces (0.06 lbs metallic copper) of LIQUI COP in 5 gallons of water per 1 000 square feet LIQUI COP may be used alone or in combination with other registered fungicides as a maintenance spray Observe all precautions and limitations on the label of each product used in tank mixes Do not apply more than 80 fl oz (0.48 lbs metallic copper) of LIQUI COP per 1 000 sq ft per year Do not apply at less than 10 day intervals

NOTE Phytotoxicity may occur depending upon varietal differences Apply the recommended rate to a small area and observe for 7 to 10 days for signs of injury If phytotoxicity occurs discontinue use Do not apply in a spray solution with a pH of less than 6.5

FROST INJURY PROTECTION

Bacterial Ice Nucleation Inhibitor – Application of LIQUI COP to all plants listed on this label at rates indicated just prior to anticipated frost conditions will sustain control of ice nucleating bacteria (*Pseudomonas syringae*, *Erwinia herbicola* and *Pseudomonas fluorescens*) and may therefore provide protection against light frost

Not recommended for those geographic areas where weather conditions favor severe frost

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage or disposal

PESTICIDE STORAGE Store in a secure locked area away from pets and out of the reach of children Store only in original container and place in a locked storage area

PESTICIDE DISPOSAL/CONTAINER HANDLING Nonrefillable container Do not reuse or refill this container If empty Place in trash or offer for recycling if available If partly filled Call your local solid waste agency for disposal instructions Never place unused product down any indoor or outdoor drain

IMPORTANT READ BEFORE USE

Read the entire Directions for Use Conditions Disclaimer of Warranties and Limitations of Liability before using this product If terms are not acceptable return the unopened product container at once

By using this product user or buyer accepts the following Conditions Disclaimer of Warranties and Limitations of Liability

CONDITIONS The Directions for Use of this product are believed to be adequate and must be followed carefully However it is impossible to eliminate all risks associated with the use of the product Crop injury ineffectiveness or other unintended consequences may result because of such factors as weather conditions presence of other materials or the manner of use of application all of which are beyond the control of Lawn and Garden Products Inc To the extent consistent with applicable law all such risks shall be assumed by the user or buyer


DISCLAIMER OF WARRANTIES To the extent consistent with applicable law Lawn and Garden Products Inc makes no other warranties express or implied of merchantability or of fitness for a particular purpose or otherwise that extend beyond the statements made on this label No agent of Lawn and Garden Products Inc is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein To the extent consistent with applicable law Lawn and Garden

13/
18

Products Inc disclaims any liability whatsoever for special incidental or consequential damages resulting from the use or handling of this product

LIMITATIONS OF LIABILITY To the extent consistent with applicable law the exclusive remedy of the user or buyer for any and all losses injuries or damages resulting from the use or handling of this product whether in contract warranty tort negligence strict liability or otherwise shall not exceed the purchase price paid or at Lawn and Garden Products Inc s election the replacement of product

BATCH CODE _____



14/18

Sublabel B – READY-TO-SPRAY LABEL
FOR NON-COMMERCIAL/RESIDENTIAL USES

LIQUI-COP RTS

COPPER FUNGICIDE GARDEN SPRAY

Controls Many Diseases Attacking
Citrus and Fruit & Nut Trees

Ready To Spray Bottle
Just Attach to Garden Hose & Spray

Delivers 4 Tsp of LIQUI-COP RTS per Gallon of Water
Makes up to 48 Gallons of Spray Solution

FOR RESIDENTIAL USE ONLY

ACTIVE INGREDIENT	
Copper diammonia diacetate complex*	27 15%
[Bis (acetato O) Diamminocopper]	
CAS No 13822 80 5	
OTHER INGREDIENTS	<u>72 85%</u>
TOTAL	100 00%

* Metallic Copper Equivalent 8 0%
Contains 0 193 lbs Copper per quart

EPA Reg No 54705 7

EPA Est No 48498 CA 1

NET CONTENTS 1 Quart (32 FL OZ)



Manufactured for
LAWN AND GARDEN PRODUCTS, INC
P O Box 35000 Fresno CA 93745 (559) 499 2100
info@montereylawngarden.com

KEEP OUT OF REACH OF CHILDREN	
CAUTION	
FIRST AID	
IF IN EYES	Hold eye open and rinse slowly and gently with water for 15 20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing eye Call a poison control center or doctor for treatment advice
IF ON SKIN OR CLOTHING	Take off contaminated clothing Rinse skin immediately with plenty of water for 15 20 minutes Call a poison control center or doctor for treatment advice
IF SWALLOWED	Call poison control center or doctor immediately for treatment advice Have person sip a glass of water if able to swallow Do not induce vomiting unless told to do so by the poison control center or doctor Do not give anything by mouth to an unconscious person

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact the National Pesticide Information Center at 1 800 858 7378 for emergency medical treatment information.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION Causes skin and eye irritation. Harmful if swallowed, absorbed through the skin or inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Wear protective eyewear (goggles, face shield, or safety glasses), long-sleeved shirt and long pants, shoes and socks, and chemical-resistant gloves when handling this product. Wash thoroughly with soap and water after handling and before eating, drinking, or using tobacco. Remove contaminated clothing and wash clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may contaminate water through runoff. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact adults, children, or pets, either directly or through drift. Do not allow adults, children, or pets to enter treated area until sprays have dried.

The control of fungal disease is based on PREVENTION. Copper pesticides are considered preventative, not curative, of plant diseases. Plant surfaces must be completely covered with fungicide to successfully prevent infection.

Seek advice on identifying specific disease problems and solutions from your local Agricultural Extension Agent or other knowledgeable persons.

INSTRUCTIONS

SHAKE WELL BEFORE USING

APPLICATION Thoroughly spray tree or bush to the point of runoff, including upper and lower surfaces of any foliage. Do not overspray. Do not apply at less than 7 day intervals.


NOTE This product may be reactive on metal and masonry surfaces such as galvanized roofing. AVOID contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.

HOW TO USE THIS SPRAYER Shake Container Well Before Getting Started

- 1 Before you start, the blue top water flow valve should be in the OFF position and the red product flow valve should be in the up OFF position.
- 2 Shake container well and attach it to your garden hose.
- 3 Turn on the water from the faucet.
- 4 To apply, remove side pin (labeled PULL OUT) and rotate red product flow valve to ON, then while holding sprayer at waist level and pointing in a direction away from face and body, push blue water flow valve forward to activate water.


NOTE LIQUI COP RTS is a dark blue liquid. When the spray turns clear, it is time to change to a new bottle of product.

- 5 When you are finished spraying or if you have to stop spraying at any time, press blue water flow valve with thumb to the rear to shut off water and return side red valve to the upright OFF position and replace side pin.
- 6 Turn off the water at the faucet.



16/18

7 Remove the container from the garden hose then rinse thoroughly and store according to storage instructions



FRUITS AND NUTS

FRUITS AND NUTS	DISEASE CONTROLLED	INSTRUCTIONS
ALMONDS	Brown Rot	Apply at delayed dormant bud swell stage
	Shot Hole	Apply as a dormant spray at leaf fall to protect buds and shoots from infection during rainy periods Reapply every 3 to 4 weeks up until late bud swell (the period in Spring before buds begin to open) Do not apply after full bloom
APPLES	Apple Scab (Black Spot) Bacterial Canker Blossom and Shoot Blast	Apply post harvest before fall rains
APRICOTS	Brown Rot Blossom Blight	Apply when buds show red color until brown bud jackets have fallen
	Shot Hole	Apply in late fall or early winter before winter rains or snow Repeat in late winter or early spring before buds break Do not apply after full bloom
AVOCADOS	Anthraco	Begin application when the flower buds begin to swell and continue at monthly intervals until August
BLUEBERRIES	Bacterial Canker	Apply before fall rains and again 4 weeks later
	Cane Canker	Apply before fall rains and again 4 weeks later In the spring during wet weather apply at 10 14 day intervals beginning at leaf emergence
CANE BERRIES	Anthraco Bacterial Blight Leaf and Cane Spot Purple Blotch Yellow Rust	Apply in the fall after harvest
CERRIES	Brown Rot Blossom Blight	Apply as a full cover spray at popcorn stage and at full bloom
CITRUS	Greasy Spot Melanose Pink Pitting Scab	Apply as pre bloom and post bloom sprays
CURRENTS GOOSEBERRIES	Anthraco Leaf Spot (Cane Blight)	Make 3 applications starting after harvest before bloom and after petal fall
LIMES	Greasy Spot	Apply in June and continue at monthly intervals through August
MANGOS (Not for use in California)	Anthraco	Apply at 30 day intervals from the time the flowers are 2 inches in length until all fruits are set
OLIVES	Peacock Spot	Make first application before winter rains fall A second application should be made in early spring if disease is severe
PEACHES NECTARINES	Bacterial Spot	Apply as a dormant spray
	Blossom Brown Rot	Apply as a dormant or delayed dormant spray Do not apply at or after full bloom
	Leaf Curl Shot Hole	Apply at leaf fall to protect buds and shoots from infection during rainy periods Reapply up until late bud swell Do not apply after full bloom
PEARS	Blossom Blast	Apply as a dormant spray Apply only at bud break to control of every infection
PECANS	Shuck and Kernel Rot Zonate Leaf Spot	For suppression apply at 2 - 4 week intervals starting at kernel growth and continuing until shucks open Use shorter interval if frequent rainfall occurs
PISTACHIOS	Alternaria Late Blight	Apply at 50% and full bloom followed by up to 3 applications at 30 day intervals
	Botrytis Blight Botryosphaeria Panicle Shoot Blight Septoria Leaf Blight	Make initial application at bud swell and repeat on a 14 - 28 day schedule as dictated by disease conditions If disease conditions are severe use the higher rate and shorter spray interval
PLUMS PRUNES	Bacterial Blast Bacterial Canker Shot Hole	Apply as a dormant spray and at early green bud to full popcorn stages
WALNUTS	Walnut Blight	Make first application at early pre bloom Make second application at late pre bloom Make additional applications if disease conditions persist at 7 day intervals

Delayed Dormant - The period just before blossom buds break open

17/18

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage or disposal

PESTICIDE STORAGE Store in a secure locked area away from pets and out of the reach of children Store only in original container and place in a locked storage area

PESTICIDE DISPOSAL/CONTAINER HANDLING Nonrefillable container Do not reuse or refill this container **If empty** Place in trash or offer for recycling if available **If partly filled** Call your local solid waste agency for disposal instructions Never place unused product down any indoor or outdoor drain

IMPORTANT READ BEFORE USE

Read the entire Directions for Use Conditions Disclaimer of Warranties and Limitations of Liability before using this product If terms are not acceptable return the unopened product container at once

By using this product user or buyer accepts the following Conditions Disclaimer of Warranties and Limitations of Liability

CONDITIONS The Directions for Use of this product are believed to be adequate and must be followed carefully However it is impossible to eliminate all risks associated with the use of the product Crop injury ineffectiveness or other unintended consequences may result because of such factors as weather conditions presence of other materials or the manner of use of application all of which are beyond the control of Lawn and Garden Products Inc To the extent consistent with applicable law all such risks shall be assumed by the user or buyer

DISCLAIMER OF WARRANTIES To the extent consistent with applicable law Lawn and Garden Products Inc makes no other warranties express or implied of merchantability or of fitness for a particular purpose or otherwise that extend beyond the statements made on this label No agent of Lawn and Garden Products Inc is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein To the extent consistent with applicable law Lawn and Garden Products Inc disclaims any liability whatsoever for special incidental or consequential damages resulting from the use or handling of this product

LIMITATIONS OF LIABILITY To the extent consistent with applicable law the exclusive remedy of the user or buyer for any and all losses injuries or damages resulting from the use or handling of this product whether in contract warranty tort negligence strict liability or otherwise shall not exceed the purchase price paid or at Lawn and Garden Products Inc s election the replacement of product




BATCH CODE _____

OPTIONAL HOSE-END SPRAYER DIRECTIONS

How To Use This Sprayer Shake container well before getting started

The sprayer attached to the container is ready to use Simply attach to your garden hose and follow these instructions

- 1 Connect sprayer to hose
- 2 Turn on water
- 3 To begin spraying point nozzle in the direction you want to spray
- 4 Bend plastic tab back and turn knob to ON position
- 5 Spray evenly to area to be treated
- 6 To stop spraying turn knob to OFF position
- 7 Turn off water
- 8 Relieve water pressure by bending plastic tab back and turning knob to ON position until water slows to a drip Then turn knob back to OFF position
- 9 Disconnect sprayer from hose



18/18

OPTIONAL HOSE-END SPRAYER DIRECTIONS

How To Use This Sprayer Shake container well before getting started

CONNECT

- 1 Shake well before using
- 2 Connect sprayer to hose
- 3 Turn on water

SPRAY

- 1 Before spraying point nozzle in the direction you want to spray
- 2 [Bend small plastic tab back and] Turn knob [clockwise] to ON position
- 3 Begin spraying

FINISH

- 1 To stop spraying [turn knob counter clockwise] to OFF position
- 2 Turn off water
- 3 Relieve water pressure by [bending plastic tab back and] turning knob to ON position until water slows to a drip Then turn knob back to OFF position
- 4 Disconnect sprayer from hose

[Diagram of Hose End Sprayer to be inserted]

