

FRONT LABEL

**ACTROL CHEMICAL PRODUCTS**

Houston, Texas 77074

AGRICULTURAL**FUNGICIDE****CHAMPION[®]**
WETTABLE POWDER**KEEP OUT OF
REACH OF CHILDREN
DANGER – PELIGRO****SEE BACK PANEL FOR
ADDITIONAL PRECAUTIONARY
STATEMENTS, AND
STATEMENT OF
PRACTICAL TREATMENT****ACTIVE INGREDIENT**

CUPRIC HYDROXIDE 77%

INERT INGREDIENTS 23%

TOTAL 100%

(METALLIC COPPER EQUIVALENT 50%)

**NET WEIGHT 20 LBS.**

EPA REG. NO. 55146-1

EPA EST. NO. 55896-SC-1

BEST AVAILABLE COPY

CAUTION: Phytotoxicity may occur from late applications. (Discontinue use when green tip is 1/2-inch). **Crown or Collar Rot.** Mix 4 pounds in 100 gallons of water. Apply 4 gallons of suspension as a trench on the lower trunk area of each tree. Apply either in early spring or in fall after harvest each year. Do not use if soil pH is below 5.5 or copper toxicity may result. (EXCEPT CALIFORNIA)

APRICOTS, *Coryneum Blight (Shot Hole) & Blossom Brown Rot.* Apply at popcorn to full bloom using 2 to 3 pounds per 100 gallons (300-400 GPA). To avoid spray injury, do not apply after bloom.

AVOCADOS, Scab. Apply when bloom buds begin to swell at 2 pounds CHAMPION WETTABLE POWDER per 100 gallons or 8-10 pounds per acre depending on equipment. Continue application at monthly intervals for 5 to 6 applications. Follow recommendations of State Agricultural Experiment Stations.

CHERRY, Dead Bud (*Pseudomonas syringae*) & *Coryneum Blight.* Apply 6 pounds plus 1 pint superior-type oil per 100 gallons in October (before heavy fall rains) and again in January. In orchards where the disease is severe, a spray should also be applied in August. **Brown Rot Blossom Blight.** For adequate control apply 2 to 3 pounds per 100 gallons as a full cover spray at popcorn and full bloom.

CITRUS, Melanose, Scab, Greasy Spot, & Pink Pitting. Apply as pre-bloom and post bloom sprays. Use 0.75 to 1 pounds per 100 gallons depending on disease severity. May be used in concentrate sprays at equivalent rates. For aerial application use 6 pounds per 10 gallons per acre. **Brown Rot.** Use 0.5 to 1 pounds per 100 gallons as dilute spray or at equivalent rate as concentrate spray depending on disease severity. Begin application in fall before or just after first heavy rains. Apply to skirts of trees to a height of at least 4 feet. Apply also to bare ground one foot beyond skirt. (California only. In areas subject to copper injury, add 1.3 to 1 pound of high quality lime per pound of product). **Citrus Canker.** Spray flushes 7-14 days after shoots begin to grow. For dilute sprays, apply 1.5 pounds per 100 gallons of water. Spray trees thoroughly. For concentrate sprays, apply at equivalent rates in sufficient water to obtain good coverage. Young fruit may require an additional application (under slight disease pressure, the early spring and late fall flushes may not need spraying). Under heavy disease pressure, each flush of new growth should be sprayed.

FILBERTS, Bacterial Blight. Apply 4 to 6 pounds plus 1 pint superior-type oil per 100 gallons in late August or early September. In seasons of heavy rainfall, apply another spray when three-fourths of the leaves have dropped.

MANGO, (Florida) Anthracnose. Apply monthly after fruit set until harvest at 2 pounds per 100 gallons or 8-10 pounds per acre depending on equipment. Consult Extension Service for State recommendations.

OLIVES, (California) Peacock Spot. Make first application at 2 to 3 pounds per 100 gallons or at 8-12 pounds per acre depending on equipment before winter rains fall. A second application in early spring should be made if disease is severe.

PEACHES & NECTARINES, Leaf Curl & *Coryneum Blight (Shot Hole).* Dormant application: apply at leaf fall. Dilute: Apply 2 to 4 pounds per 100 gallons (minimum of 8 pounds per acre). Use 16 pounds per acre when rainfall is very heavy and disease pressure is high. Concentrate: Apply at equivalent rates in approximately 100 gallons of water. May be used with agricultural spray oil. **Brown Rot Blossom Blight.** Apply at 2 to 3 pounds per 100 gallons as a full cover spray at pink bud. Application at this time affords some control of Leaf Curl and *Coryneum Blight*. **Bacterial Spot.** Dormant application: apply at 2 pounds per 100 gallons. Post bloom: apply 1.4 pound per 100 gallons at first and second cover sprays. Do not spray later than three weeks prior to harvest. Do not use at rates above those recommended. Caution: Slight defoliation and spotting of leaves may occur from use in cover sprays.

PEARS, Fireblight. Apply at 0.25 pounds per 100 gallons or 1 pound per acre at 5 day intervals throughout bloom period. ***Pseudomonas Blight.*** Apply before fall rains at a rate of 3-4 pounds per 100 gallons (300-400 GPA) and again at dormant before spring growth starts. Excessive dosages may cause fruit russet.

WALNUT, Walnut Blight. Apply first spray at early pre-bloom when catkins are partially expanded. Make three additional applications during bloom and early nut set stage at 7-10 day intervals. Additional applications may be necessary when frequent rainfall occurs. Dilute: Apply 2 pounds per 100 gallons of water (minimum of 8 pounds per acre). Concentrate: Apply at equivalent rates in 50-100 gallons of water per acre. One pint of summer oil emulsion may be added per 100 gallons of spray. Do not apply more than 12.5 pounds per acre per application.

TROPICAL CROPS

BANANAS, (Florida) Anthracnose. Apply at 3 pounds per 100 gallons of water or 12 pounds per acre in 50-100 gallons of water. Apply at 7-10 day intervals starting at first fruit set and continuing until harvest. In areas where the disease is severe, apply at 2 to 3 pounds per 100 gallons or 8-10 pounds per acre at 7-10 day intervals. Apply at 2 to 3 pounds per 100 gallons or 8-10 pounds per acre at 7-10 day intervals. Apply at 2 to 3 pounds per 100 gallons or 8-10 pounds per acre at 7-10 day intervals.

CACAO, Black Pod. Begin applications at the start of season and continue while infection conditions persist. Spray made as often as 14 to 21 days in high rainfall areas. Rates from 2 to 4.5 pounds per acre depending on disease severity. For drier areas where 2 to 4 applications are recommended, critical infection periods and at long intervals, use 6.5 pounds per acre according to disease pressure incident planting density.

COFFEE, Iron Spot (*Cercospora coffeicola*) & Pink Disease (*Colletotrichum salmonicolum*). Apply at 2 pounds per acre as a concentrate dilute spray. Begin treatment at start of wet season and continue at monthly intervals for three applications. **Leaf Rust (Brazil) A.** 3.5 to 5.5 pounds per acre for average density plantation. High density plantations may require 7 to 8 pounds per acre. Make application from September to March depending on altitude and local recommendations. Apply at three to four week intervals depending on disease severity and rainfall conditions.

VEGETABLE CROPS

BEANS, Bacterial Blight (*Halo* & Common). For protective sprays, apply first application when plants are six inches high. Apply on 7-14 day schedule depending on local conditions. Use 1 to 3 pounds per acre depending on disease severity.

BROCCOLI, BRUSSELS SPROUT, CABBAGE & CAULIFLOWER, Downy Mildew. Apply 0.5 to 1 pound in a minimum of 25 GPA at 7 day intervals (CABBAGE ONLY). **Black Rot (*Xanthomonas*) & Black Leaf Spot (*Alternaria*).** Apply at 2 pounds per acre in a minimum of 25 GPA at 7 to 10 day intervals. (Caution: A slight reddening of older leaves may occur on broccoli and a slight flecking of wrapper leaves may occur on cabbage at the 2 pound rate). For control of diseases of these crops, begin applications after transplants are set in the field, or shortly after emergence of field-seeded crops or when conditions favor disease development.

CANTALOUPE, HONEYDEWS & MUSKMELONS, Downy Mildew. Apply weekly at 2 pounds per acre starting before disease appears.

CARROTS, Carrot Blight (*Cercospora*). When disease threatens, apply 2 pounds per acre at 7 to 14 day intervals depending on disease severity.

CELERY, Early Late & Bacterial Blights. Apply as soon as plants are first established in the field at 2 pounds per acre, then every 5-7 days depending on disease severity and weather. One or two quarts of a suitable agricultural spray oil per acre may be used as a sticker-spreader.

CUCUMBERS, Angular Leaf Spot & Downy Mildew. Apply weekly once the plants begin to vine. Use at 1.5 to 2 pounds per acre.

EGGPLANT, *Alternaria Blight*, Anthracnose, Phomopsis. Use 2 pounds per acre before disease appears. Repeat at 7 to 10 day intervals. (EXCEPT CALIFORNIA).

LETTUCE, Downy Mildew. Apply 1 to 2 pounds per 100 gallons of water. Begin treatment when disease first appears and repeat every 7-10 days as needed to suppress disease.

ONION, Purple Blotch & Downy Mildew. Apply at 2 pounds per acre when plants are 4 to 6 inches high and repeat at 7 to 10 day intervals.

PEAS, Powdery Mildew. Begin spray treatment when disease symptoms first appear. Use 1.5 to 3 pounds per acre according to disease severity. Repeat applications at weekly intervals.

PEPPERS, Bacterial Spot. When disease threatens, apply 2 to 3 pounds per acre in sufficient water for adequate coverage at 7 to 14 day intervals depending on disease severity.

POTATOES, Early & Late Blight. Apply at 7 to 10 day intervals starting when plants are 6 inches high until two weeks before harvest. Apply 1 to 1.5 pounds per acre where disease is light. Use 3-4 pounds per acre where disease is more severe.

PUMPKIN & SQUASH, Powdery Mildew. Begin application when plants are three weeks old or when first disease symptoms appear. Use at weekly intervals at 1.5 to 3 pounds per acre depending on disease severity.

TOMATOES, Early Blight. When disease threatens, apply 2 to 3 pounds per acre at 7-10 day intervals. **Bacterial Speck.** Apply at 2 pounds per acre at 10-30 day intervals beginning when the disease threatens. Use more frequent applications when disease pressure is high. **Bacterial Spot.** When disease threatens, apply 2 to 4 pounds per acre at 7-10 day intervals, more frequently when disease is severe. May be combined with mancozeb or mancozeb. Do not apply within five days of harvest when either mancozeb or mancozeb are used.

WATERMELON, Anthracnose & Downy Mildew. Apply as soon as plants are established and at weekly intervals thereafter. Apply at 2 to 3 pounds per acre at 7-10 day intervals. Use at 2 to 3 pounds per acre at 7-10 day intervals.



AGTROL CHEMICAL PRODUCTS

1324 Southwest Freeway, Suite 100
Houston, Texas 77074
713 995 0111

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is a reason to believe that written warnings cannot be understood by workers. Written warnings for this product must include the signal word DANGER-PELIGRO, the name of this product which is CHAMPION WETTABLE POWDER and the date of application. Do not enter treated area without protective clothing until spray has dried. In case of accidental exposure refer to the Statement of Practical Treatment on this label. Because certain states require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

FROST INJURY PROTECTION. Bacterial Ice Nucleation Inhibitor. Application of CHAMPION WETTABLE POWDER made to all crops listed on this label at rates indicated below just prior to anticipated frost conditions will afford control of ice nucleating bacteria (*Pseudomonas Syringae*, *Erwinia Herbicola* and *Pseudomonas Fluorescens*) and may therefore provide protection against light frost. Not recommended in those geographical areas where weather conditions favor severe frost.

HOMEOWNER USE

One level tablespoon of CHAMPION WETTABLE POWDER per gallon of water is equivalent to 1 pound per 100 gallons. Make application according to specific instructions given on this label for those crops where rates are given in pounds per 100 gallons. (For example, 2 pounds of CHAMPION WETTABLE POWDER per 100 gallons are equivalent to 2 level tablespoons of CHAMPION WETTABLE POWDER per gallon.)

BERRIES, VINES AND HOPS

BLACKBERRIES. (*Santiams Logans Boysens Marions Auroras Cascades Chenaieims and Thornless Evergreens*) **Leaf & Cane Spot.** Apply delayed dormant spray after training in spring at 4 pounds plus 1 quart superior-type oil per 100 gallons. Apply again in late spring at 2 pounds plus one quart superior-type oil per 100 gallons. Make fall spray application after harvest using 4 pounds plus 1 quart superior-type oil per 100 gallons.

CRANBERRY. **Fruit Rot.** Apply at 8 pounds per acre beginning in late bloom. One or two additional applications made at 10 to 14 day intervals may be required depending on disease pressure. Follow the advice of the State Agricultural Extension Service.

CURRENTS & GOOSEBERRY. **Leaf Spot.** Make three applications at 10 pounds per acre starting after harvest before bloom and after petal fall.

GRAPES. **Black Rot Powdery Mildew & Downy Mildew.** Apply 2 pounds plus 2-6 pounds hydrated lime per acre as a dilute or concentrate spray. Use for the last one or two late summer applications following early season application of another fungicide. Follow State schedule for exact timing. (Caution: Slight to severe foliage injury may occur on copper-sensitive varieties such as Concord Delaware Niagara and Rosette).

HOPS. **Downy Mildew.** Apply 2 pounds as a fungicide crown treatment after pruning but before training; as needed. After training, additional fungicide treatments are needed at about 10 day intervals. Discontinue use 2 weeks before harvest.

STRAWBERRIES. **Leaf Spot & Leaf Blight.** Apply at 2 to 3 pounds in 100 gallons per acre. Begin application when plants are established and continue on a weekly schedule throughout season. Discontinue applications if signs of phytotoxicity appear.

FIELD CROPS

ALFALFA. **Cercospora & Leptosphaerulina Leaf Spots.** Apply at 2 pounds per acre 10-14 days before each harvest or earlier if disease threatens. Apply with ground or aerial equipment. Spray injury may occur with sensitive varieties such as Labontan.

PEANUTS. **Cercospora Leaf Spot.** Begin spraying 40-45 days after planting or when disease symptoms first appear. Make ground or aerial applications at 1.5 to 3 pounds per acre. For aerial application use 3-10 gallons of water. Continue applications at 10-14 day intervals. Use sufficient water to get adequate coverage. One to two quarts of six pound per gallon flowable sulfur may be added.

SUGAR BEETS. **Cercospora Leaf Spot.** Start spray when disease threatens and continue for 4 to 5 applications. Spray every 10-14 days depending on weather conditions at 2 to 5 pounds per acre depending on disease severity. Addition of suitable agricultural spray oils recommended at 2 quarts per acre.

WHEAT & BARLEY. **Septoria Leaf Blotch & Helminthosporium Spot.** Apply at 1 to 2 pounds per acre. Make first application at early heading and follow with second application 11 days later.

ORNAMENTALS

LIVE OAK. **Bark Moss.** Texas only. Apply at 6 pounds per 100 gallons of water in spring after heavy rain using 10 gallons of spray per foot tree height. Make sure to wet trunks thoroughly. A second application may be required after 10 months.

PHILODENDRON. **Bacterial Leaf Spot.** Apply weekly before disease appears at 1.5 pounds of CHAMPION WETTABLE POWDER plus 1.5 pounds of a coordination product of maneb and zinc (80% active compound) per 100 gallons of water.

SYCAMORE. **Anthraxnose.** Make two applications using 2 to 3 pounds per 100 gallons as a full cover spray. Make first application at bud crack and second application 7-14 days later at 10% leaf expansion.

To the following crops apply CHAMPION WETTABLE POWDER at 1 pound per 100 gallons as a full cover spray beginning at first sign of disease. Repeat at intervals of 7 to 14 days depending on rainfall and disease severity.

ARALIA. **Xanthomonas & Cercospora Leaf Spots, Alternaria**

AZALEA*. **Cercospora Leaf Spot, Botrytis Blight, Phytophthora Dieback & Powdery Mildew**

BEGONIA. **Xanthomonas Leaf Spot, Anthracnose**

BULBS (EASTER LILY, TULIP, GLADIOLUS). **Botrytis Blight**

CARNATION*. **Alternaria Blight, Pseudomonas Leaf Spot & Botrytis Blight**

CHRYSANTHEMUM*. **Septoria Leaf Spot & Botrytis Blight**

COTONEASTER. **Botrytis Blight**

EUONYMUS. **Botrytis Blight, Anthracnose**

IVY*. **Xanthomonas Leaf Spot**

PACHYSANDRA. **Volutella Leaf Blight**

PERIWINKLE. **Phomopsis Stem Blight**

PYRACANTHA. **Fireblight, Scab**

ROSE*. **Powdery Mildew, Black Spot**

YUCCA (ADAMS NEEDLE). **Cercospora & Septoria Leaf Spots**

* On some varieties a discoloration may occur on foliage or blooms. To prevent residues on commercial plants, do not spray just before selling season.

CAUTION: Phytotoxicity may occur on certain varieties. Apply on a few plants at the recommended rate and observe for a few days to see if phytotoxicity will occur.

TREE CROPS

ALMONDS. **Coryneum Blight & Blossom Brown Rot.** Use 2 to 3 pounds CHAMPION WETTABLE POWDER per 100 gallons of water (300-400 GPA) in late dormant before foliage buds swell. If frequent rainfall occurs, a second application should be made during the early bloom stage (popcorn). Apply 5-8 pounds per acre in 7.5-20 GPA by air. To avoid plant injury, do not use above rate after full bloom. **Bacterial Blast (Pseudomonas).** Use 3 to 4 pounds CHAMPION WETTABLE POWDER per 100 gallons and apply at dormant to early pink bud. For blast control in sprinkler irrigated orchards or where disease is severe, apply 2-4 CHAMPION WETTABLE POWDER sprays at 0.25 pound per 10 gallons at 2 weeks post bloom intervals. Just before spring frost, light leaf injury may occur from post bloom spray.

APPLE. **Anthraxnose, European Canker, Pseudomonas.** Apply before fall rains at 3-4 pounds per 100 gallons (300-400 GPA). Use on new varieties may cause discoloration. To avoid pink before spraying. **Fireblight.** Apply at 2 to 4 pounds per 100 gallons of water as a full cover spray. Make application between silver tip and green tip.

(Continued)

ACCEPTED
with COMMENTS
in EPA Letter Dated:

NOV 13 1986

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.**CHAMPION® WETTABLE POWDER AGRICULTURAL FUNGICIDE**
PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****DANGER — PELIGRO**

Corrosive — Causes irreversible eye damage. May be fatal if inhaled. Harmful if swallowed or absorbed through skin. Do not get in eyes or inhale dust. Wear goggles or face shield and dust mask while handling, mixing or loading. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

(PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que le etiqueta haya sido explicado ampliamente.)

ACCEPTED

NOV 13 1986

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under
EPA Reg. No. 55146-1**STATEMENT OF PRACTICAL TREATMENT**

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF IN EYES: Flush with copious volumes of water. Get medical attention immediately.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

GENERAL INSTRUCTIONS

CHAMPION WETTABLE POWDER can be used with all types of spraying equipment. The volume per acre will differ depending on the specific crop and the equipment used. Use CHAMPION WETTABLE POWDER as per instructions on this label.

APPLYING SPRAY MIXTURE The directions given under each crop are for applying dilute spray mixture unless otherwise shown. The amount of CHAMPION WETTABLE POWDER applied per acre in concentrate and aerial sprays should be the same as the amount applied per acre in dilute sprays. The required amount should be mixed with enough water to thoroughly cover the crop with spray mixture. The volume of water needed will depend upon the spray equipment used and the size of the crop being sprayed. Read the information below about applying dilute, concentrate, and aerial sprays.

APPLYING DILUTE HIGH-VOLUME SPRAYS On vegetable crops use 25 to 100 gallons of spray mixture per acre, on fruit and nut trees use 250 to 800 gallons per acre. As much as 1500 gallons per acre may be needed for large trees.

APPLYING CONCENTRATE GROUND SPRAYS On vegetable crops use 5 to 20 gallons of spray mixture per acre, on fruit and nut trees use 25 to 250 gallons per acre.

APPLYING AERIAL SPRAYS Use 3 to 30 gallons per acre.

No additional surfactants are needed. Add CHAMPION WETTABLE POWDER slowly to the spray tank followed by any sticker-spreaders, insecticides, nutrients etc. Observe all cautions and limitations on the label of all products used in mixtures. The specific instructions given on this label are based on general applications and circumstances. The recommendations of the State Agricultural Extension Service should be closely followed as to timing, frequency and number of sprays per season.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste dump facility.

CONTAINER DISPOSAL Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY STATEMENT

AGTROL CHEMICAL PRODUCTS warrants that the product conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions. THERE ARE NO OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING A WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE. This warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller and buyer assumes all risk of any such use.

BEST AVAILABLE COPY