EPA Form 8570-6 (Rev. 5-76) PARVIOUS EDITION MAN

ATTACHMENT IS APPLICABLE
SIGNATURE OF APPROVING OFFICIAL

d. Under Statement of Practical Treatment change "If in Eyes. . . "
to read "If in Eyes. . . Get medical attention immediately."

 $\mathcal{J}_{M_{i}}$

- e. Add paragraph II A.7 from PR Notice 83-2 Farmworker Safety.
- f. Delete the claim to prevent frost injury. Haneb would not be expected to be beneficial for this use so the use of this product would result in an unnecessary additional per licide burden on the environment.
- g. For beans, add "Do not apply to succulent beans within 4 days of harvest."
- h. For celery, the sentence relating to removal of residues should read "Remove residues by stripping, trimming, and washing."
- i. Indicate when to use the high and low rates for the crops other than beans.
- 3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

The additional brand name listed below is also being accepted under this registration. The labels must be identical with the basic label except for brand name.

Additional Brand Name

Manpower

Henry H. Jacoby Product Hanager (21) Fungicide-Herbicide Branch Registration Division (TE-767C)

Enclosures

FRONT LAUEL



AGTROL CHEMICAL PRODUCTS HOUSTON Texas 77074

AGRICULTURAL

FUNGICIDE

भागका प्रा

STATEMENTS, AND STATEMENT OF PRACTICAL TREATMENT

The area being treated must be vacated by unprotected persons. Do not enter area without protective clothing until spray has dried.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce womiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF IN EYES: Flush with copious volumes of water. Get medical attention if irritation persists.

...

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste dump facility.


CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incinerate, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY STATEMENT

AGTROL CHEMICAL PRODUCTS warrants that the product conforms to the chemical description on the label and is reasonably it for the purposes set forth on the label when used according to directions under normal use conditions. **THERE ARE NO OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING A WARRANTY OF FITNESS FOR A PAPTICULAR PURPOSE. This warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably forseeable to seller and buyer assumes all risk of any such use.



DIRECTIONS FOR USE



RE-ENTRY STATEMENT

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Do not enter treated area without protective clothing until spray has dried. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

FROST INJURY PROTECTION: Bacterial Ice Nucleation Inhibitor: Application of MANPOWER made to all crops listed on this label at rates indicated below just prior to anticipated frost conditions, will afford control of ice nucleating bacteria (Pseudomonas Syringae, Erwina Herbicola and Pseudomonas Fluorescens) and may therefore provide protection against light frost. Not recommended in those geographical areas where weather conditions favor severe frost.

GENERAL INSTRUCTIONS

MANPOWER can be used with all types of spraying equipment. The volume per acre will depend on the equipment used. Use MANPOWER according to instructions on this label. For high volume sprays (dilute) use 25-100 gallons per acre (GPA). For concentrate ground sprays use 5-20 GPA and for aerial spraying use 3-15 GPA. No additional surfactants are needed. Add MANPOWER slowly to the spray tank followed by any other sticker-spreaders, insecticides, nutrients, etc. Observe all cautions and limitations on the label of all products used in mixtures. The specific instructions given on this label are based on general applications and circumstances. The recommendations of the State Agricultural Extension Service should be closely followed as to timing, frequency and number of sprays per season.

VEGETABLE CROPS

BEANS, Bacterial Blight (Halo and Common), Rust, Downy Mildew. Apply first when plants are small on 7 to 14 day schedule depending on local conditions, use 4-6 pints per acre depending on disease condition.

CELERY, Early, Late and Bacterial Blight. Apply as soon as plants are established in the field. Repeat as often as necessary or every 7-10 days in the field depending on disease severity. Use 4-6 pints per acre. Remove residues by washing. Do not apply within 14 days of harvest.

PEPPERS, Bacterial Spot, Anthracnose, Cercospora Leaf Spot. Apply 4-6 pints in sufficient water to obtain good coverage. Begin when disease threatens and apply every 7-10 days depending on disease severity.

TOMATO, Early Blight, Late Blight, Anthracnose. Apply at 4-6 pints when disease threatens at 7-10 day intervals. Bacterial Speck. Apply at 6 pints at 10-30 day intervals when disease threatens. Apply more frequently when disease pressure is high. Bacterial spot. Begin when disease threatens. Apply 6-8 pints per acre at 7-10 day intervals or more frequently when disease is severe. Do not apply within five days of harvest.



