

FRONT LABEL

<b>ACCEPTED</b>
<b>MAY 13 1987</b>
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No.



**AGTROL CHEMICAL PRODUCTS**

Houston, Texas 77074

This product must not be used on golf courses and sod farms

# DIAZINON AG 500 INSECTICIDE

## For Agricultural/Commercial Use Only

**ACTIVE INGREDIENTS:**

DIAZINON [O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate]	47.5%
XYLENE	39.4%
INERT INGREDIENTS:	13.1%
Total	100.0%

**DIAZINON AG 500 CONTAINS 4 LBS. DIAZINON PER GALLON**

### KEEP OUT OF REACH OF CHILDREN

### WARNING

See side panel for additional precautionary statements.

#### PRACTICAL TREATMENT: (FIRST AID)

**IF SWALLOWED:** Call a physician or Poison Control Center immediately. Gastric lavage is indicated if material was taken internally.

**DO NOT INDUCE VOMITING** unless other treatment is not available. Vomiting may cause aspiration pneumonia.

If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

**IF INHALED:** Remove victim to fresh air. Give artificial respiration if victim stops breathing. Get medical attention immediately.

**IF ON SKIN:** Remove contaminated clothing and wash contaminated skin with soap and water.

**IF IN EYES:** Flush eyes with plenty of water for at least 15 minutes. Get medical attention immediately.

EPA REG. NO. 55146-37  
EPA EST. NO.

**NET CONTENTS:  
1 GALLON**

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**WARNING**

Keep out of reach of children. May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly before smoking or eating and after handling. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not contaminate ornamental fishponds. Do not use on humans, household pets or livestock. Keep children and pets out of treated areas until spray has dried.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use, pour, spill or store near heat or open flame.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds, and other wildlife feeding in treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes, and estuaries (except for effluence from treated watercourses). Do not apply to water that will be used for recreational purposes and human and livestock consumption. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, crab and other aquatic life are important resources. This product must not be used in areas where impact on threatened endangered species is likely. Contact your State Fish and Game Agency before applying this product. Apply this product only as specified on this label.

**GENERAL INFORMATION**

Diazinon AG 500 is an emulsifiable solution which when diluted with water according to the directions is intended for the control of insect pests of agricultural crops. Do not use in barns or dwellings. Do not use on livestock.

In the directions for use which follow, best control is obtained when application is made at first sign of insects. Application should be repeated as necessary to maintain control.

**NOTE:** The California Department of Agriculture has set a 5-day reentry for Diazinon on citrus groves, orchards and nutcrimes. Five days should elapse between the time of application and the time a worker may enter the field to engage in any activity requiring substantial contact with treated foliage. When a mixture of two or more organic phosphorus pesticides is applied, the interval may be prolonged by adding to the longest applicable interval an additional 50% of that interval.

**DIRECTIONS FOR USE -**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE DISPOSAL:** Pesticide containers or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved State procedures under Subtitle C of the Resource Conservation and Recovery Act.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures.

**VEGETABLE POLAR INSECTS:**

Unless otherwise stated, foliar applications of Diazinon AG 500 should always be applied to vegetable crops as a spray in a minimum of 2 gal. of water per acre for aerial applications and a minimum of 10 gals. of water per acre for ground applications.

**BEANS:** Pole Beans, Snap Beans, Lima Beans: Black Bean Aphids, Mexican Bean Beetles, Leafminers. Apply 1/4 pts. per acre. Cucumber Beetles. Apply 1/2 pt. in 100 gals. of water per acre. Do not apply within 7 days of bean harvest. Bean forage or bean hay may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay.

**LIMA BEANS:** Diterous Leafminers, Leafhoppers, Mites. Apply 1/2 pt. per acre. Do not apply within 7 days of harvest. Bean forage or bean hay may be fed to dairy cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay.

**BROCCOLI, CABBAGE, CAULIFLOWER:** Aphids, Diamondback Moths, Imported Cabbageworms. Apply 1/2 to 1 pt. per acre. Applications at 7-10 day intervals will usually be required. Do not apply to broccoli or cauliflower within 5 days, or to cabbage within 7 days of harvest.

**BRUSSELS SPROUTS:** Aphids, Diamond Moths, Imported Cabbageworms. Apply 1/2 pt. per acre. Generally applications at 7-10 day intervals will be required. Do not apply to brussels sprouts within 7 days of harvest.

**CARROTS:** Aphids. Apply 1/2 pt. per acre. Do not apply within 10 days of harvest.

**CELERY:** Aphids, Flea Beetles. Apply 1/2 pt. in 100 gals. of water per acre. Do not apply within 10 days of harvest. Plants should be market trimmed (tops removed) before being shipped or used. Tops should not be used for feed or food.

**COLLARDS, KALE, SWISS CHARD, TURNIP TOPS:** Aphids, Imported Cabbageworms, Diamondback Moths, Harlequin Cabbage Bugs. Apply 1/2 pt. per acre. Do not apply to collards, kale or turnip tops within 10 days, or to Swiss Chard within 12 days of harvest.

**CUCUMBERS:** Aphids, Serpentine Leafminers, Thrips. Apply 1/2 pt. per acre. Do not apply within 7 days of harvest.

**LETTUCE, ENDIVE:** Aphids, Diterous Leafminers. Apply 1/2 pt. per acre. Do not apply within 10 days of harvest.

**MELONS, SQUASH, CANTALOUPE, CASABAS, CHERSHAWNS, HONEYDEW MELONS, MUSKMELONS, PERSIAN MELONS, AND HYBRIDS OF THESE; WATERMELONS AND THEIR HYBRIDS, SUMMER AND WINTER SQUASH:** Aphids, Leafhoppers, Mites. Apply 1-1 1/2 pts. per acre. Serpentine Leafminers, Thrips. Apply 1/2 pt. per acre. Do not apply to melons and winter squash within 3 days, or to summer squash within 7 days of harvest.

**MUSHROOM HOUSES:** Mushroom Flies, Aphids and Sciarid. Mix 1/2 Gal. in 50 Gals. of water, 0.5% to 1.0% actual Diazinon and spray on outside and inside walls and ventilators of mushroom houses after compost in beds has been pasteurized by heating. Do not spray compost in beds or after spawning. To treat wooden sides of beds, pests, doors and door frames after pasteurizing mix 1/2 pt. in 6 gals. of water and apply with a brush. Do not treat compost.

**CARROTS:** Onion Thrips. Apply 1/2 pt. per acre. Do not apply within 10 days of harvest.

**PEAS:** Aphids, Leafminers. Apply 1/2 pt. per acre. Peas may be picked immediately. Pea vines and pea hay may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay.

**PEPPERS:** Aphids, Serpentine Leafminers. Apply 1/2 pt. per acre. Do not apply within 5 days of harvest.

**POTATOES:** Aphids, Flea Beetles, Colorado Potato Beetles, Diterous Leafminers. Apply at the rate of 1/4 gals. per acre. Banded Cucumber Beetles, Leafhoppers, Southern Armyworms. Apply 1/2 pt. per acre. Do not apply within 35 days of harvest.

**RADISHES, PARSNIPS, TURNIPS:** Aphids, Flea Beetles, Diterous Leafminers. Apply 1/2 pt. per acre. Do not apply within 10 days of harvest.

**SPINACH, BEETS:** Aphids, Diterous Leafminers. Apply 1/2 pt. per acre. Do not apply to spinach within 10 days, or to beets within 14 days of harvest.

**SQUASH (Summer and winter):** See Melons and Squash section.

**BEST AVAILABLE COPY**

**SWEET CORN:** Corn Earworms: Apply 1-1 1/2 pts. per acre when silks first appear. Two or three repeat applications at 7-3 day intervals will usually be necessary. Corn may be picked immediately following last application. Corn borers may be fed to dairy and beef cattle and sheep. Refer also to Corn under Field and Forage For Insects and Field and Forage Soil Insects.

**TYRANTFLIES:** Aphids, Destructive Leafhoppers: Apply 1/2 pt. per acre To control Leafhoppers, weekly applications are usually required. Banded Cucumber Beetles, Fall Armyworms, Southern Armyworms: Apply 1/2-1 pt. per acre. Vinegar Flies (Drosophila spp.): Apply 1/2-1 pt. per acre. Weekly applications are usually required. Do not apply within 1 day of harvest.

**WATERBESS:** (Narrow only) Cyclamen Mites: Mix 1 qt. per 100 gals. of water and apply to one acre as a thorough cover spray. Spray should be directed into the crown and growing base of plants. If infestation is heavy, make a second application no later than 10 days after the first application. Repeat applications as necessary. Do not apply within 5 days of harvest.

**VEGETABLE SOIL INSECTS:** Unless otherwise stated soil applications of Dazelon AG 500 should always be applied to vegetable crops as a spray in a minimum of 5 gals. of water per acre for ground applications.

**CUTWORMS:** (Surface and Subterranean)

Broccoli	Endive (Escarole)	Potatoes
Brussels Sprouts	Kale	Snap Beans
Cabbage	Lettuce	Sprach
Cantaloupes	Lima Beans	Summer Squash
Cauliflower	Mushrooms	Sweet Corn
Celery	Parley	Swiss Chard
Collards	Peas	Tomatoes
Cucumbers	Peppers	Watermelons
		Winter Squash

Broadcast 2-4 qts. per acre prior to planting, and immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.

**GARDEN STYPTERYLARS:** (Caterpillars)

Cabbage	Peas	Snap Beans
Carrots	Pole Beans	Tomatoes
Corn	Radishes	Turnips
Lettuce	Red Beets	

Broadcast 10 qts. per acre just prior to planting immediately work into soil 4-8 inches. This preplant treatment will also control other soil insects, such as cutworms (surface and subterranean), mole crickets, root maggots, and wireworms.

**MOLE CRICKETS:**

Broccoli	Endive (Escarole)	Radishes
Brussels Sprouts	Kale	Tomatoes
Cabbage	Lettuce	Turnips
Cauliflower	Mustard	
Carrots	Peppers	

Broadcast 1 qt. per acre when insects are present. Make application to plant beds of field 1-2 days prior to sowing of seed or transplanting and immediately work into soil 1-2 inches.

**ONION MAGGOTS:** Onions  
Furrow Application: Apply 1 qt. per acre in sufficient water to drench the seed furrow at planting time. Broadcast Application: Broadcast 2-4 qts. per acre just prior to planting, immediately work into soil 3-4 inches.

**ROOT MAGGOTS:**  
Broccoli, Brussels Sprouts, Cabbage, Cauliflower: Broadcast Application: Broadcast 2-3 qts. per acre and immediately work into the soil 3-4 inches, before seeding, either in the seedbed or in the field or before transplanting. Transplant water (drench) application: Mix 1/2-1 pt. in 50 gals. of transplant water when 200-300 gals. of water are used per acre. May be applied at the rate of 1-1 1/2 cups per plant, either by hand or by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Transplant water treatment may result in soil-oil reduction due to plant stress at time of transplanting.  
Radishes: Mix 1 qt. in 50 gals. of water. Apply 80-100 gals. per acre (1-1 1/4 gals. per 1,000 linear feet of row) in the seed furrow at planting time.

**WIREWORMS:**

Broccoli	Endive (Escarole)	Potatoes
Brussels Sprouts	Kale	Snap Beans
Cabbage	Lettuce	Sprach
Cantaloupes	Lima Beans	Summer Squash
Carrots	Mushrooms	Sweet Corn
Cauliflower	Onions	Sweet Potatoes
Celery	Parley	Swiss Chard
Collards	Peas	Tomatoes
Cucumbers	Peppers	Watermelons
		Winter Squash

Broadcast 3-4 qts. per acre prior to planting, and immediately work into soil 4-8 inches. Potatoes: Broadcast 1-4 qts. per acre prior to planting immediately disk or harrow into soil 4-8 inches.

**SOUTHERN POTATO WIREWORMS:** (Southeastern U.S.) Potatoes for use to control southern potato wireworms in potatoes grown in mineral soils in the Southeast, apply 2-4 qts. per acre as a soil treatment prior to planting and immediately mix thoroughly by disk or other suitable means into the top 3-6 inches of soil depending on location of wireworms.

**FRUIT AND NUT CROPS:**  
NOTE: UNLESS OTHERWISE STATED THE FOLLOWING RATES SHOULD BE APPLIED AS A DILUTE FOLIAR COVER SPRAY.

**ALMONDS:** Mites: Twig Borers: Apply 1/4 qts. per 100 gal. of water. Perilaria Scales: San Jose Scales: Mix 1/4 qts. Dazelon AG 500 plus 2-3 qts. dormant oil per 100 gals. of water and apply as a dormant spray. Almond hulls may be used for food for insects.

**APRICOTS:** Aphids: Brown Mites (Clover Mites): Two-spotted Spider Mites: Olive Scale Crawlers: San Jose Scale Crawlers: Mix 1 qt. per 100 gals. of water. Apply when aphids, mites, and scale crawlers first appear. Olive Scale Crawlers: Mix 1/2 qt. of Dazelon AG 500 plus 1/4 gal. light medium horticultural oil per 100 gals. of water. Apply when scale crawlers are present. Apricot Mite: Mix 1 qt. per 100 gals. of water. Apply as a cover. Do not apply within 10 days of harvest.

**BLUEBERRIES:** Cranberry Flea Beetles: Cherry Fruitworm: Blueberry Maggot: Aphid: Thrips: Two-spotted Mites: Mix 1 qt. per 100 gals. water. Apply up to 200 gals. of spray per acre. Do not apply within 7 days of harvest.

**CANEBERRIES:** Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries: Leafhoppers: Thrips: Aphids: Two-spotted Mites: Raspberry Sawflies: Apply 2 qts. per acre in a minimum of 100 gals. of water. Raspberry Fruitworms: Apply at the above rates when buds are in buds separate and repeat application immediately before blossoms open. Dewberry Mites: Apply 1-2 qts. per acre in a minimum of 100 gals. of water. Apply as a dormant spray in early spring when canes are still on the ground and buds are closed. Make a second application when the buds are well developed but still closed. Do not apply within 7 days of harvest.  
Raspberries Fruitworm Crown Borers (Raspberry Root Borers): Apply 2 qts. per acre in a minimum of 100 gals. of water as a drench to the crown area and to the lower canes. Apply either in the spring before berries begin to form or in the fall after harvest.

**CHERRIES:** (Two-spotted Bud Moths: Fruit Tree Leafrollers: Mix 1 qt. per 100 gals. of water. Apply in cover sprays as necessary beginning with petal fall.  
Cherry Fruit Flies: Mix 1/2 qt. per 100 gals. of water. Makes 3 or 4 applications beginning when adult cherry fruit flies begin to emerge and repeat at 10-day intervals.  
Leafhoppers: Apply 1 qt. per 100 gals. of water in 3-5 applications beginning at petal fall and continuing through the growing season or apply as infestations develop.  
Black Cherry Aphids: Mix 1 qt. per 100 gals. of water. Apply as a cover spray when aphids appear and repeat after 10 days.  
San Jose Crawlers: Mix 1 qt. per 100 gals. of water. Apply during the pre-blossom stage or when crawlers first appear. Do not apply within 10 days of harvest.  
Cherry Rust Mites: Mix 1/2 qt. per 100 gals. of water. Apply as a supplemental spray after harvest.

**CITRUS FRUITS: GRAPEFRUIT, LEBBORS, LIMES, ORANGES, TANGELOS, TANGERSINES, CITRUS CITRUS, KUMQUATS, AND HYBRIDS OF THESE:**  
Citrus Aphids: Fruit Tree Leafrollers: Cottoncushion Scale Crawlers: Soft Scale Crawlers: Apply 1-1 qt. per 100 gals. of water (maximum 1,000 gals. per acre) or a thorough cover spray when aphids and crawler stages of scale insects first appear. Fruit Tree Leafrollers use 1/2 qt. per 100 gals. of spray (maximum 1,000 gals. per acre) when an average of 50 percent of eggs in egg masses have hatched in the spring. Citrus Thrips: Apply 1 qt. in 100 gals. of water (maximum 1,000 gals. per acre) when thrips first appear.

**Limes: Lemons: Scales:** Apply 1-2 qts. per 100 gals. of water (maximum 1,000 gals. per acre) as a pre-blossom spray and again as a summer spray. In heavy green scale infestation, use the higher dosage rate. Thorough coverage of tree trunks, limbs and buds is essential. Do not apply during bloom. Do not spray in citrus within 21 days of harvest. Sprays and craw may be mixed at application rates recommended on this label. Do not apply where these are important resources.



**CRANBERRIES:** Black-headed Fireworms (Rhagoletis cerasana) Apply 2 etc. in 400 gals. of water per acre.  
 Cranberry Fruitworms: Apply 3 etc. in 400 gals. of water per acre. Do not apply within 7 days of harvest.

**FIGS:** Vinegar Flies (Drosophila spp.) Orinid Fruit Borer: Apply 1 pt. per 100 gals. of water. Do not apply within 5 days of harvest.

**FRUITS:** Filbert Leafrollers: Aphids: Apply 1-2 etc. in 250-400 gals. of water per acre as a thorough cover spray.

**GRAPES:** Pacific Solder Mites, Leafhoppers, Grape Berry Moths, Grape Leafhoppers: Apply 1-2 etc. per acre in a minimum of 100 gals. of water as a thorough cover spray. Do not apply within 6 days of harvest.

**Drosophila spp.** Apply 12-18 fl. oz. per acre in a minimum of 100 gals. of water as a thorough cover spray. Do not apply when insects first appear. Make additional applications of weekly intervals as necessary. Do not apply within 7 days of harvest.

**OLIVES:** Olive Scars: Mix 1/2 pt. of Diazinon AG 500 plus 1/4 gal. of medium summer horticultural oil per 100 gals. of water. Apply as a cover spray in late winter. Do not apply during bloom to avoid injury to pollinating insects or within 75 days of harvest.

**PEACHES, NECTARINES:** Aphids, Brown Mites (Clover Mites), Two-spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers: Mix 1 pt. per 100 gals. of water. Apply when aphids, mites, and scale crawlers first appear.

**Olive Scale Crawlers:** Mix 1/2 pt. of Diazinon AG 500 plus 1/4 gal. light medium horticultural oil per 100 gals. of water. Apply when scale crawlers are present.

**Oriental Fruit Moths:** Mix 1 pt. per 100 gals. of water and apply at 100% petal fall. Make 2 additional applications at 10-15 day intervals. Time the first application to peak adult emergence. If possible, time to coincide with both crawler and pupal activity. Under conditions of heavy infestations, two applications may be necessary.

**Do not apply to peaches within 20 days or to nectarines within 10 days of harvest.**

**PEACHES:** Peach Tree Weevils, Pecan Nut Casebearer, Walnut Caterpillars: Apply 1-3 etc. per acre as a "shock spray" in a minimum of 100 gals. of water. Do not apply after nutset crack.

**White Peach Scale (Peaches Only):** As a post-harvest spray apply 1 1/2-2 etc. per 100 gals. of water. Time to coincide with both crawler and pupal activity. Under conditions of heavy infestations, two applications may be necessary.

**Do not apply to peaches within 20 days or to nectarines within 10 days of harvest.**

**PECAN NUT CASEBEARER:** Mix 1 pt. per 100 gals. of water. Do not apply after nutset crack.

**PINEAPPLE:** Scars: Apply 5 etc. of Diazinon AG 500 plus 2 gals. of light or medium horticultural oil in 500 gals. of water per acre to the vegetative stage on the plants following harvest of pineapple.

**Walnuts:** Apply 2 etc. per acre in a minimum of 100 gals. of water. May be applied on day of harvest.

**Prunus foliage and twigs may be fed to livestock 7 days following last application.**

**PLUMS, PRUNES:** Brown Mites (Clover Mites), European Red Mites: Apply 1 pt. per 100 gals. of water.

**Self-cult Plum Aphids, Mealy Plum Aphids, Thistle Aphids:** Apply 1/2 pt. per 100 gals. of water. Do not apply within 10 days of harvest.

**STRAWBERRIES:** Aphids, Two-spotted Solder Mites: Apply 1 pt. per 100 gals. of water per acre. Cyclamen Mites: Apply 1 pt. per 100 gals. of water per acre directing spray to plant crowns. Plants should be sufficiently irrigated to assure thorough coverage of the foliage. Make additional applications if necessary after harvest.

**Worm Crickets:** Broadcast 1 pt. per acre when insects are present. Apply 1-2 days before transplanting and immediately mix into the top 1-2 inches of soil.

**Strawberry Leafrollers:** Mix 1/2 pt. per 100 gals. of water. Apply 200 gals. of spray per acre when blossoms show color. Do not apply within 5 days of harvest.

**NALBUTS:** Aphids, Mites: Apply 1 pt. per acre as a foliage spray in a minimum of 100 gals. of water.

**Lugging Moths, Scale Crawlers, Walnut Caterpillars:** Apply 3 etc. per acre as a foliage spray in a minimum of 100 gals. of water.

**Do not apply after the nuts open.**

**FIELD AND FORAGE FOLIAR INSECTS**

Unless otherwise stated, foliar applications of Diazinon AG 500 should always be applied to field and forage crops as a spray in a minimum of 2 gals. of water per acre for aerial applications and a minimum of 10 gals. of water per acre for ground applications.

**ALFALFA, CLOVER:** Spotted Alfalfa Aphids, Yellow Clover Aphids, Pea Aphids, Leafhoppers, Plant Bug, Hemlock Mites: Apply 1/2 pt. per acre.

**Grasshoppers:** Apply 1 pt. per acre.

Following the above applications: dairy and beef cattle and sheep (1) may be grazed immediately following application; (2) may be fed green alfalfa or clover immediately following cutting; and (3) may be fed hay provided alfalfa or clover is not cut for hay for 7 days following treatment.

Alfalfa meadow larvae for control in the spring, apply 2-3 etc. per acre when feeding damage to growing tips becomes evident. Repeat application if necessary if damage is already extensive. Cut the hay and make application to the stubble. Do not graze dairy and beef animals or sheep for 2 days. Do not cut for hay for 10 days following treatment.

Do not spray livestock directly with either air or ground equipment. Do not treat during bloom to avoid injury to pollinating insects.

**CLOVER:** Striped Ground Crickets: Broadcast 2 etc. per acre following seeding in the fall but prior to clover emergence.

**CORN:** Use a minimum of 2 gal. of water for aerial applications and 5 gals. of water for ground applications.

**Corn Rootworm Adults:** Apply 1/2 pt. per acre.

**Corn Leaf Aphids:** Apply 2 etc. per acre. Complete coverage of the upper portions of the plants is essential. Repeat applications as necessary.

**Mites:** Apply 1 pt. per acre when infestations first appear. Repeat applications as necessary.

**Leaf Beetles:** Apply 1 pt. per acre when insects first appear. Repeat applications as necessary.

**Grasshoppers:** Apply 1 pt. per acre.

**Leaf Beetles:** Apply 2-3 etc. per acre. Add 1/2 gal. of soluble mineral oil per acre to the spray mixture. Consult State Agricultural Experiment Stations or State Extension Service Specialists for proper timing of sprays. Corn may be grazed immediately following last application. Corn forage may be fed to dairy and beef cattle and sheep.

BEST AVAILABLE COPY

