

55146-40


PM 23

1066

FRONT LABEL

AGTROL CHEMICAL PRODUCTS
Houston, Texas 77074

COC 50 WP
WETTABLE POWDER
AGRICULTURAL FUNGICIDE



ACTIVE INGREDIENT		
Copper * Expressed as Metallic.....	50.0%	
INERT INGREDIENTS.....	50.0%	
	TOTAL	<u>100.0%</u>

* Copper as metallic derived from Copperoxychloride


KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

(PRECAUCION AL USURIO: Si usted no lee ingles, no use este producto hasta que le stiqueta sido explicodo ampliamente.)

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
AND STATEMENT OF PRACTICAL TREATMENT

Net Weight 20 lbs.



EPA REG. NO. 55146-

EPA EST. NO.

BEST AVAILABLE COPY

BACK LABEL - 1

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. (Indicate specific oral warnings which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure.) When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: DANGER - PELIGRO. Area treated with ~~COC 50 WP~~ on (date of application). ~~Do not enter without~~ appropriate protective clothing until spray has dried. In case of accidental exposure see Statement of Practical Treatment.


Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

GENERAL INSTRUCTIONS

COC 50 WP contains copperoxychloride together with specially selected and stabilized mineral fillers and appropriate surface active agents. COC 50 WP is spray dried to produce a fine uniform sprayable powder that wets out and goes into suspension readily in water.

DO NOT apply as a dry powder. Fill the spray tank half full of water, begin agitation and add spray adjuvants if needed. As tank continues to fill, add required amount of this product. Constant agitation during spraying operation is necessary to maintain adequate suspension.

This product is compatible with most other insecticides and/or fungicides EXCEPT lime sulfur and dithiocarbamates. When used in combination with oil, OBSERVE THE PRECAUTIONS ON THE OIL LABEL.



BEST AVAILABLE COPY

COC 50 WP is prepared for use against certain diseases of crops as listed below. Unless otherwise noted, dosages are given in pounds of this product per 100 gallons of water to be applied as a full coverage spray with conventional spray equipment. Maximum dosages refer to pounds of this product per acre per application. When rates are given as ranges, use lower rates when disease conditions are light to moderate and higher rates when conditions are severe.

The following directions for use are based on general applications. The recommendations of State Agricultural Extension Services should be closely followed as to timing, frequency and number of sprays per season.

FRUITS

CITRUS: *Greasy Spot, Melanose, Scab* - Apply 1.25 - 2.0 lbs./100 gallons in pre-bloom and post-bloom spray. Higher rate used under conditions of greater disease severity. May be used in concentrate sprays at equivalent rates. For aerial sprays use a minimum of 9 lbs. in 10 - 15 gallons per acre. Do not apply in areas where copper injury is known to occur or where fumigation with hydrogen cyanide gas is practiced.

PEACHES: *Leaf Curl*. As a fall spray, apply 4 to 4 1/2 lbs. from November 14 to December 15. If this application cannot be made, make applications just before bud swell. California Only: To control Leaf Curl, use 12 to 16 lbs. in a minimum of 40 gallons of water per acre by ground or a minimum of 20 gallons of water by air applications. Apply as a dormant spray up to bud swell.

FIELD AND ROW CROPS

For disease control on the following crops, use 3 to 4 lbs. COC 50 WP per acre (unless otherwise indicated) in 25 to 100 gallons of water per acre at 7 to 10 day intervals to protect new growth throughout the growing season.

CELERY: *Bacterial Blight, Early Blight (Cercospora), Late Blight (Septoria)*.



CARROTS: *Leaf Blight, Leaf Spot.*

MELONS, CUCUMBERS: *Angular Leaf Spot, Anthracnose, Downy Mildew, Leaf Blight, Scab and aids in control of Bacteria Wilt.*

PEPPERS, TOMATOES (In cold frames, greenhouses, etc.): Damping-Off - Use 4 lbs. per 100 gallons of water or 2 cunces in 3 gallons. Keep suspension well agitated. Apply at 4 to 7 day intervals, starting when plants break ground. Spray plants lightly but thoroughly. Also apply to soil surrounding the plants.

POTATOES: *Early and Late Blight.*

SUGAR BEETS: *Cercospora Leaf Spot*- Apply 6 lbs. per acre beginning when disease first appears and repeat at 7 to 10 day intervals.


TOMATOES: *Early and Late Blight, Septoria Leaf Spot* - Use 4 lbs. if Late Blight infection is severe, increase dosage up to 7 lbs. on large plants. For Bacterial Spot, (seed bed treatments only) use 5 to 6 lbs. per acre every 5 to 7 days after first leaves appear.

ORNAMENTALS

AZALEAS: *Bud Blight, Twig Blight* - Apply 4 lbs. per 100 gallons of water. Begin when disease is first reported in area. Repeat at 7 to 10 day intervals or as required for control.

VINCA: *Canker, Dieback* - Apply 4 lbs. per 100 gallons of water. Begin when disease is first reported in area. Repeat at 7 to 10 day intervals or as required for control.

BEST AVAILABLE COPY



BACK LABEL - 2

COC 50 WP WETTABLE POWDER

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER - PELIGRO

Corrosive. Causes irreversible eye damage. Harmful if swallowed or absorbed through skin. Harmful if inhaled. Do not get in eyes or inhale dust. Wear goggles or face shield and dust mask while handling, mixing, or loading. Avoid contact with skin, eyes, or clothing. May cause skin sensitization reactions in certain individuals. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink promptly a large quantity of milk, egg white, gelatin solution. If these are not available, drink large quantities of water. Avoid alcohol. NOTE TO PHYSICIAN: Probable mucousal damage may contraindicate the use of gastric lavage.


IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF IN EYES: Flush with copious volumes of water. Get medical attention immediately.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent sites. Do not allow rinsate from cleaning of equipment or disposed material to enter surface or ground water. Apply this product only as specified on this label.



BEST AVAILABLE COPY

STORAGE AND DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, Stay out of smoke.

WARRANTY STATEMENT

AGTROL CHEMICAL PRODUCTS warrants that this product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions. THERE ARE NO OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING A WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE. This warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller and buyer assumes all risk of any such use.

BEST AVAILABLE COPY

