

AGTROL CHEMICAL PRODUCTS
Houston, Texas 77074

FRONT LABEL

GIBGRO 5% Powder

(GIBBERELIC ACID)

KEEP OUT OF THE REACH OF CHILDREN

CAUTION


SEE BACK PANEL AND ACCOMPANYING INFORMATION FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

ACTIVE INGREDIENT		W/W
Gibberellin A.....		5.0%
INERT INGREDIENTS.....		95.0%
WATER.....		0.0%

Contains a total of 16 g. Gibberellin Acid

Net Contents - 16 g. (0.56 oz.) (220 g.)

EPA Reg. No. 55146 - 5



EPA Est. No.

BACK LABEL

GIBGRO 5% Powder

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Powder causes eye irritation. In case of contact with eyes flush thoroughly with water.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

ENVIRONMENTAL HAZARDS

Keep out of lakes, ponds or streams. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL AND CHEMICAL HAZARDS

Powder causes eye irritation. If powder gets in eyes, flush thoroughly with water.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: In case of contact with eyes, flush thoroughly with water. Get medical attention if irritation persists.

IF SWALLOWED: Seek medical aid.

IF ON SKIN: Immediately flush skin with plenty of water. Get medical attention if irritation persists.

IF INHALED: Move to fresh air.

DIRECTIONS FOR USE

General Classification.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

IMPORTANT

Before application, read accompanying GIBGRO 5% Powder Spray Guide carefully and use only as directed. DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

2

RE-ENTRY STATEMENT

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep containers tightly closed when not in use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY STATEMENT

AGTROL CHEMICAL PRODUCTS warrants that the product conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions. THERE ARE NO OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING A WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE. This warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller and buyer assume all risk of any such use.

STORAGE AND DISPOSAL

See container label.

RE-ENTRY STATEMENT

Do not enter treated area without protective clothing until sprays have dried.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must include the following information: Inform workers of areas or fields that must not be entered without appropriate protective clothing until sprays have dried. In case of accidental exposure, wash with plenty of water. If there is any irritation in eyes after washing, get medical attention. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: CAUTION. Area treated with Agtrol. Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure, wash with plenty of water. If there is any irritation in eyes after washing, get medical attention.

NOTE: Gibberellic Acid is an extremely potent plant growth regulator. For best results, read all directions for use thoroughly. Consult your local experiment station specialist, distributor, or the Agtrol agricultural specialist in your area for the spray schedule best suited to your conditions.

GENERAL DIRECTIONS FOR USE

Discard any unused spray material at the end of each day. Prepare solution concentrations by mixing the required amount of product with water only in a clean spray tank. For best results, applications should be made during cooler parts of the day.

Use only as directed. Good spray practices should be followed. The label should be read thoroughly and understood before making applications. Effectiveness requires that all parts of plant or crop must receive spray or desired result will not occur, so spray thoroughly. When a range of rates is indicated, use the concentration and spray volume recommended locally.

Gibberellic Acid is a naturally occurring compound, produced by Agtrol Chemical Products in a biological process.

Data concerning the compatibility of GibGro with other agricultural compounds is not available.

SPRAY GUIDELINES FOR GRAPES

For all grapes, application is recommended by ground sprayer. Use 100 to 500 gallons as a dilute spray according to foliage density, or 30 to 80 gallons as a concentrate spray, unless specified otherwise. Do not exceed maximum rates. It is important to wet all berries thoroughly.

Thompson Seedless.

For cluster elongation ("Stretch"), looser cluster forms, and reducing cost of thinning, when used in conjunction with established girdling and thinning practices: apply 8 to 16 grams A before bloom when flower clusters are 3 to 5 inches long.*

For decreased berry set ("Thinning"), reducing hand-thinning costs, and hastened maturity: apply 8 to 16 grams A during bloom as one application or as two applications of equal amounts when the bloom period is extended with the second application made 3 to 7 days after the first application.*

For larger berries ("Sizing") and larger clusters when used in conjunction with established girdling and thinning practices: apply 32 to 80 grams A when average berry size is 4-5 millimeters in diameter or as two applications of equal amounts with the first made at or 2 to 3 days after shatter, followed during the next two weeks by the second application. Timing of the second spray will be dictated by experience in the vineyard to be sprayed and temperatures occurring during the interim between sprays. Potential effect will be reduced if the second spray occurs more than two weeks after the first application.*

* Refers to actual Gibberellic Acid. See Conversion Table to convert to amount of formulated GibGro needed.

THOMPSON SEEDLESS FOR RAISINS

For decreasing berry set, with increased raisin quality, and hastened maturity: apply 0.75 to 6 grams*/A when most bunches are in 60% to 80% bloom.

FLAME SEEDLESS

For decreased berry set ("Thinning") and reducing hand-thinning costs: apply 3 to 7.5 grams*/A during bloom. Higher amounts may cause an excess of shot berries or overthinning.

For larger berries ("Sizing") and larger clusters when used in conjunction with established girdling and thinning practices: apply 20 to 48 grams*/A as one application when berry diameter reaches 6 to 8 millimeters, or as two applications of equal amounts with the first made when berry diameter reaches 6 to 8 millimeters, followed during the next 5 to 10 days by the second application. Timing of the second spray will be dictated by experience in the vineyard to be sprayed and rate of berry growth during the interim between sprays.

PERLETTES

For larger berries ("Sizing") and larger clusters when used in conjunction with established girdling and thinning practices: Apply 32 to 48 grams*/A when average berry size is 4 to 5 millimeters in diameter, or as two applications of equal amount with the first made at 1 to 3 days after shatter, followed during the next two weeks by the second application. Timing of the second spray will be dictated by experience in the vineyard to be sprayed and temperatures occurring during the interim between sprays. Potential effect will be reduced if the second spray occurs more than two weeks after the first application.

* Refers to actual Gibberellic Acid. See Conversion Table to convert to amount of formulated GibGro needed.

OTHER SEEDLESS VARIETIES SUCH AS SEEDLESS TOKAY, INTERLOCKEN
SERIES AND RELATED HYBRIDS

For larger berries and larger clusters when used in conjunction with established girdling and thinning practices: apply 8 to 48 grams*/A as one application at or just after shatter (usually 2 to 3 days later) or as two applications of equal amounts not to exceed a total of 48 grams*/A, with the first made at or just after shatter, followed during the next two weeks by the second application. Timing of the second spray with split application will be dictated by experience in the vineyard to be sprayed and temperatures occurring during the interim between sprays. Potential effect will be reduced if the second spray occurs more than two weeks after the first application.

EMPEROR

For reducing berry shrivel. This can also increase berry size: apply 20 grams*/A as one application in 200 to 250 gallons/A approximately two weeks after completion of shatter following bloom. This timing should correspond to a period when the predominant berry diameter ranges from 10 to 15 millimeters.

* Refers to actual Gibberellic Acid. See Conversion Table to convert to amount of formulated GilGro needed.

BLACK CORINTH (ZANTE CURRANT)

For improving berry size: apply spray containing 1 to 8 grams*/A 3 to 5 days after full bloom, but before shatter begins.

SPRAY GUIDELINES FOR CITRUS

NAVEL ORANGES

(California) To delay aging of the rind and reduce rind disorders (e.g. rind staining, water spotting, sticky or tacky surface, puffy rind and rupture under pressure) and to produce a more orderly harvesting pattern: EARLY SPRAY (before any color change), apply to groves where harvest is not anticipated before March 1. The delay in rind aging is greatest when the early spray is applied before a color change. This spray timing produces the firmest rind possible. Apply one spray two weeks prior to color break which normally occurs August to November. Apply until February 15 in San Joaquin Valley, March 1 in Southern California. On large mature trees, apply 10 to 40 grams*/A in 400 to 500 gallons/A dilute, or 50 to 100 gallons/A concentrate.

NOTE: Do not apply to grove that may be harvested before March 1 as a reduction in grade may result due to the delayed coloring. Do not apply in white wash sprays in which lime or other caustic material has produced a high pH in the spray tank.

LATE SPRAY: (after marketable color is reached) apply to groves where harvest may be before March 1 (or not known). Apply one spray just after marketable color has developed which is normally from October through December. On large mature trees, apply 16 to 48 grams*/A in 400 to 500 gallons/A dilute with thorough coverage or 50 to 100 gallons/A concentrate.

NOTE: Sprays applied in late January/February may cause reduced production the following year. Do not apply within 10 days of harvest. Do not spray navel orange trees between February 15 and August 1.

* Refers to actual Gibberellic Acid. See Conversion Table to convert to amount of formulated GibGro needed.

VALENCIA ORANGES

(California) *To reduce rind creasing and to delay aging and softening of the rind:* apply a single spray in August or September to trees with a target crop of young fruit. On large mature trees, apply 40 to 80 grams* A in approximately 500 gallons/A dilute or 100 gallons A concentrate.

NOTE: Some increased regreening, or slower color development, should be expected in the target crop. Some increased regreening of mature fruit, if present, may occur.

LEMONS

(California) *To decrease the amount of small tree ripe fruit and to produce a more desirable production pattern in relation to market demand:* apply in a single spray when target crop is 1/2 to 3/4 full size but still green. Typically October through December. Use 20 grams* in 500 to 700 gallons/A on large, mature trees.

When applied two years in a row, an even larger difference in harvest pattern and maturity occurs.

NOTE: Do not apply within one month of harvest. Do not apply in spring or summer.

TANGERINE HYBRIDS

(Florida) *To increase fruit set and yield in tangerine hybrids with pollination problems such as the Orlando, Robinson, and Minneola:* apply spray during full bloom. Be sure to wet the leaves sufficiently.

Fruits are generally seedless: use 8 to 30 grams* in 400 to 500 gallons/A on large mature trees.

NOTE: A slight increase in mature leaf drop occurs at concentrations above 25 ppm. Fruit sizes may be reduced and color development slightly retarded.

* Refers to actual Gibberellic Acid. See Conversion Table to convert to amount of formulated GibGro needed.

(California) To delay disorders associated with rind aging of the *Minneola tangelo*; e.g., puffiness and softening and to increase peel strength: apply 20 to 40 grams* A in 400-500 gallons dilute spray two weeks prior to color break.

NOTE: Do not apply if early harvest is planned. Do not apply after coloring as pre-harvest rind staining may occur. Application during coloring may cause variation in rind color development.

GRAPEFRUIT

(Florida and Texas) To delay disorders associated with rind aging; e.g., puffiness, softening and orange coloration, to prevent pre-harvest drop of mature fruit and to increase peel strength and reduce water loss during storage: apply a single spray to fully colored fruit during the November through January period. Use 20 to 56 grams* in 500 to 700 gallons/A containing a suitable non-ionic surfactant at the the manufacturer's recommended rate. It is advisable to spot pick heavy crops to aid early marketing and to avoid reduction of yields which generally follow late held crops.

NOTE: Applications made after dormancy when trees begin to break dormancy may adversely affect crop. Do not use concentrated sprays. Results may vary due to seasonal and environmental conditions.

GRAPEFRUIT, STAR RUBY VARIETY

(Texas) To reduce early-season drop of small fruit of Star Ruby Variety thereby increasing yields: apply a single spray during the bloom period.

Use GibGro 4L (25 fluid ounces) or GibGro 2L (50 fluid ounces) (25 grams*) in 250 gallons water final spray mixture per acre. A suitable surfactant may be used to enhance efficacy.

NOTE: Do not tank-mix with other chemicals.

Do not apply concentrated solution.

Results may vary season to season depending on environmental conditions.

Maintain a well-balanced fertilization and watering program.

* Refers to actual Gibberellic Acid. See Conversion Table to convert to amount of formulated GibGro needed.

SPRAY GUIDELINES FOR FRUIT CROPS

BLUEBERRIES

For improving fruit set. For set problems due to insufficient natural honeybee pollination on varieties such as Coville, Jersey, Stanley, Earlieblue, Weymouth and others: make a single foliage spray application at full bloom (when over 75 percent of all flowers are fully open). For Weymouth, application can be delayed up to two weeks after full bloom to affect sizing of shot berries. Use GibGro 4L (80 fluid ounces) or GibGro 2L (160 fluid ounces) in 100 gallons of water. Use of a spreader-sticker is recommended. Apply to the point of run-off, thoroughly wetting all parts of the plant. Total gallonage will depend on size and density of the plants.

NOTE: Do not exceed 300 gallons/A. Although some varieties bloom closer to harvest than others - in no case should application be made closer than 40 days before harvest. Do not apply to plants in a low state of vigor.

SWEET CHERRIES

To delay harvesting, increase fruit weight, colored, firmer fruit and to increase size of fruit. Apply when the fruit is light green to start colored. Apply to the point of run-off wet the entire tree. Use 10 to 40 gallons/A on medium to large trees.

NOTE: Do not apply within one week of harvest.

RED TART CHERRIES

(All states except California) To maintain and extend high fruiting capacity of bearing tart cherry trees and reduce occurrence of "blind" nodes by stimulating lateral vegetative buds to develop a more productive balance of lateral shoots and spurs: apply a foliar spray containing GibGro 4L (4-8 fluid ounces) or GibGro 2L (8-16 fluid ounces) in 100 gallons finished spray from 14 to 28 days after bloom (or up to 14 days after shuck split). Use full coverage sprays of 50 to 150 gallons/A on medium to large bearing trees. Be sure entire trees receive good coverage. Use of a good horticultural wetting agent at the manufacturer's recommended rate will aid foliar wetting. GibGro must be applied annually to insure vegetative development and subsequent yield improvement year after year.

* Refers to actual Gibberellic Acid. See Conversion Table to convert to amount of formulated GibGro needed.

NOTE: GibGro works by affecting lateral bud differentiation which is apparent the year after application. Therefore, changes in shoot, spur and flower production will not be evident until 2 or 3 years after program initiation. Once this period is satisfied, response will be yearly provided annual applications have been made.

NOTE: Do not spray within one month of harvest. Adjust GibGro rate to complement vigor to trees. If trees are vigorous, use lowest recommended rates. Use higher rate for trees low in vigor and showing weak shoot and spur production. Excessive application rates on any tree will increase vegetative growth at the expense of fruit production the following year.

GibGro will not improve growth of trees under stress (nutritional, moisture, winter injury) or other factors inhibiting normal growth and development resulting from physical damage or unsound orchard practices. Best results from GibGro will be obtained when combined with good cultural practices.

YOUNG TART AND SWEET CHERRY TREES

(All states except California) *To reduce flowering and fruiting in young tart and sweet cherry trees to minimize the competitive effect of early fruiting on tree development:* apply GibGro liquid two to four weeks after bloom. Mix 25 to 40 ounces of GibGro 4L or 40 to 60 ounces of GibGro 1L in 100 gallons of water. Apply a foliar spray of 25 to 50 gallons per acre, assuming a tree density of 100 trees per acre equivalent, or apply about one quart of spray volume per tree.

Under conditions of low vigor, two applications are recommended. If two spray applications are made, allow at least a seven-day interval between sprays.

NOTE: DO NOT SPRAY TREES IN THE FIRST YEAR. Treat in the second season for reduction of flowering in the third season and again in the third season if reduction of flowering and fruiting is desired in the fourth season.

OLYMPUS STRAWBERRIES

(N.W. US Only; propagation stock) *To increase runner production of mother plants of the Olympus cultivar:* apply a single spray to mother plants 10 to 30 days after planting. At the time of spraying, plants should have 1 to 6 leaves. Apply 100 gallons/A to thoroughly wet new foliage to the point of run-off. Use 20 grams*/A.

NOTE: Not for use on fruiting plants. Treatments may not be effective on plantings set out after mid-May.

* Refers to actual Gibberellic Acid. See Conversion Table to convert to amount of formulated GibGro needed.

FORCING RHUBARB

To increase yield of marketable forced rhubarb and to break dormancy on plants receiving insufficient chilling: apply 2 fluid ounces (60 ml) of a solution containing 20 grams* in 10 gallons to each cleaned crown, when the rest period is not completely broken. When the rest period is broken by cold weather, apply 2 fluid ounces (60 ml) of a solution containing 10 grams* in 10 gallons.

NOTE: Keep forcing house temperatures at 40° to 50°F for 24 hours after application. If house is warmer than 50°F, the crowns should be covered with plastic. Temperatures in the forcing house above 50°F will result in lower yields and poor stalk color.

SPRAY GUIDELINES FOR VEGETABLE CROPS

ARTICHOKES

(California) To accelerate maturity of artichokes and to shift the harvest to an earlier date: apply spray at bud initiation time, normally six weeks prior to anticipated harvest. Be sure the entire plant (leaves, stems and buds) are covered to prevent run-off. Use 10 grams* in 100 to 125 gallons/A.

NOTE: Do not apply within seven days of harvest.

CELERY

To increase plant height and yield, and overcome stress due to cold weather conditions, or saline soils and to obtain earlier maturity: apply spray one to four weeks prior to harvest. Lower concentrations are applied at the three to four-week interval. Higher concentrations at the one to two-week interval. Use 2.5 to 10 grams* in 25 to 50 gallons/A.

NOTE: Do not apply earlier than four weeks before harvest as Gibberellic Acid may induce bolting (seed stalk formation).

Applications made less than one week preharvest may result in residues.

Celery plants must be harvested when mature to ensure quality.

* Refers to actual Gibberellic Acid. See Conversion Table to convert to amount of formulated GibGro needed.

LETTUCE FOR SEED

To obtain uniform bolting and increase seed production, apply the following spray schedule:

Growth Stage	ppm*	g.*/A	Gallons/Acre
4 leaf stage	10	0.4	10
8 leaf stage	10	1.6	20
12 leaf stage	10	4	100

NOTE: Do not feed crop wastes to livestock.

SEED POTATOES

To stimulate uniform sprouting - for maximum production, more uniform development, fewer late maturing plants, and to break dormancy of newly harvested potatoes that have not had a full rest period: Dip freshly dug seed pieces in a solution containing 0.2 to 0.4 gram* in 100 gallons prior to planting.

NOTE: If soil temperature is very high, avoid treating rested seed and use the minimum concentration for dormant soil.

SPINACH

(All States except California) To facilitate harvest, increase yield and improve quality of fall and over-winter spinach: apply a single spray 10 to 14 days before each anticipated harvest on fall or over-winter spinach, ideally when daytime temperatures are 40° to 70°F and during early morning hours when dew is present on crop.

Use GibGro 4L (6 to 8 fluid ounces) or GibGro 2L (12 to 16 fluid ounces) - 6 to 8 grams*/A. Mix in 10 to 50 gallons/A by ground sprayer or in a minimum of 5 to 10 gallons/A by air.

Maximum benefit from GibGro is obtained when below normal temperatures predominate following application and growth would be otherwise slowed in untreated spinach.

NOTE: Since Gibberellic Acid can promote bolting, do not apply to spinach after the mid-winter period or if temperatures may be expected to exceed 75°F within several days of application. Do not apply on spring-planted spinach.

* Refers to actual Gibberellic Acid. See Conversion Table to convert to amount of formulated GibGro needed.

SPRAY GUIDELINES FOR FLORICULTURAL CROPS

POMPOM CHRYSANTHEMUMS

(Florida) *For elongating peduncles on pom-pom chrysanthemums:* apply a single spray 4 to 5 weeks after initiation of short day conditions.

Use GibGro 4L (1/2 to 1 fluid ounce) or GibGro 2L (1 to 2 fluid ounces) - (1/2 to 1 gram*) in 12 gallons for application to 1,000 sq. ft. of bed (20 to 40 fluid ounces GibGro 4L or 40 to 80 fluid ounces GibGro 2L equivalent to 20 to 40 grams* in 500 gallons/A).

Apply with overhead nozzles directing the spray to the flower buds.

NOTE: Overuse or incorrect timing may cause long, spindly and weak stems.

STATICE

(Florida) *To promote earlier flowering and to increase flower yield:* apply a single drench spray when plants are more than 10 inches in diameter (approximately 20 to 22 days after normal seeding time). Use 40 to 50 grams* in 15 gallons to provide 10 ml (1/2 oz) solution per plant.

NOTE: Do not exceed specified rates. For best results, repeated sprays. Accelerated flowering is induced by extended photoperiod, adequate nutrition and reduced night temperature. Treatment with gibberellins lessens the requirement for the cold requirement and/or the long photoperiod.

SPRAY GUIDELINES FOR ADDITIONAL CROPS

BERMUDAGRASS GOLF TURF

(Florida) *To initiate or maintain growth and prevent color change during periods of cold stress and light frosts on golf course Bermudagrass (e.g., Tifdwarf, Tifgreen, etc.):* apply 10 grams* weekly or 25 grams* biweekly in 25 to 100 gallons/A.

Use GibGro 4L (1/4 to 2/3 fluid ounces) or GibGro 2L (1/2 to 1 1/3 fluid ounces) - (1/4 to 2/3 gram*) in approximately 6 gallons appropriate for the spray equipment for application to 1,000 sq. ft. (10 1/2 to 26 1/2 fluid ounces/A GibGro 4L or 21 to 53 fluid ounces/A GibGro 2L equivalent to 10 to 25 grams*/A in 25 to 100 gallons/A).

* Refers to actual Gibberellic Acid. See Conversion Table to convert to amount of formulated GibGro needed.

NOTE: Do not exceed specific rates.

Do not apply during extended warm periods where night temperatures exceed 65°F.

Maintain adequate moisture and proper fertilization programs recommended in local area.

Discontinue treatments if thinning is observed.

Do not apply the high rate more frequently than every two weeks. More frequent mowing may be necessary.

Do not use on dormant turf.

HOPS

(For seeded and seedless Fuggle hops and similar varieties adapted to Oregon and the Northwest) *To increase yield and pickability:* apply spray when vine growth is five to eight feet in length. Use 4 to 6 grams* in 100 to 150 gallons/A.

NOTE: Do not apply within three weeks of harvest.

SUGAR CANE

(Hawaii) Use GibGro 20% or GibGro 5% Powder. *For increase in sucrose yield:* apply 28 to 84 grams* in 7 to 10 gallons/A of spray by airplane. Uniform coverage is essential for maximum response. Use 56 grams as a single treatment, or 28 grams two or three times in separate applications with 30 to 45 day intervals. Application may be made to cane during the first and/or second year of culture. Young cane should be at least three months old to avoid possible tiller reduction. Application should not be made less than 4 months prior to harvest.

Application should be made when growth rate is depressed by temperature. Cane grown below 1,500 feet elevation will benefit from applications made during November through March.

Cane should be treated when there is sufficient soil moisture from rain or irrigation to sustain a high growth rate for at least 30 days following each treatment. Lack of water will negate treatment effect.

* Refers to actual Gibberellic Acid. See Conversion Table to convert to amount of formulated GibGro needed.

Four forms of GibGro brand Gibberellic Acid are now available to better serve the needs of individual growers.

GibGro 20% Powder (110 grams/bottle)
Active Ingredient:
Gibberellic Acid*..... 20% W/W
Equivalent to 32 grams* of Gibberellic Acid per bottle.
EPA Reg. No. 55146-53

GibGro 5% Powder (320 grams bottle)
Active Ingredient:
Gibberellic Acid*..... 5% W/W
Equivalent to 16 grams* of Gibberellic Acid per bottle.
EPA Reg. No. 55146-56

GibGro 4I Liquid Concentrate
(20 fl. oz/bottle and 128 fl. oz bottle)
Active Ingredient:
Gibberellic Acid*.....4% W/W
Equivalent to approximately 1.0 gram* of Gibberellic Acid
per fluid ounce of product.
EPA Reg. No. 55146-55

GibGro 2L Liquid Concentrate
(128 fl. oz/bottle)
Active Ingredient:
Gibberellic Acid*.....2% W/W
Equivalent to approximately .5 gram* of Gibberellic Acid
per fluid ounce of product.
EPA Reg. No. 55146-57

CONVERSION TABLE POWDERS

GRAMS OF ACTUAL GIBBERELIC ACID PER ACRE	AMOUNT OF GIBGRO POWDER FORMULATION PER ACRE	
Desired Actual Gibberellic Acid Concentration (Grams*) in Finished Spray (Per Acre)	GibGro 20% Powder Contains (2.0 Grams*/10 Grams Formulated Product)	GibGro 5% Powder Contains (0.5 Grams*/10 Grams Formulated Product)
0.5	2.5 grams	10.0 oz.
1.0	5 grams	20 oz.
2.0	10 grams	40 oz.
4.0	20 grams (1/8 bottle)	80 oz.
5.0	25 grams	100 oz.
8.0	40 grams (1/4 bottle)	160 oz.
10.0	50 grams	200 oz.
12.0	60 grams	240 oz.
16.0	80 grams (2/5 bottle)	320 oz.
20.0	100 grams	400 oz.
25.0	125 grams	500 oz.
32.0	160 grams (2/3 bottle)	640 oz.
40.0	200 grams	800 oz.
48.0	240 grams (1 1/2 bottle)	960 oz.
50.0	250 grams	1000 oz.

* Refers to actual Gibberellic Acid.

CONVERSION TABLE LIQUIDS

GRAMS OF ACTUAL GIBBERELIC ACID PER ACRE TO AMOUNT OF GIBGRO LIQUID FORMULATION PER ACRE

Desired Actual Gibberellic Acid Concentration (Grams*) in Finished Spray (Per Acre)	GibGro 2L Liquid Contains (0.5 Grams*/Fluid Ounce of Formulated Product)	GibGro 4L Liquid Contains (1.0 Gram*/Fluid Ounce of Formulated Product)
0.5	1 oz.	0.5 oz.
1.0	2 oz.	1 oz.
2.0	4 oz.	2 oz.
4.0	8 oz.	4 oz.
5.0	10 oz.	5 oz.
8.0	16 oz.	8 oz.
10.0	20 oz.	10 oz.
12.0	24 oz.	12 oz.
16.0	32 oz.	16 oz.
20.0	40 oz.	20 oz.
25.0	50 oz.	25 oz.
32.0	64 oz.	32 oz.
40.0	80 oz.	40 oz.
48.0	96 oz.	48 oz.
50.0	100 oz.	50 oz.

* Refers to actual Gibberellic Acid.

NOTICE TO USER

Seller makes no warranty, express or implied, of merchantability, fitness or otherwise concerning use of this product other than as indicated on the label. User assumes all risks of use, storage or handling not in strict accordance with accompanying directions.