

(A)	United States Environmental Protection Agency Office of Pesticide Programs (H7505C) Washington, DC 20460 Application for Pesticide:	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number Please Attach An Original Form
-----	---	--	--

Section I		
1. Company/Product Number 55146-57	2. EPA Product Manager C. Giles-Parker	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Nufarm Champion Formula 2 Dry Flowable Fungicide	PM# 22	
5. Name and Address of Applicant (Include ZIP Code) Nufarm Inc. Agent for Nufarm Americas Inc., AGT Division 1333 Burr Ridge Parkway, Suite 125A Burr Ridge, IL 60527 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. Product Name	

Section II	
<input type="checkbox"/> Amendment - Explain Below <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application <input type="checkbox"/> Other - explain below

NOTIFICATION
NOV 30 2004

Explanation: Use additional page(s) if necessary. (For Section I and Section II).

Notification to add marketing statement to mango.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Sections 12 and 14 of FIFRA.

Section III			
1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No <small>*Certification must be submitted.</small>	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If "Yes," Unit Package Wt. No. Per Container	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If "Yes," Package Wt. No. Per Container	2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label or <input checked="" type="checkbox"/> Container		4. Size(s) of Retail Container 1 qt.; 1/2, 1, 2 1/2, 5, 30, 55, 110, 220 gal.; bulk	5. Location of Label Directions <input checked="" type="checkbox"/> On Label or <input checked="" type="checkbox"/> On Labeling accompanying product
6. Manner In Which Label Is Affixed To Product <input checked="" type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Other (Self-Adhesive Integrated Label/Booklet) <input checked="" type="checkbox"/> Paper Glued <input type="checkbox"/> Stenciled			

Section IV		
1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Theodore D. Head	Title Product Registration Manager / Agriculture	Telephone No. (Include Area Code) (708) 764-3230

Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.	6. Date Application Received (Stamped)
Signature 	3. Title Product Registration Manager / Agriculture
4. Typed Name Theodore D. Head	5. Date 11-15-04

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the WPS.

Do not enter or allow worker entry into treated areas during the REI of 24 hours provided the following instructions are followed:

For at least 7 days following application an eye-flush container, designed specifically for flushing eyes, must be available at the WPS decontamination site for workers entering the area treated with copper hydroxide.

Notify workers of the application by warning them orally that residues in the treated areas may be highly irritating to their eyes and to take precautions such as refraining from rubbing their eyes and if they get residues in their eyes they should immediately flush their eyes using the eye-flush container.

PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated, such as plants, soil, or water, is long sleeved shirt and coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the WPS for agricultural pesticides 40 CFR part 170. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Keep unprotected persons out of treated area until sprays have dried.

FROST INJURY PROTECTION: Bacterial Ice Nucleation Inhibitor: Application of Champ® Dry Prill made to all crops listed on this label at rates and stages of growth indicated below just prior to anticipated frost conditions will afford control of ice nucleating bacteria (*Pseudomonas Syringae*, *Erwinia Herbicola* and *Pseudomonas Fluorescens*) and may therefore provide protection against light frost. Use higher rates when bacterial infection is severe. Not recommended in those geographical areas where weather conditions favor severe frost.

GENERAL INSTRUCTIONS

CHAMP Dry Prill can be used with all types of spraying equipment. The volume per acre will differ depending on the specific crop and the equipment used. Use CHAMP Dry Prill as per instructions on this label.

APPLYING SPRAY MIXTURE: The directions given under each crop are for applying dilute spray mixture unless otherwise shown. The amount of CHAMP Dry Prill applied per acre in concentrate and aerial sprays should be the same as the amount applied per acre in dilute sprays. The required amount should be mixed with enough water to thoroughly cover the crop with spray mixture and is to be applied to the point of "runoff". The volume of water needed will depend upon the spray equipment used and the size of the crop being sprayed. Read the information below about applying dilute, concentrate, and aerial sprays.

APPLYING DILUTE HIGH-VOLUME SPRAYS: On vegetable crops use 25 to 100 gallons of spray mixture per acre; on fruit and nut trees use 250 to 800 gallons per acre. As much as 1,500 gallons per acre may be needed for large trees.

APPLYING CONCENTRATE GROUND SPRAYS: On vegetable crops use 5 to 20 gallons of spray mixture per acre; on fruit and nut trees use 25 to 250 gallons per acre.

APPLYING AERIAL SPRAYS: Use 3 to 30 gallons per acre.

NOTE: This product may be reactive on metal and masonry surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.

NOTE: Do not tank mix CHAMP Dry Prill with Alette® fungicide unless appropriate precautions have been taken to buffer the spray solution or severe phytotoxicity may result.

3/10
NOTE: CHAMP Dry Prill should not be applied in a spray solution having a pH of less than 6.5 as phytotoxicity may occur.

NOTE: Environmental conditions such as extended periods of wet weather, acid rain, etc. which alter the pH of the leaf surface may affect the performance of CHAMP Dry Prill resulting in possible phytotoxicity or loss of effectiveness.

NOTE: Reduced effect on pests or crop injury may result from tank mixing agricultural chemicals especially where several products are involved. Unless recommended on this label or by state/local expert, or the user has small scale direct experience, tank mixing should not be undertaken.

NOTE: Agricultural chemicals may be reactive with soft metals and some synthetic materials such as plastics, rubbers, etc. When working with equipment containing these materials the equipment must be thoroughly flushed with clean water after each day's use.

CHEMIGATION: Apply this product only through center pivot, motorized lateral move, end tow, traveler, big gun, plastic solid set, or plastic hand move sprinkler irrigation systems that do not contain aluminum components. Do not apply this product through any other type of irrigation system unless specifically set forth above or as may be specified in the future as additional systems not containing aluminum components come into use. Crop injury, lack of effectiveness or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), backflow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. For nonpublic water sprinkler chemigation systems, the system must contain a functional checkvalve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shutdown. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. For non-public water sprinkler chemigation systems, the irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when windspeed favors drift beyond the area intended for treatment.

It is recommended that the pesticide supply tank be equipped with a means for continuous agitation either by recirculation or a mechanical agitator. Charge the supply tank with the appropriate amount of water and add the pesticide slowly followed by any sticker-spreaders, insecticides, nutrients, etc. Observe all cautions and limitations on the label of all products used in the mixtures. For fixed position irrigation systems such as center pivot, big gun, etc., the pesticide should be applied towards the end of the irrigation period. Exact timing will depend on the desired pesticide application rate and calibration of the system. For moving systems, the pesticide should be applied continuously. In all cases, thorough coverage of the crop should be achieved.

Posting of areas to be chemigated is required when 1) any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, day care centers,

4/10

hospitals, inpatient clinics, nursing homes or any public areas such as schools, parks, playgrounds, or other public facilities not including public roads, or 2) when the chemigated area is open to the public such as golf courses or retail greenhouses.

Posting must conform to the following requirements. Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive areas. When there are no usual points of entry, signs must be posted in the corners of the treated areas and in any other location affording maximum visibility to sensitive areas. The printed side of the sign should face away from the treated area towards the sensitive area. The signs shall be printed in English. Signs must be posted prior to application and must remain posted until foliage has dried and soil surface water has disappeared. Signs may remain in place indefinitely as long as they are composed of materials to prevent deterioration and maintain legibility for the duration of the posting period.

All words shall consist of letters at least 2 1/2 inches tall, and all letters and the symbol shall be a color which sharply contrasts with their immediate background. At the top of the sign shall be the words KEEP OUT, followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word STOP. Below the symbol shall be the words PESTICIDES IN IRRIGATION WATER.

This sign is in addition to any sign posted to comply with the WPS.

NOTE: IRRIGATION SYSTEMS AND ASSOCIATED PIPING SHOULD BE THOROUGHLY FLUSHED WITH CLEAN WATER FOLLOWING APPLICATION OF COPPER BASED FUNGICIDES. FLUSHING MUST BE DONE IN A MANNER WHICH WILL NOT WASH THE PRODUCT FROM THE FOLIAGE AND REDUCE DISEASE CONTROL.

No additional surfactants are needed unless specified for an individual crop. Add CHAMP Dry Prill to the spray tank followed by any stickerspreaders, insecticides, nutrients, etc. Observe all cautions and limitations on the label of all products used in mixtures. The specific instructions given on this label are based on general applications and circumstances. The recommendations of the State Agricultural Extension Service should be closely followed as to timing, frequency and number of sprays per season.

NOTE: APPLICATION TO PLANT SURFACES SPRAYED WITH AND WHICH HAVE LOW pH CHEMICAL RESIDUE MAY ALSO RESULT IN CROP INJURY.

BERRIES, VINES AND HOPS

BLACKBERRY

(Santiam, Logans, Boyesen, Marion, Aurora, Cascade, Chehalem and Thornless Evergreen) *Leaf Spot, Cane Spot, Purple Blotch, Anthracnose, Yellow Rust, and Pseudomonas Blight:* apply delayed dormant spray after training in Spring at 2 2/3 pounds per acre. Make Fall spray application after harvest. Add 1 quart of crop oil per acre. *Leaf Spot, Cane Spot, Purple Blotch, Anthracnose, and Yellow Rust:* apply when leaf buds begin to open and repeat when flower buds show white at 1 2/5 pounds per acre. Add 1 quart of crop oil per acre.

NOTE: Crop injury may occur if applied to foliage under certain conditions such as hot or prolonged moist periods. Discontinue applications if signs of crop injury appear.

BLUEBERRY

(Except California) *Bacterial Canker:* apply at 3 3/4 to 4 2/3 pounds per acre. Make first application before the Fall rains, preferably the first week in October and a second application four weeks later.

CRANBERRY

Fruit Rot: apply at 5 3/5 pounds per acre beginning in late bloom. One or two additional applications made at 10 to 14 day intervals may be required, depending on disease pressure. *Rose Bloom:* apply three sprays of 5 3/5 pounds per acre CHAMP Dry Prill at 10 to 14 day intervals as soon as symptoms are observed. *Bacterial Stem Canker:* apply 5 3/5 pounds per acre post harvest and again in the Spring before bud burst. One or two additional applications at 10 to 14 day intervals may be required depending on disease severity. *Tip Blight (Monilinia), Stem Blight, Leaf Blight, Red Leaf Spot:* apply at 5 3/5 pounds per acre as a delayed dormant spray in the Spring. Repeat at 10 to 14 day intervals as needed through pre-bloom. *Upright Dieback:* apply at 5 1/3 pounds per acre as a prebloom application. A second application can be made 10 to 14 days later if required.

CURRANT & GOOSEBERRY

Anthracnose, Leaf Spot: make three applications at 7 pounds per acre, starting after harvest, before bloom and after petal fall. Continue on a 10 to 14 day schedule during wet conditions in the Spring.

GRAPE

Black Rot, Phomopsis, Powdery Mildew and Downy Mildew: apply

1 1/3 to 2 2/3 pounds per acre. Begin application at bud break with additional applications made throughout the season depending upon disease severity.

NOTE: Slight to severe foliage injury may occur in copper-sensitive varieties such as Concord, Delaware, Niagara and Rosette. Use lower rate of CHAMP Dry Prill and test for sensitivity when treating these varieties or others to be sensitive to copper. Hydrated lime may be added at a rate up to 1/2 pound per 100 gallons of spray solution to decrease the severity of phytotoxicity. Mix CHAMP Dry Prill and water first before adding lime or incompatibility may occur.

HOPS

Downy Mildew: apply 1 1/3 pounds per acre as a fungicide crown treatment after pruning, but before training. After training, additional fungicide treatments are needed at about 10 day intervals.

NOTE: Discontinue use two weeks before harvest.

RASPBERRY

(Except California) *Leaf and Cane Spot, Purple Blotch, Anthracnose, Yellow Rust, Pseudomonas Blight:* apply at 2 2/3 pounds per acre as a delayed dormant spray after training in the Spring. Make a Fall application after harvest. Add one quart of crop oil per acre. *Leaf and Cane Spot, Purple Blotch, Anthracnose, Yellow Rust:* apply at 1 2/5 pounds per acre when leaf buds begin to open and repeat when flower buds show white. Add one quart of crop oil per acre.

NOTE: Crop injury may occur if applied to foliage under certain conditions such as hot or prolonged moist periods. Discontinue applications if signs of crop injury appear.

STRAWBERRY

Leaf Spot & Leaf Blight: apply 1 1/3 to 2 pounds per acre. Begin application when plants are established and continue on a weekly schedule throughout season.

NOTE: Discontinue applications if signs of phytotoxicity appear.

FIELD CROPS

ALFALFA

Cercospora & Leptosphaerulina Leaf Spots: apply 1 1/3 pounds per acre 10 to 14 days before each harvest or earlier if disease threatens.

NOTE: Spray injury may occur with sensitive varieties such as Lahontan.

PEANUT

Cercospora Leaf Spot: begin spraying 35 to 40 days after planting or when disease symptoms first appear. Apply at 1 to 2 pounds per acre. Continue applications at 10 to 14 day intervals. One to two quarts of 6 pounds per gallon flowable sulfur may be added. Reduce spray interval to seven days during humid weather. Use higher rates when conditions favor disease.

SUGAR BEET

Cercospora Leaf Spot: start spray when disease threatens and continue for four to five applications. Spray every 10 to 14 days depending on weather conditions at 1 1/3 to 3 1/3 pounds per acre depending on disease severity.

WHEAT, BARLEY, OATS

Septoria Leaf Blotch & Helminthosporum Spot Blotch: apply 1 to 1 1/3 pounds per acre. Make first application by early heading and follow with second application 10 days later or as necessary. Use higher rates when conditions favor disease.

TREE CROPS

ALMOND, APRICOT, CHERRY, PLUM & PRUNE

Coryneum Blight (Shot Hole) (Stigmata carpophila), Bacterial Canker, Blossom Brown Rot, Dead Bud (Pseudomonas syringae), Bacterial Blast (Pseudomonas): use 5 1/3 to 8 pounds CHAMP Dry Prill per acre as a dormant application before foliage buds swell. For CHERRIES, where disease is severe, an additional application at leaf fall may be required. **ALMOND ONLY:** for *Bacterial Blast* control in sprinkler irrigated orchards or where disease is severe, apply 3/4 pound per acre post-bloom, at two week intervals or just prior to sprinkling. *Coryneum Blight (Shot Hole) (Stigmata carpophila), Blossom Brown Rot:* for early bloom (popcorn) application prior to full bloom, apply at 4 to 5 1/2 pounds per acre.

NOTE: To avoid plant injury, do not use above rate after full bloom. For *blast* control in sprinkler irrigated orchards or where disease is severe, apply 2 to 4 CHAMP Dry Prill sprays, or as many as required, at 1 to 2 pounds per acre at two week post bloom intervals or just before sprinkling.

NOTE: In sensitive varieties of ALMONDS, such as Peerless, Mission, and Neplus slight leaf injury may occur from post bloom spray.

APPLE

Anthracnose, European Canker, Blossom Blast, Shoot Blast (Pseudomonas): apply before Fall rains at 8 to 10 1/2 pounds per acre.

NOTE: Use on yellow varieties may cause discoloration. To avoid, pick before spraying.

Fireblight, Scab: apply at 5 1/2 to 10 1/2 pounds per acre. Make application as a full cover spray between silver-tip and green-tip.

NOTE: Phytotoxicity may occur from late application. After green-tip apply at 2/3 pound per acre. *Crown or Collar Rot*: mix 2 3/4 pounds in 100 gallons of water. Apply 4 gallons of suspension as a drench on the lower trunk area of each tree. Apply either in early Spring or in Fall after harvest each year.

NOTE: Do not use if soil pH is below 5.5 or copper toxicity may result.

AVOCADO

Anthracnose, Blotch, Scab: apply when bloom buds begin to swell at 5 1/3 to 8 pounds per acre. Continue application at monthly intervals for five to six applications. Use higher rate when conditions favor disease.

CITRUS

Melanose, Scab, Algal Spot: apply 2 3/4 to 8 pounds per acre, depending on disease severity as a pre-bloom and post-bloom spray. *Greasy Spot, Pink Pitting*: apply 1 1/3 to 4 pounds per acre using higher rates when conditions favor disease. *Brown Rot*: apply 2 2/3 to 5 1/3 pounds per acre beginning in Fall and continuing as needed. Apply to skirts of trees to a height of at least four feet. Use higher rates when conditions favor disease.

NOTE: In California, in areas subject to copper injury, add 1/3 to 1 pound of high quality lime per pound of CHAMP Dry Prill. *Citrus Canker (SUPPRESSION ONLY)*: apply 8 pounds per acre, spraying flushes 7 to 14 days after shoots begin to grow. Young fruit may require an additional application. Number and timing of applications will be dependent on disease pressure. Under heavy disease pressure, each flush of new growth should be sprayed. *Alternaria Brown Spot (SUPPRESSION ONLY)*: apply 5 1/3 to 6 2/3 pounds per acre to susceptible varieties on the first flush in the Spring and every additional flush. Application to fruit should start after two-thirds of the petals have fallen and be repeated at 21 day intervals. *Phytophthora Foot Rot*: mix 2/3 pound of CHAMP Dry Prill with 1 gallon of water and paint trunks of trees from the soil surface to the lowest scaffold limbs. Apply in May prior to Summer rains and/or in the Fall prior to wrapping trees for freeze protection. Treatment serves as protection for up to one year, but does not cure existing infections. *Phytophthora Brown Rot, Septoria Spot*: apply 2 2/3 to 5 1/3 pounds per acre to the entire tree in the Fall before or just after the first rain and continue as needed.

NOTE: In California in areas subject to copper injury, add 1/3 to 1 pound of high quality lime per pound of CHAMP Dry Prill.

NOTE: Do not use CHAMP Dry Prill on Citrus seedlings grown in greenhouses or shadehouses.

CITRUS FIELD NURSERY GROWN

Melanose, Scab, Greasy Spot, Pink Pitting, Brown Rot and Citrus Canker (SUPPRESSION ONLY): apply 1 1/3 pounds of CHAMP Dry Prill in 100 gallons of water (2 2/3 to 5 1/3 pounds per acre) at 28 day intervals.

FILBERT

Bacterial Blight: apply 10 2/3 to 16 pounds per acre as a post-harvest spray. In seasons of heavy rainfall, apply another spray when three-fourths of the leaves have dropped. Add 1 pint of superior-type oil per 100 gallons of water. *Eastern Filbert Blight*: apply 10 2/3 to 16 pounds per acre in sufficient water to obtain thorough coverage. Make initial application at budswell to budbreak. Additional sprays should be made on a 10 to 14 day interval depending on disease severity or when conditions are conducive for disease development. Add 1 pint of superior-type oil per 100 gallons of water.

KIWIFRUIT

Blossom Blight (Bud Rot), Leaf Spot (Phomopsis), Erwinia herbicola, Pseudomonas syringae, Pseudomonas fluorescens: make two to three applications at 1 1/3 to 2 pounds per acre during dormant season. Do not apply at time of or after leaf emergence.

MACADAMIA

(Except California) *Blossom Blight & Raceme Blight, Anthracnose*: apply 3 to 6 pounds per acre depending on disease pressure in 50 to 300 gallons of water during peak raceme development and bloom periods. For aerial application apply 3 to 6 pounds per acre in 10 to 30 gallons of water.

OLIVE

(California) *Peacock Spot, Olive Knot*: make first application at 5 1/3 to 8 pounds per acre before Winter rains fall. A second application in early Spring should be made if disease is severe.

PEACH, NECTARINE

Leaf Curl, Coryneum Blight (Shot Hole), Bacterial Canker, Bacterial Blast (Pseudomonas), Bacterial Blight (Xanthomonas): apply 5 1/3 to 10 2/3 pounds per acre after leaf fall as a dormant application. Use the higher rate when rainfall is very heavy and disease pressure is high. May be used with an agricultural spray oil. *Blossom Brown Rot, Leaf Curl, Coryneum Blight (Shot Hole)*: apply 5 1/3 to 8 pounds per acre as a full cover spray at pink bud. Application at this time affords some control of *Leaf Curl* and *Coryneum Blight*. *Bacterial Spot*: apply 5 1/3 pounds per acre as a dormant spray.

(Except California) As a post-bloom spray: apply up to 1/2 pound per acre at first and second cover sprays.

NOTE: Do not apply three weeks prior to harvest. Use only recommended rates. Spotting of leaves and defoliation may occur from use in cover sprays.

PEAR

Fireblight: apply at 2/3 pound per acre at five day intervals throughout bloom period. *Pseudomonas Blight*: apply before Fall rains at a rate of 8 to 10 2/3 pounds per acre and again at dormant before Spring growth starts.

NOTE: Excessive dosages may cause fruit russet.

PECAN

Shuck Rot, Kernel Rot (Phytophthora cactorum), Zonate Leaf Spot (Cristulariella pyramidalis) (SUPPRESSION ONLY): apply 1 1/3 to 2 2/3 pounds per acre at two to four week intervals when kernel growth begins through shuck opening. Apply in sufficient water to ensure thorough coverage.

PISTACHIO

Botrytis Blight, Botryosphaeria Panicle and Shoot Blight, Septoria Leaf Blight, Late Blight (Alternaria alternata): apply 2 2/3 to 5 1/3 pounds per acre beginning at budswell. Repeat at 14 to 28 day intervals depending on disease conditions. If disease conditions are severe, use the high rate and the short spray interval.

QUINCE

Fire Blight: apply 2/3 pound per acre at five day intervals throughout the bloom period. Apply in sufficient water to provide thorough coverage.

WALNUT

Walnut Blight: apply 5 1/3 to 8 pounds per acre. Apply first spray at early pre-bloom prior to or when catkins are partially expanded. Make additional applications during bloom and early nutlet stage as needed. Additional applications may be necessary when frequent rainfall occurs.

NOTE: When applied as a dilute spray, 1 pint of summer oil emulsion may be added per 100 gallons of spray. Adequate control may not be obtained when copper tolerant species of *Xanthomonas* bacteria are present.

TROPICAL CROPS

BANANA

Sigatoka: apply by air at 1 1/3 pounds per acre in 3 gallons of water containing 1/2 gallon agricultural oil. Apply on a 14 day schedule throughout the wet season. Apply at 21 day intervals during dry periods. *Black Pitting*: apply at 2 2/3 pounds per 100 gallons directly to the fruit stem and include the basal portion of the leaf crown. Apply during the first and second weeks after fruit emergence.

CACAO

Black Pod: begin applications at the start of the rainy season and continue while infection conditions persist. Sprays should be made as often as 14 to 21 days in high rainfall areas at varying rates from 1 1/3 to 5 1/3 pounds per acre depending on disease severity. For drier areas, where two to four applications are recommended during critical infection periods and at long intervals, use 4 to 5 2/3 pounds per acre, according to disease pressure incidence and planting density.

COFFEE

Coffee Berry Disease (Collectotrichum coffeanum): apply first spray at 4 to 5 1/3 pounds per acre after flowering and before onset of long rains and then at 21 to 28 days interval until picking. Use higher rates when rainfall is heavy and disease pressure is high. *Bacterial Blight (Pseudomonas syringae)*: apply 4 to 5 1/3 pounds per acre. Begin spray program before onset of the long rains and continue throughout the rainy season at 14 to 21 day intervals. The critical time of spraying to control this disease is just before, during, and after flowering(s), especially when coinciding with wet weather. Use the higher rates when rainfall is heavy and disease pressure is high. *Leaf Rust (Hemileia vastatrix)*: apply 1 1/3 to 2 2/3 pounds per acre before the onset of rain and then at 21 day intervals while the rains continue. Use higher rates when rainfall is heavy and disease

pressure is high. *Iron Spot (Cercospora coffeicola)* and *Pink Disease (Corticium salmonicolor)*: apply 1 1/3 pounds per acre as a concentrate or dilute spray. Begin treatment at the start of wet season and continue at monthly intervals for three applications.

GUAVA

Anthracoze, Red Algae: apply 2 pounds per acre beginning just prior to flowering and repeat weekly until just prior to harvest.

LITCHI

Anthracoze: apply 2 pounds per acre beginning just prior to flowering and repeat weekly until just prior to harvest.

MAMEY SAPOTE

Anthracoze, Algal Leaf Spot: apply 4 to 5 pounds per acre when conditions favor disease development. Repeat at 14 to 30 day intervals as needed.

MANGO

(Florida & Puerto Rico) *Anthracoze*: apply monthly after fruit set until harvest at 5 1/3 to 6 2/3 pounds per acre.

PAPAYA

Anthracoze: apply 2 to 6 2/3 pounds per acre beginning before disease is expected to appear. Repeat at 10 to 14 day intervals or at 5 to 7 day intervals during periods of heavy rainfall. Use the higher rates when conditions favor disease. The addition of a suitable spreader-sticker may be desirable especially during periods of heavy rains.

PASSION FRUIT

Anthracoze: apply 4 pounds per acre beginning just prior to flowering and repeat weekly.

SUGAR APPLE (Annona)

Anthracoze: apply 8 pounds per acre beginning just prior to flowering and repeat weekly.

VEGETABLE CROPS

BEAN (dry, green)

Brown Spot, Bacterial Blight (Halo & Common): for protective sprays, apply first application when plants are six inches high. Apply on 7 to 14 day schedule depending on local conditions. Use 2/3 to 2 pounds per acre depending on disease severity.

CRUCIFERS

Broccoli, Brussels Sprout, Cabbage, Cauliflower, Kale, Collard Greens, Mustard Greens and Turnip Greens

Black Rot (Xanthomonas), Black Leaf Spot (Alternaria), Downy Mildew: apply 1/3 to 2/3 pound per acre at seven day intervals after transplants are set in the field. Use higher rate when conditions favor disease.

NOTE: Reddening of older leaves may occur on broccoli at the higher rate and flecking of wrapper leaves may occur on cabbage.

CUCURBITS

Cantaloupe, Cucumber, Honeydew, Muskmelon, Pumpkin, Squash, and Watermelon

Alternaria Leaf Spot, Angular Leaf Spot, Anthracnose, Downy Mildew, Powdery Mildew, Gummy Stem Blight, Watermelon Bacterial Fruit Blotch (SUPPRESSION): apply at 1 1/3 pounds per acre beginning when conditions are favorable for disease development and repeat at 5 to 7 day intervals, as needed depending on disease severity.

NOTE: Crop injury may occur from application at shorter intervals. Discontinue use if injury occurs.

CARROT

Alternaria Leaf Spot, Carrot Blight (Cercospora): when disease threatens apply 1 1/3 pounds per acre at 7 to 14 day intervals depending on disease severity.

CELERY & CELERIAC

Cercospora Early Blight, Septoria Late Blight & Bacterial Blights: apply as soon as plants are first established in the field at 1 1/3 pounds per acre, then every 5 to 7 days depending on disease severity and weather.

EGGPLANT

(Except California) *Alternaria Blight, Anthracnose, Phomopsis*: use 1 1/3 pounds per acre before disease appears. Repeat at 7 to 10 day intervals.

ENDIVE, ESCAROLE

Downy Mildew: apply 2/3 to 1 1/3 pounds per acre. Begin treatment when disease first appears and repeat every 7 to 10 days as needed to suppress disease.

GARLIC, LEEK, ONION

Purple Blotch & Downy Mildew: apply at 1 1/3 pounds per acre when plants are 4 to 6 inches high and repeat at 7 to 10 day intervals. *Bacterial Blight*: apply 2/3 to 1 pound per acre.

LETTUCE

Downy Mildew: apply 2/3 to 1 1/3 pounds per acre. Begin treatment when disease first appears and repeat every 7 to 10 days as needed to suppress disease.

PEA

Powdery Mildew: begin spray treatment when disease symptoms first appear. Use 1 to 2 pounds per acre according to disease severity. Repeat applications at weekly intervals.

PEPPER

Bacterial Spot: when disease threatens, apply 1 1/3 to 2 pounds per acre in sufficient water for adequate coverage at 5 to 10 day intervals depending on disease severity.

POTATO

Early Blight & Late Blight: apply at 3 to 10 day intervals starting when plants are six inches high. Apply 2/3 to 1 pound per acre in those locations where disease is light and up to 2 to 2 2/3 pounds per acre where disease is severe. *Colorado Potato Beetle (SUPPRESSION ONLY)*: application of CHAMP Dry Prill at rates and timing recommended for control of early and late blight may provide suppression of the Colorado Potato Beetle.

SPINACH

Anthracoze, Cercospora Leaf Spot, Downy Mildew, White Rust Blue Mold: apply 1 1/3 to 2 2/3 pounds per acre. Begin treatment when disease first appears and repeat every 7 to 10 days as needed to suppress disease.

NOTE: Flecking may occur on spinach leaves.

TABLE BEET, BEET GREENS

Cercospora Leaf Spot: apply 1 1/3 to 2 2/3 pounds per acre when conditions favor disease. Repeat treatment at 10 to 14 day intervals as needed. The addition of an agricultural spray oil is recommended.

TOMATO

Early Blight, Late Blight: when disease threatens, apply 1 1/3 to 2 pounds per acre at 7 to 10 day intervals, or as necessary. *Bacterial Speck*: apply at 1 1/3 pounds per acre at 10 to 30 day intervals beginning when the disease threatens. Use more frequent applications when disease pressure is high. *Bacterial Spot, Anthracnose, Gray Leaf Mold & Septoria Leaf Spot*: when disease threatens, apply 1 1/3 to 2 2/3 pounds per acre at 7 to 10 day intervals, more frequently when disease is severe.

WATERCRESS

Cercospora Leaf Spot: apply 1 1/3 pounds per acre when plants are established in the field. Repeat at 7 to 14 day intervals up to four applications per crop in at least 50 gallons of water per acre.

MISCELLANEOUS

ATEMOYA

Anthracoze: apply 2 pounds per acre just prior to flowering and repeat weekly until just prior to harvest.

CARAMBOLA

Anthracoze: apply 4 pounds per acre just prior to flowering and repeat weekly until just prior to harvest.

CHIVES

Downy Mildew: apply 1 1/3 pounds per acre when plants are established in the field. Repeat at 7 to 10 day intervals as needed.

DILL

Phoma Leaf Spot, Rhizoctonia Foliage Blight: apply 1 3/4 pounds per acre when plants are established in the field. Repeat at 7 to 10 day intervals as needed.

DOUGLAS FIR

Rhabdocline Needlecast: apply 1 1/3 pounds per acre at bud break and repeat at three to four week intervals. Apply in a tank mix with other registered pesticide if disease pressure is moderate to severe.

GINSENG

Alternaria Leaf & Stem Blight: CHAMP Dry Prill may be applied at 1 3/4 pounds per acre as a tankmix with two pounds Iprodione 50WP in 100 gallons of water per acre. Begin Iprodione 50WP-CHAMP Dry Prill applications as soon as plants have emerged in Spring. Applications should be repeated every seven days until plants become dormant in Fall. Apply fungicides at least eight hours before rain, giving the fungicides time to dry on the plants. Use of a spreader-sticker is advised.

NOTE: *Alternaria Leaf & Stem Blight* is most severe in humid conditions such as those found in the dense canopies of two, three and four year old ginseng. Complete and thorough spray coverage is required for control.

PARSLEY

Bacterial Blight (Pseudomonas spp.): apply 2 pounds per acre when plants are first established in the field and repeat at 5 to 7 day intervals.

PERSIMMON

Cercospora Leaf Spot: apply 1 1/3 pounds per acre beginning in May/June, during leaf flush, and repeat at 14 day intervals throughout the season depending on disease severity.

TURFGRASS

For algae control in turfgrass in areas such as sod farms, golf courses, cemeteries, home lawns, and industrial or municipal turf areas (including parks, playgrounds, athletic fields). Apply 1 pound of CHAMP Dry Prill per 1,000 square feet in 5 gallons of water to control algae. CHAMP Dry Prill may be used alone or in combination with other registered fungicides as a maintenance spray.

NOTE: Phytotoxicity may occur depending upon varietal differences. If injury occurs discontinue use. Do not apply in spray solutions with a pH of less than 6.5.

GREENHOUSE AND SHADEHOUSE CROPS

CHAMP Dry Prill may be used in greenhouses and shadehouses to control diseases on crops listed on this label. Specific directions are provided below for certain crops and the grower should be aware that the sensitivity of crops grown under such conditions differ greatly from field conditions. The user must determine if CHAMP Dry Prill can be used safely prior to commercial application by testing a small area and observing the results for 7 to 10 days.

One level tablespoon of CHAMP Dry Prill per 1,000 square feet is equivalent to 1 pound per acre. Begin application at first sign of disease and repeat at 7 to 14 day intervals as needed.

CUCUMBER

Angular Leaf Spot, Downy Mildew: apply 1 1/4 to 1 1/2 tablespoons weekly when plants begin to vine.

EGGPLANT

Alternaria Blight, Anthracnose, Phomopsis: apply 1 1/2 tablespoons at first sign of disease and repeat at 7 to 14 day intervals as needed.

PEPPER

Bacterial Spot: apply 1 1/2 to 2 1/4 tablespoons when conditions first favor disease and at 5 to 10 day intervals as needed.

TOMATO

Anthracnose, Bacterial Speck, Bacterial Spot, Early Blight, Grey Leaf Mold, Late Blight, Septoria Leaf Spot: apply 1 1/2 to 2 1/4 tablespoons when conditions first favor disease and at 7 to 10 day intervals as needed.

NOTE: Do not use CHAMP Dry Prill on Citrus seedlings grown in greenhouses or shadehouses.

ORNAMENTALS

PECAN, LIVE OAK

(Texas and Florida) Ball Moss: apply at 5 1/2 pounds per 100 gallons of water, in Spring after heavy rain, using 1 1/2 gallons of spray per foot of tree height. Make sure to wet tufts thoroughly. A second application may be required after 12 months.

NOTE: CHAMP Dry Prill maybe injurious to ornamentals grown under live oaks.

PHILODENDRON

Bacterial Leaf Spot: apply weekly before disease appears at 1 pound of CHAMP Dry Prill per 100 gallons of water.

SYCAMORE

Anthracnose: make two applications using 1 1/3 to 2 pounds per 100 gallons as a full cover spray. Make first application at bud crack and second application 7 to 14 days later at 10% leaf expansion.

For control of bacterial and fungal diseases on foliage, flowers, and stems of ornamentals grown in Greenhouses, Shadehouses, Fields and Nurseries (container, bench or bed-grown): apply CHAMP Dry Prill at 2/3 pound per 100 gallons as a full cover spray beginning at first sign of disease. Repeat at intervals of 7 to 14 days (or shorter) depending on rainfall and disease severity. Due to the large number of species, widely varying growth conditions, and varieties of ornamentals and nursery plants it is impossible to test every one for sensitivity to CHAMP Dry Prill. The user should apply the recommended rate of CHAMP Dry Prill in a small area and check for any symptoms of phytotoxicity in 7 to 10 days prior to large-scale application.

Do not tank mix with Aliette® fungicide without buffering the spray solution.

One half tablespoon of CHAMP Dry Prill per gallon of water is equivalent to .9 pounds per 100 gallons.

AGLAONEMA

Bacterial Leaf Spot

ALTHEA (Rose of Sharon)

Bacterial Leaf Spot

ARALIA

Xanthomonas & Cercospora Leaf Spots, Alternaria

ARBORVITAE

Alternaria Twig Blight, Cercospora Leaf Spot

AZALEA

(a) Cercospora Leaf Spot, Botrytis Blight, Phytophthora Dieback, Powdery Mildew

BEGONIA

Bacterial Leaf Spot (Xanthomonas spp., Erwinia spp., Pseudomonas spp.)

BOSTON FERN

Bacterial Leaf Spot

BOUGAINVILLEA

Anthracnose, Bacterial Leaf Spot

BULBS

(EASTER LILY (b), TULIP), Botrytis Blight, Anthracnose

CAMELIA

Anthracnose, Bacterial Leaf Spot

CAMPHOR TREE

Pseudomonas Leaf Spot

CANNA

Pseudomonas Leaf Spot

CARNATION

(a) Alternaria Blight, Pseudomonas Leaf Spot & Botrytis Blight

CHINESE TALLOW TREE

Bacterial Leaf Spot (Xanthomonas spp., Pseudomonas spp.)

CHRYSANTHEMUM

(a) Septoria Leaf Spot, Botrytis Blight

COTONEASTER

Botrytis Blight

DAHLIA

Alternaria Leaf Spot, Cercospora Leaf Spot, Botrytis Grey Mold

DATE PALM

Pestalotia Leaf Spot

DIANTHUS

Bacterial Spot, Bacterial Soft Rot

DOGWOOD

Anthracnose

DRACAENA

Bacterial Leaf Spot

DUMB CANE

Bacterial Leaf Spot

DUSTY MILLER

Bacterial Leaf Spot (Pseudomonas cichorii spp.)

ECHINACEA

Botrytis Blight

ELM (Drake)

Xanthomonas Leaf Spot

EUONYMUS

Botrytis Blight, Anthracnose

EUROPEAN FAN PALM

Pestalotia Leaf Spot

GARDENIA

Alternaria Leaf Spot, Cercospora Leaf Spot, Botrytis Bud Rot

GERANIUM

Alternaria Leaf Spot, Cercospora Leaf Spot, Botrytis Grey Mold

GLADIOLUS

Alternaria Leaf Spot, Botrytis Grey Mold, Bacterial Leaf Blight, Botrytis Blight, Anthracnose

GOLDEN RAIN TREE*Alternaria Leaf Spot, Botrytis Grey Mold, Bacterial Leaf Blight***GRAPE IVY***Bacterial Leaf Spot***HIBISCUS***(c) Bacterial Leaf Spot***HOLLY FERN***Pseudomonas Leaf Spot***HONEY LOCUST***Bacterial Leaf Spot***IMPATIENS***Bacterial Leaf Spot***INDIA HAWTHORN***(d) Anthracnose, Entomosporium Leaf Spot***IRIS***Bacterial Leaf Spot***IVY (English, Algerian)***(a) Xanthomonas Leaf Spot***IXORA***Xanthomonas Leaf Spot***JUNIPER (Eastern Red Cedar)***Anthracnose***LANTANA***Bacterial Leaf Spot***LILAC***Cercospora Leaf Spot***LOBLOLLY BAY***Anthracnose***LOQUAT***Entomosporium maculata, Colletotrichum spp.***MAGNOLIA (Southern)***Anthracnose, Bacterial Leaf Spot, Algal Leaf Spot***MAGNOLIA (Sweet Bay)***Anthracnose***MAGNOLIA***Bacterial Leaf Spot***MANDEVILLAS***Anthracnose***MARIGOLD***Alternaria Leaf Spot, Botrytis Leaf Rot, Flower Rot, Cercospora Leaf Spot***MULBERRY CONTORTED***Bacterial Leaf Spot***MULBERRY, WEEPING***Bacterial Leaf Spot***NEPHTHYTIS***Bacterial Leaf Spot***OLEANDER***Bacterial Leaf Spot, Fungal Leaf Spot***OAK, LAUREL***Algal Leaf Spot (Cephaleuros virescens spp.)***PACHYSANDRA***Volutella Leaf Blight***PANSY***Downy Mildew***PARLOR PALM***Bacterial Leaf Spot***PEAR (flowering)***Fire Blight, Leaf Spot***PENTAS (Egyptian Star)***Bacterial Leaf Spot (Xanthomonas spp.)***PEONY***Botrytis Blight***PERIWINKLE***Phomopsis Stem Blight***PHLOX***Alternaria Leaf Spot***PHOTINIA (Red Tip, Red Leaf)***Anthracnose, Entomosporium***PISTACHIO***Anthracnose***PLANTAIN LILY***Bacterial Leaf Spot***POTHOS***Bacterial Leaf Spot***POWDER PUFF PLANT***Bacterial Leaf Spot***PURPLE OSIER WILLOW***Anthracnose***PYRACANTHA***Fireblight, Scab***QUEEN PALM***Exosporium Leaf Spot, Phytophthora Bud Rot***RHODODENDRON***Alternaria Flower Spot***ROSE***(a) Powdery Mildew, Black Spot***SNAPDRAGON***Anthracnose, Dieback, Downy Mildew***SPATHE FLOWER***Bacterial Leaf Spot***TATARIAN HONEYSUCKLE***Bacterial Leaf Spot***UMBRELLA TREE***Bacterial Leaf Spot***VERBENA***Xanthomonas Leaf Spot***VIBURNUM***Anthracnose***WASHINGTON PALM***Pestalotia Leaf Spot***WEEPING FIG***Bacterial Leaf Spot***WEEPING WILLOW***Anthracnose***YUCCA (ADAMS NEEDLE)***Cercospora & Septoria Leaf Spots*

- (a) On some varieties a discoloration may occur on foliage or blooms. To prevent residues on commercial plants, do not spray just before selling season.
- (b) Apply 2 to 3 1/2 pounds of CHAMP Dry Prill in 20 to 100 gallons of water per acre.
- (c) Hibiscus - Do not apply to plants in flower.
- (d) For India Hawthorn use 1 2/5 to 2 3/4 pounds per 100 gallons or 3/4 to 1 1/2 level tablespoons per gallon.

9/10

CHAMP® Dry Prill

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER - PELIGRO

Corrosive. Causes irreversible eye damage. Do not get in eyes. Wear protective eyewear (goggles, face shield or safety glasses). Harmful if absorbed through skin, swallowed or inhaled. Avoid contact with skin or clothing. Avoid breathing dust, vapor or spray mist. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- a) long-sleeved shirt and long pants
- b) chemical resistant gloves made of any waterproof material, such as polyvinyl chloride, nitrile rubber, or butyl rubber
- c) shoes plus socks
- d) protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- After each day of use, wash in detergent and hot water any personal clothing worn while using this product.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites.

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest Environmental Protection Agency Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY STATEMENT

Nufarm Americas Inc. warrants that the product conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions. THERE ARE NO OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING A WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE. This warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller and buyer assumes all risk of any such use.

Champ® is a registered trademark of Nufarm Americas Inc.

Aliette® is a registered trademark of Rhone Poulenc



Nufarm Americas Inc.
1333 Burr Ridge Parkway, Suite 125A
Burr Ridge, IL 60527-0866
Phone: 708.754.3330 Fax: 708.754.0314
www.us.nufarm.com

10/10

November 15, 2004

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504C)
U. S. Environmental Protection Agency
Room 266A, Crystal Mall 2
1801 S. Bell Street
Arlington, VA 22202-4501

Attention: Ms. Cynthia Giles-Parker, PM 22
Herbicide Branch

Subject: Notification to add marketing statement to
Nufarm Champion Formula 2 Dry Flowable,
EPA Reg. No. 55146-57

Dear Cynthia:

Nufarm is requesting, via notification, to add the word Puerto Rico to our Mango section of the label. Our marketing group has brought my attention that Mangos are grown in Puerto Rico as well as Florida.

Please note that no use rates or application sites have changed. We are simply adding one word to the label, Puerto Rico.

If you should have any questions, please contact me at the address and telephone number above or by e-mail to <ted.head@us.nufarm.com>. Thank you for your cooperation.

Sincerely,

A handwritten signature in black ink, appearing to read 'Theodore D. Head'.

Theodore D. Head
Product Registration Manager

