

55146-57

10-24-2007

117



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Mary Beth Endres
Registration Manager
Nufarm Americas, Inc. AGT division
150 Harvester Drive, Suite 200
Burr Ridge, IL 60527

OCT 24 2007

Subject: Champ Dry Pill (EPA Reg. No. 55146-57)
Label Amendment Submission Dated May 10, 2007

Dear Ms. Endres:

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, to reformat use directions from paragraph to table format, add Florida specific recommendation for Citrus Canker suppression as recommended by the University of Florida, and adds new company warranty statement, is acceptable provided the following changes are made:

- Non-Agricultural Use Requirements Box, delete "this product" from the sentence beginning with "Keep unprotected persons...."
- Field Crop Table, for potato (colorado potato beetle), insert the same rate for potato (early blight & late blight) or merge the cells in rate per acre column for potato.
- Turfgrass Table, within the parentheses following "turfgrass", replace the period after "sodfarms" with a comma.
- In the last sentence of the "Limitation of Liability" statement on p.16, add "To the extent consistent with applicable law," in front of "Buyer's or user's exclusive remedy ..."

One copy of the label stamped "Accepted with comments" is enclosed for your records. This label supersedes all labels previously accepted for this product. Please submit one copy of the final printed label that incorporates the required changes before the product is released for shipment. Should you have any questions or concerns, please contact Tawanda Spears via telephone at 703.308.8050 or e-mail at spears.tawanda@epa.gov.

Sincerely yours,

A handwritten signature in black ink that reads "Tawanda Spears for".

Tony Kish
Product Manager (22)
Fungicide Branch
Registration Division (7505P)

2/17

Champ[®] Dry Prill

AGRICULTURAL FUNGICIDE / BACTERICIDE

ACCEPTED
with COMMENTS
In EPA Letter Dated

OCT 24 2007

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

55146-57

ACTIVE INGREDIENT:

Copper Hydroxide*

57.6%

OTHER INGREDIENTS:

42.4%

TOTAL: 100.0%

*(Metallic Copper Equivalent 37.5%)

**KEEP OUT OF REACH OF CHILDREN
DANGER - PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 55146-57
EPA EST. NO. 62562-CHL-2

MANUFACTURED FOR
NUFARM AMERICAS INC.
AGT DIVISION
BURR RIDGE, IL 60527



NET CONTENTS LBS.

MADE IN CHILE

055146-57.20070510.PENDING

3/17

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER - PELIGRO**

Corrosive. Causes irreversible eye damage. Do not get in eyes. Wear protective eyewear (goggles, face shield or safety glasses). Harmful if absorbed through skin, swallowed or inhaled. Avoid contact with skin or clothing. Avoid breathing dust, vapor or spray mist. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- a) long-sleeved shirt and long pants
- b) chemical-resistant gloves made of any waterproof material, such as polyvinyl chloride, nitrile rubber, or butyl rubber
- c) shoes plus socks
- d) protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS	
Users Should:	
<ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. • Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. • After each day of use, wash in detergent and hot water any personal clothing worn while using this product. 	

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites.

4/17

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours provided the following instructions are followed:

For at least 7 days following application, an eye-flush container, designed specifically for flushing eyes, must be available at the Worker Protection Standard decontamination site for workers entering the area treated with copper hydroxide.

Notify workers of the application by warning them orally that residues in the treated areas may be highly irritating to their eyes and to take precautions such as refraining from rubbing their eyes and if they get residues in their eyes they should immediately flush their eyes using the eye-flush container.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is long-sleeved shirt and coveralls, chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber, or butyl rubber, shoes plus socks and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard (WPS) for agricultural pesticides 40 CFR part 170. The Worker Protection Standard (WPS) applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep unprotected persons out of treated area until sprays have dried this product.

FROST INJURY PROTECTION: Bacterial Ice Nucleation Inhibitor: Application of this product made to all crops listed on this label at rates and stages of growth indicated below just prior to anticipated frost conditions will afford control of ice nucleating bacteria (*Pseudomonas Syringae*, *Erwina Herbicola* and *Pseudomonas Fluorescens*) and may therefore provide protection against light frost. Use higher rates when bacterial infection is severe. Not recommended in those geographical areas where weather conditions favor severe frost.

GENERAL INSTRUCTIONS

This product can be used with all types of spraying equipment. The volume per acre will differ depending on the specific crop and the equipment used. Use this product as per instructions on this label.

APPLYING SPRAY MIXTURE: The directions given under each crop are for applying dilute spray mixture unless otherwise shown. The amount of this product applied per acre in concentrate and aerial sprays should be the same as the amount applied per acre in dilute sprays. The required amount should be mixed with enough water to thoroughly cover the crop with spray mixture and is to be applied to the point of runoff. The volume of water needed will depend upon the spray equipment used and the size of the crop being sprayed. Read the information below about applying dilute, concentrate, and aerial sprays.

APPLYING DILUTE HIGH-VOLUME SPRAYS: On vegetable crops use 25 to 100 gallons of spray mixture per acre; on fruit and nut trees use 250 to 800 gallons per acre. As much as 1,500 gallons per acre may be needed for large trees.

APPLYING CONCENTRATE GROUND SPRAYS: On vegetable crops use 5 to 20 gallons of spray mixture per acre; on fruit and nut trees use 25 to 250 gallons per acre.

APPLYING AERIAL SPRAYS: Use 3 to 30 gallons per acre.

NOTES:

- This product may be reactive on metal and masonry surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.
- Do not tank mix this product with Aliette® fungicide unless appropriate precautions have been taken to buffer the spray solution or severe phytotoxicity may result.
- This product should not be applied in a spray solution having a pH of less than 6.5 as phytotoxicity may occur.
- Environmental conditions such as extended periods of wet weather, acid rain, etc. which alter the pH of the leaf surface may affect the performance of this product resulting in possible phytotoxicity or loss of effectiveness.
- Reduced effect on pests or crop injury may result from tank mixing this product especially where several products are involved. Unless recommended on this label or by state/local expert, or the user has small scale direct experience, tank mixing should not be undertaken.
- Agricultural chemicals may be reactive with soft metals and some synthetic materials such as plastics, rubbers, etc. When working with equipment containing these materials the equipment must be thoroughly flushed with clean water after each day's use.

CHEMIGATION: Apply this product only through center pivot, motorized lateral move, end tow, traveler, big gun, plastic solid set, or plastic hand move sprinkler irrigation systems that do not contain aluminum components. Do not apply this product through any other type of irrigation system unless specifically set forth above or as may be specified in the future as additional systems not containing aluminum components come into use. Crop injury, lack of effectiveness or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent, in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. For nonpublic water sprinkler chemigation systems, the system must contain a functional checkvalve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shutdown. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. For non-public water sprinkler chemigation systems, the irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when windspeed favors drift beyond the area intended for treatment.

It is recommended that the pesticide supply tank be equipped with a means for continuous agitation either by recirculation or a mechanical agitator. Charge the supply tank with the appropriate amount of water and add the pesticide slowly followed by any sticker-spreaders, insecticides, nutrients, etc. Observe all cautions and limitations on the label of all products used in the mixtures. For fixed position irrigation systems such as center pivot, big gun, etc., the pesticide should be applied towards the end of the irrigation period. Exact timing will depend on the desired pesticide application rate and calibration of the system. For moving systems, the pesticide should be applied continuously. In all cases, thorough coverage of the crop should be achieved.

Posting of areas to be chemigated is required when 1) any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, day care centers, hospitals, inpatient clinics, nursing homes or any public areas such as schools, parks, playgrounds, or other public facilities not including public roads, or 2) when the chemigated areas is open to the public such as golf courses or retail greenhouses.

Posting must conform to the following requirements: Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive areas. When there are no usual points of entry, signs must be posted in the corners of the treated areas and in any other location affording maximum visibility to sensitive areas. The printed side of the sign should face away from the treated area towards the sensitive area. The signs shall be printed in English. Signs must be posted prior to application and must remain posted until foliage has dried and soil surface water has disappeared. Signs may remain in place indefinitely as long as they are composed of materials to prevent deterioration and maintain legibility for the duration of the posting period.

All words shall consist of letters at least 2-1/2 inches tall, and all letters and the symbol shall be a color which sharply contrasts with their immediate background. At the top of the sign shall be the words KEEP OUT, followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word STOP. Below the symbol shall be the words PESTICIDES IN IRRIGATION WATER.

This sign is in addition to any sign posted to comply with the Worker Protection Standard.

NOTE: IRRIGATION SYSTEMS AND ASSOCIATED PIPING SHOULD BE THOROUGHLY FLUSHED WITH CLEAN WATER FOLLOWING APPLICATION OF COPPER BASED FUNGICIDES. FLUSHING MUST BE DONE IN A MANNER WHICH WILL NOT WASH THE PRODUCT FROM THE FOLIAGE AND REDUCE DISEASE CONTROL.

No additional surfactants are needed unless specified for an individual crop. Add this product to the spray tank followed by any sticker-spreaders, insecticides, nutrients, etc. Observe all cautions and limitations on the label of all products used in mixtures. The specific instructions given on this label are based on general applications and circumstances. The recommendations of the State Agricultural Extension Service should be closely followed as to timing, frequency and number of sprays per season.

NOTE: APPLICATION TO PLANT SURFACES SPRAYED WITH AND WHICH HAVE LOW pH CHEMICAL RESIDUE MAY ALSO RESULT IN CROP INJURY.

6/17

BERRIES, VINES AND HOPS

CROP	DISEASE	RATE PER ACRE	USE INSTRUCTIONS
BLACKBERRY (Santiam, Logans, Boysen, Marion, Aurora, Cascade, Chehalem and Thornless Evergreen)	Leaf Spot, Cane Spot, Purple Blotch, Anthracnose, Yellow Rust, and Pseudomonas Blight.	2-2/3 pounds	Apply delayed dormant spray after training in Spring. Make Fall spray application after harvest. Add 1 quart of crop oil per acre.
	Leaf Spot, Cane Spot, Purple Blotch, Anthracnose, and Yellow Rust.	1-2/5 pounds	Apply when leaf buds begin to open and repeat when flower buds show white. Add 1 quart of crop oil per acre.
	NOTE: Crop injury may occur if applied to foliage under certain conditions such as hot or prolonged moist periods. Discontinue applications if signs of crop injury appear.		
BLUEBERRY (Except California)	Bacterial Canker	3-3/4 to 4-2/3 pounds	Make first application before the Fall rains, preferably the first week in October and a second application four weeks later.
CRANBERRY	Fruit Rot	5-3/5 pounds	Apply beginning in late bloom. One or two applications made at 10 to 14 day intervals may be required, depending on disease pressure.
	Rose Bloom	5-3/5 pounds	Make three applications at 10 to 14 day intervals as soon as symptoms are observed.
	Bacterial Stem Canker	5-3/5 pounds	Apply post harvest and again in the Spring before bud burst. One or two additional applications at 10 to 14 day intervals may be required depending on disease severity.
	Tip Blight (Monilinia), Stem Blight, Leaf Blight, Red Leaf Spot	5-3/5 pounds	Apply as a delayed dormant spray in the Spring. Repeat at 10 to 14 day intervals as needed through prebloom.
	Upright Dieback	5-1/3 pounds	Apply as a prebloom application. A second application can be made 10 to 14 days later if required.
CURRANT & GOOSEBERRY	Anthracnose, Leaf Spot	7 pounds	Make three applications starting after harvest, before bloom and after petal fall. Continue on a 10 to 14 day schedule during wet conditions in the Spring.
GRAPE	Black Rot, Phomopsis, Powdery Mildew, Downy Mildew	1-1/3 to 2-2/3 pounds	Begin application at bud break with additional applications made throughout the season depending on disease severity.
	NOTES: <ul style="list-style-type: none"> • Slight to severe foliage injury may occur in copper-sensitive varieties such as Concord, Delaware, Niagara and Rosette. Use lower rate of this product and test for sensitivity when treating these varieties or others known to be sensitive to copper. • Hydrated lime may be added at a rate of up to 1/2 pound per 100 gallons of spray solution to decrease the severity of phytotoxicity. Mix this product and water first before adding lime or incompatibility may occur. 		
HOPS	Downy Mildew	1-1/3 pounds	Apply as a fungicide crown treatment after pruning, but before training. After training, additional fungicide treatments are needed at about 10 day intervals.
	NOTE: Discontinue use 2 weeks before harvest.		

7/17

BERRIES, VINES AND HOPS (continued)

CROP	DISEASE	RATE PER ACRE	USE INSTRUCTIONS
RASPBERRY (Except California)	Leaf and Cane Spot, Purple Blotch, Anthracnose, Yellow Rust, Pseudomonas Blight	2-2/3 pounds	Apply as a delayed dormant spray after training in the Spring. Make a Fall application after harvest. Add one quart of crop oil per acre.
	Leaf and Cane Spot, Purple Blotch, Anthracnose, Yellow Rust	1-2/5 pounds	Apply when leaf buds begin to open and repeat when flower buds show white. Add one quart of crop oil per acre.
	NOTE: Crop injury may occur if applied to foliage under certain conditions such as hot or prolonged moist periods. Discontinue applications if signs of crop injury appear.		
STRAWBERRY	Leaf Spot & Leaf Blight	1-1/3 to 2 pounds	Begin application when plants are established and continue on a weekly schedule throughout season.
	NOTE: Discontinue applications if signs of phytotoxicity appear.		

FIELD CROPS

CROP	DISEASE	RATE PER ACRE	USE INSTRUCTIONS
ALFALFA	Cercospora & Leptosphaerulina Leaf Spots	1-1/3 pounds	Apply 10 to 14 days before each harvest or earlier if disease threatens.
	NOTE: Spray injury may occur with sensitive varieties such as Lahontan.		
PEANUT	Cercospora Leaf Spot	1 to 2 pounds	Begin spraying 35 to 40 days after planting or when disease symptoms first appear. Continue applications at 10 to 14 day intervals. One to two quarts of six pounds per gallon flowable sulfur may be added. Reduce spray interval to seven days during humid weather. Use higher rates when conditions favor disease.
POTATO	Early Blight & Late Blight	2/3 to 1 pound or 2 to 2-2/3 pounds	Apply at 3 to 10 day intervals starting when plants are 6 inches high. Apply the lower rate in those locations where disease is light and the higher rate where disease is severe.
	Colorado Potato Beetle (Suppression Only)		Application of this product at rates and timing recommended for control of <i>early blight</i> and <i>late blight</i> may provide suppression of the Colorado Potato Beetle.
SUGARBEET	Cercospora Leaf Spot	1-1/3 to 3-1/3 pounds	Start spray when disease threatens and continue for 4 to 5 applications. Spray every 10 to 14 days depending on weather conditions and depending on disease severity.
WHEAT, BARLEY, OATS	Septoria Leaf Blotch & Helminthosporum Spot Blotch	1 to 1-1/3 pounds	Make first application by early heading and follow with second application 10 days later or as necessary. Use higher rates when conditions favor disease.

8/17

TREE CROPS

CROP	DISEASE	RATE PER ACRE	USE INSTRUCTIONS
ALMOND, APRICOT, CHERRY, PLUM, PRUNE	Coryneum Blight [Shot Hole] (<i>Stigmina carpophila</i>), Bacterial Canker, Blossom Brown Rot, Dead Bud (<i>Pseudomonas syringae</i>), Bacterial Blast (<i>Pseudomonas</i>)	5-1/3 to 8 pounds	Use as a dormant application before foliage buds swell. For CHERRIES , where disease is severe, an additional application at leaf fall may be required. ALMOND ONLY: for Bacterial Blast control in sprinkler irrigated orchards or where disease is severe, apply 3/4 pound per acre postbloom, at two week intervals or just prior to sprinkling.
	Coryneum Blight [Shot Hole] (<i>Stigmina carpophila</i>), Blossom Brown Rot	4 to 5-1/2 pounds	Early bloom (popcorn) application prior to full bloom.
	NOTES: <ul style="list-style-type: none"> • To avoid plant injury, do not use above rate after full bloom. For blast control in sprinkler irrigated orchards or where disease is severe, apply 2 to 4 sprays or as many as required, at 1 to 2 pounds per acre at two week post-bloom intervals or just before sprinkling. • In sensitive varieties of ALMONDS, such as Peerless, Mission, and Neplus slight leaf injury may occur from post-bloom spray. 		
APPLE	Anthracnose, European Canker, Blossom Blast, Shoot Blast (<i>Pseudomonas</i>)	8 to 10-1/2 pounds	Apply before Fall rains.
	Fireblight, Scab	5-1/2 to 10-1/2 pounds	Make application as a full cover spray between silver-tip and green-tip. NOTE: Phytotoxicity may occur from late application. After green-tip apply at 2/3 pound per acre.
	Crown or Collar Rot	2-3/4 pounds in 100 gallons of water	Apply 4 gallons of suspension as a drench on the lower trunk area of each tree. Apply either in early Spring or in Fall after harvest each year. NOTE: Do not use if soil pH is below 5.5 or copper toxicity may result.
NOTE: Use on yellow varieties may cause discoloration. To avoid, pick before spraying.			
AVOCADO	Anthracnose, Blotch, Scab	5-1/3 to 8 pounds	Apply when bloom buds begin to swell. Continue application at monthly intervals for five to six applications. Use higher rate when conditions favor disease.
CITRUS	Melanose, Scab, Algal Spot	2-3/4 to 8 pounds	Apply, depending on disease severity, as a pre-bloom and post-bloom spray.
	Greasy Spot, Pink Pitting	1-1/3 to 4 pounds	Apply using higher rates when conditions favor disease.
	Brown Rot	2-2/3 to 5-1/3 pounds	Apply beginning in Fall and continuing as needed. Apply to skirts of trees to a height of at least 4 feet. Use higher rates when conditions favor disease.

9/17

TREE CROPS (continued)

CROP	DISEASE	RATE PER ACRE	USE INSTRUCTIONS
CITRUS (continued)	Citrus Canker (Suppression Only)	1-1/4 to 8 pounds	<p>General Recommendations: Apply 8 pounds per acre, spraying flushes 7 to 14 days after shoots begin to grow. Young fruit may require an additional application. Number and timing of applications will be dependent on disease pressure. Under heavy disease pressure, each flush of new growth should be sprayed.</p> <p>Florida Specific Recommendations: Begin applications to protect new leaf flushes. Repeat at 14 to 21 day intervals, or more often if needed, depending on disease pressure and environmental conditions. It is important to protect all subsequent leaf flushes throughout the year. Young fruit may require an additional application.</p> <p>Under dry weather conditions and low disease pressure, use 1-1/4 to 2-1/2 pounds per acre.</p> <p>Under conditions of wet weather and high disease pressure, higher rates may be required (4 to 8 pounds per acre).</p>
	Alternaria Brown Spot (Suppression Only)	5-1/3 to 6-2/3 pounds	Apply to susceptible varieties on the first flush in the Spring and every additional flush. Application to fruit should start after two-thirds of the petals have fallen and be repeated at 21 day intervals.
	Phytophthora Foot Rot	2/3 pound	Mix this product with 1 gallon of water and paint trunks of trees from the soil surface to the lowest scaffold limbs. Apply in May prior to Summer rains and/or in the Fall prior to wrapping trees for freeze protection. Treatment serves as protection for up to one year, but does not cure existing infections.
	Phytophthora Brown Rot, Septoria Spot	2-2/3 to 5-1/3 pounds	Apply to the entire tree in the Fall before or just after the first rain and continue as needed.
	<p>NOTES:</p> <ul style="list-style-type: none"> • Do not use this product on citrus seedlings grown in greenhouses or shadehouses. • In California, in areas subject to copper injury, add 1/3 to 1 pound of high quality lime per pound of this product. 		
CITRUS FIELD NURSERY GROWN	Melanose, Scab, Greasy Spot, Pink Pitting, Brown Rot and Citrus Canker (Suppression Only)	2-2/3 to 5-1/3 pounds	Apply 1-1/3 pounds of this product in 100 gallons of water at 28 day intervals.
FILBERT	Bacterial Blight	10-2/3 to 16 pounds	Apply as a postharvest spray. In seasons of heavy rainfall, apply another spray when three-fourths of the leaves have dropped. Add 1 pint of superior type oil per 100 gallons of water.
	Eastern Filbert Blight	10-2/3 to 16 pounds	Apply in sufficient water to obtain thorough coverage. Make initial application at budswell to budbreak. Additional sprays should be made on a 10 to 14 day interval depending on disease severity or when conditions are conducive for disease development. Add 1 pint of superior type oil per 100 gallons of water.
KIWIFRUIT	Blossom Blight (Bud Rot), Leaf Spot (Phomopsis), Erwinia herbicola, Pseudomonas syringae, Pseudomonas fluorescens	1-1/3 to 2 pounds	Make two to three applications during dormant season. Do not apply at time of or after leaf emergence.

10/17

TREE CROPS (continued)

CROP	DISEASE	RATE PER ACRE	USE INSTRUCTIONS
MACADAMIA (Except California)	Blossom Blight & Raceme Blight, Anthracnose	3 to 6 pounds	Apply, depending on disease pressure, in 50 to 300 gallons of water during peak raceme development and bloom periods. For aerial application apply 3 to 6 pounds per acre in 10 to 30 gallons of water.
OLIVE (California)	Peacock Spot, Olive Knot	5-1/3 to 8 pounds	Make first application before Winter rains fall. A second application in early Spring should be made if disease is severe.
PEACH & NECTARINE	Leaf Curl, Coryneum Blight (Shot Hole), Bacterial Canker, Bacterial Blast (<i>Pseudomonas</i>), Bacterial Blight (<i>Xanthomonas</i>)	5-1/3 to 10-2/3 pounds	Apply after leaf fall as a dormant application. Use the higher rate when rainfall is very heavy and disease pressure is high. May be used with an agricultural spray oil.
	Blossom Brown Rot, Leaf Curl, Coryneum Blight (Shot Hole)	5-1/3 to 8 pounds	Apply as a full cover spray at pink bud. Application at this time affords some control of <i>Leaf Curl</i> and <i>Coryneum Blight</i> .
	Bacterial Spot	5-1/3 pounds	Apply as a dormant spray. (Except California) As a post-bloom spray: apply up to 1/2 pound per acre at first and second cover sprays.
	NOTE: Do not apply three weeks prior to harvest. Use only recommended rates. Spotting of leaves and defoliation may occur from use in cover sprays.		
PEAR	Fireblight	2/3 pound	Apply at 5 day intervals throughout bloom period.
	<i>Pseudomonas</i> Blight	8 to 10-2/3 pounds	Apply before Fall rains and again at dormant before Spring growth starts.
	NOTE: Excessive dosages may cause fruit russet.		
PECAN	Shuck Rot, Kernel Rot (<i>Phytophthora cactorum</i>), Zonate Leaf Spot (<i>Cristulariella pyramidalis</i>) (Suppression Only)	1-1/3 to 2-2/3 pounds	Apply at 2 to 4 week intervals when kernel growth begins through shuck opening. Apply in sufficient water to ensure thorough coverage.
PISTACHIO	Botrytis Blight, Botryosphaeria Panicle and Shoot Blight, Septoria Leaf Blight, Late Blight (<i>Alternaria alternata</i>)	2-2/3 to 5-1/3 pounds	Apply beginning at budswell. Repeat at 14 to 28 day intervals depending on disease conditions. If disease conditions are severe, use the high rate and the short spray interval.
QUINCE	Fire Blight	2/3 pound	Apply at 5 day intervals throughout the bloom period. Apply in sufficient water to provide thorough coverage.
WALNUT	Walnut Blight	5-1/3 to 8 pounds	Apply first spray at early pre-bloom prior to or when catkins are partially expanded. Make additional applications during bloom and early nutlet stage as needed. Additional applications may be necessary when frequent rainfall occurs.
	NOTE: When applied as a dilute spray, 1 pint of Summer oil emulsion may be added per 100 gallons of spray. Adequate control may not be obtained when copper tolerant species of <i>Xanthomonas</i> bacteria are present.		

11/17

TROPICAL CROPS

CROP	DISEASE	RATE PER ACRE	USE INSTRUCTIONS
BANANA	Sigatoka	1-1/3 pounds	Apply by air. Mix this product in 3 gallons of water containing 1/2 gallon agricultural oil. Apply on a 14 day schedule throughout the wet season. Apply at 21 day intervals during dry periods.
	Black Pitting	2-2/3 pounds per 100 gallons	Apply directly to the fruit stem and include the basal portion of the leaf crown. Apply during the first and second weeks after fruit emergence.
CACAO	Black Pod	1-1/3 to 5-2/3 pounds	Begin applications at the start of the rainy season and continue while infection conditions persist. Sprays should be made as often as 14 to 21 days in high rainfall areas at varying rates from 1-1/3 to 5-1/3 pounds per acre depending on disease severity. For drier areas, where two to four applications are recommended during critical infection periods and at long intervals, use 4 to 5-2/3 pounds per acre, according to disease pressure incidence and planting density.
COFFEE	Coffee Berry Disease (<i>Collectotrichum coffeanum</i>)	4 to 5-1/3 pounds	Apply first spray after flowering and before onset of long rains and then at 21 to 28 day interval until picking. Use higher rates when rainfall is heavy and disease pressure is high.
	Bacterial Blight (<i>Pseudomonas syringae</i>)	4 to 5-1/3 pounds	Begin spray program before onset of the long rains and continue throughout the rainy season at 14 to 21 day intervals. The critical time of spraying to control this disease is just before, during, and after flowering(s), especially when coinciding with wet weather. Use the higher rates when rainfall is heavy and disease pressure is high.
	Leaf Rust (<i>Hemileia vastatrix</i>)	1-1/3 to 2-2/3 pounds	Apply before the onset of rain and then at 21 day intervals while the rains continue. Use higher rates when rainfall is heavy and disease pressure is high.
	Iron Spot (<i>Cercospora coffeicola</i>) and Pink Disease (<i>Corticium salmonicolor</i>)	1-1/3 pounds	Apply as a concentrate or dilute spray. Begin treatment at the start of wet season and continue at monthly intervals for three applications.
GUAVA	Anthracnose, Red Algae	2 pounds	Apply beginning just prior to flowering and repeat weekly until just prior to harvest.
LITCHI	Anthracnose	2 pounds	Apply beginning just prior to flowering and repeat weekly until just prior to harvest.
MAMEY SAPOTE	Anthracnose, Algal Leaf Spot	4 to 5 pounds	Apply when conditions favor disease development. Repeat at 14 to 30 day intervals as needed.
MANGO (Florida & Puerto Rico)	Anthracnose	5-1/3 to 6-2/3 pounds	Apply monthly after fruit set until harvest.
PAPAYA	Anthracnose	2 to 6-2/3 pounds	Apply beginning before disease is expected to appear. Repeat at 10 to 14 day intervals or at 5 to 7 day intervals during periods of heavy rainfall. Use the higher rates when conditions favor disease. The addition of a suitable spreader-sticker may be desirable especially during periods of heavy rains.
PASSION FRUIT	Anthracnose	4 pounds	Apply beginning just prior to flowering and repeat weekly.
SUGAR APPLE (Annona)	Anthracnose	8 pounds	Apply beginning just prior to flowering and repeat weekly.

12/17

VEGETABLE CROPS

CROP	DISEASE	RATE PER ACRE	USE INSTRUCTIONS
BEAN (Dry, Green)	Brown Spot, Bacterial Blight (Halo & Common)	2/3 to 2 pounds	For protective sprays, apply first application when plants are six inches high. Apply on 7 to 14 day schedule depending on local conditions. Adjust rates depending on disease severity.
CARROT	Alternaria Leaf Spot, Carrot Blight (Cercospora)	1-1/3 pounds	When disease threatens apply at 7 to 14 day intervals depending on disease severity.
CELERY & CELERIAC	Cercospora Early, Septoria Late Blight & Bacterial Blights	1-1/3 pounds	Apply as soon as plants are first established in the field, then every 5 to 7 days depending on disease severity and weather.
CRUCIFERS Broccoli, Brussels Sprout, Cabbage, Cauliflower, Kale, Collard Greens, Mustard Greens, and Turnip Greens	Black Rot (<i>Xanthomonas</i>), Black Leaf Spot (<i>Alternaria</i>), Downy Mildew	1/3 to 2/3 pound	Apply at 7 day intervals after transplants are set in the field. Use higher rate when conditions favor disease.
	NOTE: Reddening of older leaves may occur on Broccoli at the higher rate and flecking of wrapper leaves may occur on Cabbage.		
CUCURBITS Cantaloupe, Cucumber, Honeydew, Muskmelon, Pumpkin, Squash, and Watermelon	Alternaria Leaf Spot, Angular Leaf Spot, Anthracnose, Downy Mildew, Powdery Mildew, Gummy Stem Blight, Watermelon Bacterial Fruit Blotch (Suppression)	1-1/3 pounds	Apply beginning when conditions are favorable for disease development and repeat at 5 to 7 day intervals, as needed depending on disease severity.
	NOTE: Crop injury may occur from application at shorter intervals. Discontinue use if injury occurs.		
EGGPLANT (Except California)	Alternaria Blight, Anthracnose, Phomopsis	1-1/3 pounds	Use before disease appears. Repeat at 7 to 10 day intervals.
ENDIVE, ESCAROLE	Downy Mildew	2/3 to 1-1/3 pounds	Begin treatment when disease first appears and repeat every 7 to 10 days as needed to suppress disease.
GARLIC, LEEK, ONION	Purple Blotch & Downy Mildew	1-1/3 pounds	Apply when plants are four to six inches high and repeat at 7 to 10 day intervals.
	Bacterial Blight	2/3 to 1 pound	
LETTUCE	Downy Mildew	2/3 to 1-1/3 pounds	Begin treatment when disease first appears and repeat every 7 to 10 days as needed to suppress disease.
PEA	Powdery Mildew	1 to 2 pounds	Begin spray treatment when disease symptoms first appear. Adjust rates according to disease severity. Repeat applications at weekly intervals.
PEPPER	Bacterial Spot	1-1/3 to 2 pounds	When disease threatens, apply in sufficient water for adequate coverage at 5 to 10 day intervals depending on disease severity.
SPINACH	Anthracnose, Cercospora Leaf Spot, Downy Mildew, White Rust, Blue Mold	1-1/3 to 2-2/3 pounds	Begin treatment when disease first appears and repeat every 7 to 10 days as needed to suppress disease.
	NOTE: Flecking may occur on spinach leaves.		
TABLE BEET, BEET GREENS	Cercospora Leaf Spot	1-1/3 to 2-2/3 pounds	Apply when conditions favor disease. Repeat treatment at 10 to 14 day intervals as needed. The addition of an agricultural spray oil is recommended.

13/17

VEGETABLE CROPS (continued)

CROP	DISEASE	RATE PER ACRE	USE INSTRUCTIONS
TOMATO	Early Blight, Late Blight	1-1/3 to 2 pounds	When disease threatens, apply at 7 to 10 day intervals, or as necessary.
	Bacterial Speck	1-1/3 pounds	Apply at 10 to 30 day intervals beginning when the disease threatens. Use more frequent applications when disease pressure is high.
	Bacterial Spot, Anthracnose, Gray Leaf Mold & Septoria Leaf Spot	1-1/3 to 2-2/3 pounds	When disease threatens, apply at 7 to 10 day intervals, more frequently when disease is severe.
WATERCRESS	Cercospora Leaf Spot	1-1/3 pounds	Apply when plants are established in the field. Repeat at 7 to 14 day intervals up to four applications per crop in at least 50 gallons of water per acre.

MISCELLANEOUS

CROP	DISEASE	RATE PER ACRE	USE INSTRUCTIONS
ATEMOYA	Anthracnose	2 pounds	Apply just prior to flowering and repeat weekly until just prior to harvest.
CARAMBOLA	Anthracnose	4 pounds	Apply just prior to flowering and repeat weekly until just prior to harvest.
CHIVES	Downy Mildew	1-1/3 pounds	Apply when plants are established in the field. Repeat at 7 to 10 day intervals as needed.
DILL	Phoma Leaf Spot, Rhizoctonia Foliage Blight	1-3/4 pounds	Apply when plants are established in the field. Repeat at 7 to 10 day intervals as needed.
DOUGLAS FIR	Rhabodocline Needlecast	1-1/3 pounds	Apply at bud break and repeat at 3 to 4 week intervals. Apply in a tank mix with other registered pesticide if disease pressure is moderate to severe.
GINSENG	Alternaria Leaf & Stem Blight	1-3/4 pounds	This product may be applied as a tank mix with two pounds Iprodione 50WP in 100 gallons of water per acre. Begin Iprodione 50WP/Champ Dry Prill applications as soon as plants have emerged in Spring. Applications should be repeated every 7 days until plants become dormant in Fall. Apply fungicides at least eight hours before rain, giving the fungicides time to dry on the plants. Use of a spreader-sticker is advised. NOTE: <i>Alternaria Leaf & Stem Blight</i> is most severe in humid conditions such as those found in the dense canopies of two, three and four year old ginseng. Complete and thorough spray coverage is required for control.
PARSLEY	Bacterial Blight (<i>Pseudomonas spp.</i>)	2 pounds	Apply when plants are first established in the field and repeat at 5 to 7 day intervals.
PERSIMMON	Cercospora Leaf Spot	1-1/3 pounds	Apply beginning in May/June, during leaf flush, and repeat at 14 day intervals throughout the season depending on disease severity.

14/17

TURFGRASS

CROP	DISEASE	RATE	USE INSTRUCTIONS
TURFGRASS (such as sodfarms, golf courses, cemeteries, home lawns, and industrial or municipal turf areas (including parks, playgrounds, athletic fields))	Algae Control	1 pound per 1,000 square feet	Apply in 5 gallons of water to control algae. This product may be used alone or in combination with other registered fungicides as a maintenance spray.
	NOTE: Phytotoxicity may occur depending upon varietal differences. If injury occurs discontinue use. Do not apply in spray solutions with a pH of less than 6.5.		

GREENHOUSE AND SHADEHOUSE CROPS

This product may be used in greenhouses and shadehouses to control diseases on crops listed on this label. Specific directions are provided below for certain crops and the grower should be aware that the sensitivity of crops grown under such conditions differ greatly from field conditions. The user must determine if this product can be used safely prior to commercial application by testing a small area and observing the results for 7 to 10 days.

One level tablespoon of this product per 1,000 square feet is equivalent to 1 pound per acre. Begin application at first sign of disease and repeat at 7 to 14 day intervals as needed.

CROP	DISEASE	RATE	USE INSTRUCTIONS
CUCUMBER	Angular Leaf Spot, Downy Mildew	1-1/4 to 1-1/2 tablespoons	Apply weekly when plants begin to vine.
EGGPLANT	Alternaria Blight, Anthracnose, Phomopsis	1-1/2 tablespoons	Apply at first sign of disease and repeat at 7 to 14 day intervals as needed.
PEPPER	Bacterial Spot	1-1/2 to 2-1/4 tablespoons	Apply when conditions first favor disease and at 5 to 10 day intervals as needed.
TOMATO	Anthracnose, Bacterial Speck, Bacterial Spot, Early Blight, Grey Leaf Mold, Late Blight, Septoria Leaf Spot	1-1/2 to 2-1/4 tablespoons	Apply when conditions first favor disease and at 7 to 10 day intervals as needed.

NOTE: Do not use this product on Citrus seedlings grown in greenhouses or shadehouses.

ORNAMENTALS

CROP	DISEASE	RATE	USE INSTRUCTIONS
PECAN, LIVE OAK (Texas and Florida)	Ball Moss	5-1/2 pounds per 100 gallons of water	Apply in Spring after heavy rain, using 1-1/2 gallons of spray per foot of tree height. Make sure to wet tufts thoroughly. A second application may be required after 12 months.
	NOTE: This product may be injurious to ornamentals grown under live oaks.		
PHILODENDRON	Bacterial Leaf Spot	1 pound per 100 gallons of water	Apply weekly before disease appears.
SYCAMORE	Anthracnose	1-1/3 to 2 pounds per 100 gallons of water	Make two applications as a full cover spray. Make first application at bud crack and second application 7 to 14 days later at 10% leaf expansion.

15/17

For control of bacterial and fungal diseases on foliage, flowers, and stems of ornamentals grown in **Greenhouses, Shadehouses, Fields and Nurseries** (container, bench or bed-grown): apply this product at 2/3 pound per 100 gallons as a full cover spray beginning at first sign of disease. Repeat at intervals of 7 to 14 days (or shorter) depending on rainfall and disease severity. Due to the large number of species, widely varying growth conditions, and varieties of ornamentals and nursery plants it is impossible to test every one for sensitivity to this product. The user should apply the recommended rate of this product in a small area and check for any symptoms of phytotoxicity in 7 to 10 days prior to large-scale application.

Do not tank mix with Aliette® fungicide without buffering the spray solution.

One-half tablespoon of this product per gallon of water is equivalent to 0.9 pound per 100 gallons.

ORNAMENTAL	DISEASE
AGLAONEMA	Bacterial Leaf Spot
ALTHEA (Rose of Sharon)	Bacterial Leaf Spot
ARALIA	Xanthomonas & Cercospora Leaf Spots, Alternaria
ARBORVITAE	Alternaria Twig Blight, Cercospora Leaf Spot
AZALEA (1)	Cercospora Leaf Spot, Botrytis Blight, Phytophthora Dieback, Powdery Mildew
BEGONIA	Bacterial Leaf Spot (<i>Xanthomonas spp.</i> , <i>Erwinia spp.</i> , <i>Pseudomonas spp.</i>)
BOSTON FERN	Bacterial Leaf Spot
BOUGAINVILLEA	Anthracnose, Bacterial Leaf Spot
BULBS (Easter Lilly (2), Tulip)	Botrytis Blight, Anthracnose
CAMELLIA	Anthracnose, Bacterial Leaf Spot
CAMPHOR TREE	Pseudomonas Leaf Spot
CANNA	Pseudomonas Leaf Spot
CARNATION (1)	Alternaria Blight, Pseudomonas Leaf Spot & Botrytis Blight
CHINESE TALLOW TREE	Bacterial Leaf Spot (<i>Xanthomonas spp.</i> , <i>Pseudomonas spp.</i>)
CHRYSANTHEMUM (1)	Septoria Leaf Spot, Botrytis Blight
COTONEASTER	Botrytis Blight
DAHLIA	Alternaria Leaf Spot, Cercospora Leaf Spot, Botrytis Grey Mold
DATE PALM	Pestalotia Leaf Spots
DIANTHUS	Bacterial Spot, Bacterial Soft Rot
DOGWOOD	Anthracnose
DRACAENA	Bacterial Leaf Spot
DUMB CANE	Bacterial Leaf Spot
DUSTY MILLER	Bacterial Leaf Spot (<i>Pseudomonas cichorii spp.</i>)
ECHINACEA	Botrytis Blight
ELM (Drake)	Xanthomonas Leaf Spot
EUONYMUS	Botrytis Blight, Anthracnose
EUROPEAN FAN PALM	Pestalotia Leaf Spot
GARDENIA	Alternaria Leaf Spot, Cercospora Leaf Spot, Botrytis Bud Rot
GERANIUM	Alternaria Leaf Spot, Cercospora Leaf Spot, Botrytis Grey Mold
GLADIOLUS	Alternaria Leaf Spot, Botrytis Grey Mold, Bacterial Leaf Blight, Botrytis Blight, Anthracnose
GOLDEN RAIN TREE	Alternaria Leaf Spot, Botrytis Grey Mold, Bacterial Leaf Blight
GRAPE IVY	Bacterial Leaf Spot
HIBISCUS (3)	Bacterial Leaf Spot
HOLLY FERN	Pseudomonas Leaf Spot
HONEY LOCUST	Bacterial Leaf Spot
IMPATIENS	Bacterial Leaf Spot
INDIA HAWTHORN (4)	Anthracnose, Entomosporium Leaf Spot

16/17

ORNAMENTAL	DISEASE
IRIS	Bacterial Leaf Spot
IVY (English, Algerian) (1)	Xanthomonas Leaf Spots
IXORA	Xanthomonas Leaf Spots
JUNIPER (Eastern Red Cedar)	Anthracnose
LANTANA	Bacterial Leaf Spot
LILAC	Cercospora Leaf Spot
LOBLOLLY BAY	Anthracnose
LOQUAT	<i>Entomosporium maculata</i> , <i>Colletotrichum spp.</i>
MAGNOLIA (Southern)	Anthracnose, Bacterial Leaf Spot, Algal Leaf Spot
MAGNOLIA (Sweet Bay)	Anthracnose
MAGNOLIA	Bacterial Leaf Spot
MANDEVILLAS	Anthracnose
MARIGOLD	Alternaria Leaf Spot, Botrytis Leaf Rot, Flower Rot, Cercospora Leaf Spot
MULBERRY - CONTORTED	Bacterial Leaf Spot
MULBERRY - WEEPING	Bacterial Leaf Spot
NEPHTHYTIS	Bacterial Leaf Spot
OLEANDER	Bacterial Leaf Spot, Fungal Leaf Spot
OAK, LAUREL	Algal Leaf Spot, (<i>Cephaleuros virescens spp.</i>)
PACHYSANDRA	Volutella Leaf Blight
PANSY	Downy Mildew
PARLOR PALM	Bacterial Leaf Spot
PEAR (flowering)	Fire Blight, Leaf Spot
PENTAS (Egyptian Star)	Bacterial Leaf Spot (<i>Xanthomonas spp.</i>)
PEONY	Botrytis Blight
PERIWINKLE	Phomopsis Stem Blight
PHLOX	Alternaria Leaf Spot
PHOTINIA (Red Tip, Red Leaf)	Anthracnose, Entomosporium
PISTACHIO	Anthracnose
PLANTAIN LILY	Bacterial Leaf Spot
POTHOS	Bacterial Leaf Spot
POWDER PUFF PLANT	Bacterial Leaf Spot
PURPLE OSIER WILLOW	Anthracnose
PYRACANTHA	Fireblight, Scab
QUEEN PALM	Exosporium Leaf Spot, Phytophthora Bud Rot
RHODODENDRON	Alternaria Flower Spot
ROSE (1)	Powdery Mildew, Black Spots
SNAPDRAGON	Anthracnose, Dieback, Downy Mildew
SPATHE FLOWER	Bacterial Leaf Spot
TATARIAN HONEYSUCKLE	Bacterial Leaf Spot
UMBRELLA TREE	Bacterial Leaf Spot
VERBENA	Xanthomonas Leaf Spot

17/17

ORNAMENTAL	DISEASE
VIBURNUM	Anthracnose
WASHINGTON PALM	Pestalotia Leaf Spot
WEeping FIG	Bacterial Leaf Spot
WEeping WILLOW	Anthracnose
YUCCA (ADAMS NEEDLE)	Cercospora & Septoria Leaf Spots

- (1) On some varieties a discoloration may occur on foliage or blooms. To prevent residues on commercial plants, do not spray just before selling season.
- (2) Apply 2 to 3-1/2 pounds of this product in 20 to 100 gallons of water per acre
- (3) Hibiscus – Do not apply to plants in flower.
- (4) For India Hawthorn use 1-2/5 to 2-3/4 pounds per 100 gallons or 3/4 to 1-1/2 tablespoons per gallon.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool dry place.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest Environmental Protection Agency Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. **THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE.** UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV051007A)

Champ® is a registered trademark of Nufarm Americas, Inc.
 Aliette® is a registered trademark of Bayer.
 © 2007 Nufarm Americas, Inc.