

55146-66

176

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number
222104

Application for Pesticide - Section I

1. Company/Product Number 55146-66	2. EPA Product Manager Theresa A. Stowe	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) GibGro 10% Powder	PM# 22	
5. Name and Address of Applicant (Include ZIP Code) AGIROL CHEMICAL PRODUCTS 7322 Southwest Freeway, Suite 1400 Houston, Texas 77074 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)
Response to PR Notice 95-3.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	Other (Specify) _____	
		If "Yes" Package wgt	No. per container		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Linda C. Watson	Title Consultant/Regulatory Affairs	Telephone No. (Include Area Code) (912) 482-2771
2. Signature 		6. Date Application Received (Stamped)
3. Title Consultant/Regulatory Affairs		
4. Typed Name Linda C. Watson	5. Date October 10, 1995	

GibGro 10% POWDER

(GIBBERELIC ACID)

ACTIVE INGREDIENT:

Gibberellic Acid 10.0%

INERT INGREDIENT: 90.0%

TOTAL 100.0%

Contains a total of 16 g. of Gibberellic Acid

Net Weight - 160 grams (5.64 ounces)

EPA Reg. No. 55146-66

EPA Est. No. 65663-TX-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye injury. Harmful if inhaled or absorbed through skin. Avoid breathing dust or spray mist. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear: Long-sleeved shirt and long pants, waterproof gloves and shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear: Coveralls, waterproof gloves and shoes plus socks.

IMPORTANT

Before application, read accompanying GibGro 10% Powder Spray Guide carefully and use only as directed. **DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.**

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep containers tightly closed when not in use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY STATEMENT

AGTROL CHEMICAL PRODUCTS warrants that the product conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions. **THERE ARE NO OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING A WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE.** This warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller and buyer assumes all risk of any such use.

GibGro 10% Powder

(Gibberellic Acid)

SPRAY GUIDE

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.
IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.
IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye injury. Harmful if inhaled or absorbed through skin. Avoid breathing dust or spray mist. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear: Long-sleeved shirt and long pants, waterproof gloves and shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear: Coveralls, waterproof gloves and shoes plus socks.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep containers tightly closed when not in use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTE: Gibberellic Acid is an extremely potent plant growth regulator. For best results, read all directions for use thoroughly. Consult your local experiment station specialist, distributor, or the Agtrol agricultural specialist in your area for the spray schedule best suited to your conditions.

Discard any unused spray material at the end of each day. Prepare solution concentrations by mixing the required amount of product with water only in a clean, empty spray tank.

Use only as directed. The label should be read thoroughly and understood before making applications. Effectiveness requires that all parts of plant or crop must receive spray or desired result will not occur, so spray thoroughly. When a range of rates is indicated, use the concentration and spray volume recommended locally.

Data concerning the compatibility of GibGro with other agricultural compounds is not available, except where specified.

SPRAY GUIDELINES FOR GRAPES

For all grapes, application is recommended by ground sprayer. Use 100 to 500 gallons per acre as a dilute spray according to foliage density, or 30 to 80 gallons per acre as a concentrate spray, unless specified otherwise. Do not exceed maximum rates. It is important to wet all berries thoroughly.

THOMPSON SEEDLESS GRAPES

For cluster elongation ("Stretch"), looser cluster forms, and reducing cost of thinning, when used in conjunction with established girdling and thinning practices: apply 8 to 16 grams a.i. per acre before bloom when flower clusters are 3 to 5 inches long.

For decreased berry set ("Thinning"), reducing hand-thinning costs, and hastened maturity: apply 8 to 16 grams a.i. per acre per application during bloom as one application or as two applications of equal amounts when the bloom period is extended with the second application made 3 to 7 days after the first application.

For larger berries ("Sizing") and larger clusters when used in conjunction with established girdling and thinning practices: apply 32 to 80 grams a.i. per acre per application in 1 to 3 applications beginning when average berry size is 4-5 millimeters in diameter. Applications should be applied within a 14 day period. Timing of the second and third spray will be dictated by experience in the vineyard to be sprayed and temperatures occurring during the interim between sprays. Potential effect will be reduced if the second and/or third spray occurs more than two weeks after the first application.

NOTE: Do not apply more than 208 grams a.i. per acre per growing season for all uses.

THOMPSON SEEDLESS GRAPES FOR RAISINS

For cluster elongation ("Stretch") and looser cluster forms, allowing better air circulation to aid in the control of bunch rot and increase light penetration aiding in sugar development: Apply 8 to 16 grams a.i. per acre before bloom when flower clusters are 3 to 5 inches long.

For decreasing berry set, ("Thinning") with increased raisin quality, and hastened maturity: apply 0.75 to 6 grams a.i. per acre when most bunches are in 60% to 80% bloom.

FLAME SEEDLESS GRAPES

For decreased berry set ("Thinning") and reducing hand-thinning costs: apply 3 to 7.5 grams a.i. per acre during bloom. Higher amounts may cause an excess of shot berries or overthinning.

For larger berries ("Sizing") and larger clusters when used in conjunction with established girdling and thinning practices: apply 20 to 48 grams a.i. per acre per application in 1 to 3 applications beginning when average berry size reaches 6 to 8 millimeters in diameter. Applications should be applied within a 14 day period. Timing of the second and third spray will be dictated by experience in the vineyard to be sprayed and temperatures occurring during the interim between sprays. Potential effect will be reduced if the second and/or third spray occurs more than two weeks after the first application.

NOTE: Do not apply more than 103.5 grams a.i. per acre per growing season for all uses.

PERLETTE GRAPES

For larger berries ("Sizing") and larger clusters when used in conjunction with established girdling and thinning practices: Apply 32 to 80 grams a.i. per acre per application in 1 to 3 applications beginning when average berry size is 4 to 5 millimeters in diameter. Applications should be applied within a 14 day period. Timing of the second and third spray will be dictated by experience in the vineyard to be sprayed and temperatures occurring during the interim between sprays. Potential effect will be reduced if the second and/or third spray occurs more than two weeks after the first application.

NOTE: Do not apply more than 160 grams a.i. per acre per growing season for all uses.

OTHER SEEDLESS VARIETIES SUCH AS SEEDLESS TOKAY, INTER-LOCKEN SERIES AND RELATED HYBRIDS

For larger berries and larger clusters when used in conjunction with established girdling and thinning practices: apply 8 to 48 grams a.i. per acre as one application at or just after shatter (usually 2 to 3 days later) or as two applications of equal amounts not to exceed a total of 48 grams a.i. per acre, with the first made at or just after shatter, followed during the next two weeks

by the second application. Timing of the second spray with split application will be dictated by experience in the vineyard to be sprayed and temperatures occurring during the interim between sprays. Potential effect will be reduced if the second spray occurs more than two weeks after the first application.

EMPEROR GRAPES

For reducing berry shrivel. This can also increase berry size: apply 20 grams a.i. per acre as one application in 200 to 250 gallons/A approximately two weeks after completion of shatter following bloom. This timing should correspond to a period when the predominant berry diameter ranges from 10 to 15 millimeters.

BLACK CORINTH (ZANTE CURRANT) GRAPES

For improving berry size: apply spray containing 1 to 8 grams a.i. per acre 3 to 5 days after full bloom, but before shatter begins.

SPRAY GUIDELINES FOR CITRUS

NAVEL ORANGES

(California) To delay aging of the rind and reduce rind disorders (e.g. rind staining, water spotting, sticky or tacky surface, puffy rind and rupture under pressure) and to produce a more orderly harvesting pattern: **EARLY SPRAY** (before color change). The delay in rind aging is greatest when the early spray is applied before a color change. This spray timing produces the firmest rind possible. Apply one spray two weeks prior to color break which normally occurs August to November. Apply 10 to 40 grams a.i. per acre as a concentrate or dilute spray in sufficient gallonage to insure thorough wetting.

NOTE: Do not apply to groves that may be harvested early as a reduction in grade may result due to the delayed coloring. Do not apply in white wash sprays in which lime or other caustic material has produced a high pH in the spray tank.

LATE SPRAY: (after color break) Apply one spray just after marketable color has developed which is normally from October through December. Apply 16 to 48 grams a.i. per acre as a concentrate or dilute spray in sufficient gallonage to insure thorough wetting.

NOTE: Do not spray Navel orange trees from January through July. Sprays applied in January/February may cause reduced production the following year. Do not apply within 10 days of harvest.

NOTE: A slight increase in mature leaf drop may occur in trees under stress.

VALENCIA ORANGES

(California) To reduce rind creasing and to delay aging and softening of the rind: apply a single spray in August or September to trees with a target crop of young fruit. Apply 40 to 80 grams a.i. per acre as a concentrate or dilute spray in sufficient gallonage to insure thorough wetting.

NOTE: Slower color development should be expected in the target crop. Increased regreening of mature fruit, if present, may occur. After marketable color is achieved, treatment effects may be reduced the longer treated fruit remain on the tree.

LEMONS

(California except desert valleys) To decrease the amount of small tree ripe fruit and to produce a more desirable production pattern in relation to market demand: apply one spray when target crop is 1/2 to 3/4 full size but still green. Use 10-20 grams a.i. per acre as a concentrate or dilute spray in sufficient gallonage to insure thorough wetting.

When applied two years in a row, an even larger difference in harvest pattern and maturity occurs.

NOTE: Do not apply within one month of harvest. Do not apply in spring or summer.

TANGERINE HYBRIDS

(Florida) To increase fruit set and yields on tangerine hybrids with pollination problems such as the Orlando, Robinson, Minneola and Sunburst: apply spray during full bloom. Be sure to wet the leaves sufficiently.

Fruits are generally seedless. Use 8 to 30 grams a.i. in 400 to 500 gallons/A on large mature trees.

NOTE: A slight increase in mature leaf drop occurs at concentrations above 25 ppm. Fruit sizes may be reduced and color development slightly retarded.

(California) To delay disorders associated with rind aging of the Minneola tangelo; e.g., puffiness and softening and to increase peel strength: apply 20 to 40 grams a.i. per acre as a dilute spray in sufficient gallonage to insure thorough wetting.

NOTE: Do not apply if early harvest is planned. Do not apply after coloring as pre-harvest rind staining may occur. Application during coloring may cause variation in rind color development.

GRAPEFRUIT

(Florida and Texas) To delay disorders associated with rind aging; e.g., puffiness, softening and orange coloration, to prevent preharvest drop of mature fruit and to increase peel strength and reduce water loss during storage: apply a single spray to fully colored fruit during the November through January period. Use 20 to 56 grams a.i. in 500 to 700 gallons/A containing a suitable non-ionic surfactant at the manufacturer's recommended rate. It is advisable to spot pick heavy crops to aid early marketing and to avoid reduction of yields which generally follow late held crops.

NOTE: Applications made after January or when trees begin to break dormancy may adversely affect new crop. Do not use concentrate sprays. Results may vary season to season depending on environmental conditions.

GRAPEFRUIT, STAR RUBY VARIETY

(Texas) To reduce early-season drop of small fruit of Star Ruby Variety thereby increasing yields: apply a single spray during the bloom period. Use 25 grams a.i. in 250 gallons water final spray mixture per acre. A suitable surfactant may be used to enhance efficacy.

NOTE: Do not tank-mix with other chemicals. Do not apply concentrated

solution. Results may vary season to season depending on environmental conditions. Maintain a well-balanced fertilization and watering program.

SPRAY GUIDELINES FOR FRUIT CROPS

BLUEBERRIES

For improving fruit set. For set problems due to insufficient natural honeybee pollination on varieties such as Coville, Jersey, Stanley, Earlieblue, Weymouth and others: make a single foliage spray application at full bloom (when over 75 percent of all flowers are fully open). For Weymouth, application can be delayed up to two weeks after full bloom to affect sizing of shot berries. Use 80 grams a.i. in 100 gallons of water. Use of a spreader-sticker is recommended. Apply to the point of run-off, thoroughly wetting all parts of the plant. Total gallonage will depend on size and density of the plants.

NOTE: Do not exceed 300 gallons/A. Although some varieties bloom closer to harvest than others—in no case should application be made closer than 40 days before harvest. Do not apply to plants in a low state of vigor.

SWEET CHERRIES

To delay harvesting, to produce a brighter colored, firmer fruit and to increase size; apply spray when the fruit is light green to straw colored. Apply 16 to 48 grams a.i. per acre using sufficient water to obtain complete coverage of the tree.

NOTE: Do not apply within one week of harvest.

RED TART CHERRIES

(All states except California) To maintain and extend high fruiting capacity of bearing tart cherry trees and reduce occurrence of "blind" nodes by stimulating lateral vegetative buds to develop a more productive balance of lateral shoots and spurs. GibGro must be applied annually to insure vegetative development and subsequent yield improvement year after year.

Timing: Apply a single foliar spray between 14 to 28 days after bloom. Research and commercial experience has determined 21 days after full bloom to be optimum. Best timing is further defined as that stage when 3-5 terminal leaves have fully expanded, or, at least 1-3 inches of terminal shoot extension has occurred.

Concentration: 10 to 25 ppm. The most commonly used rate is 15 ppm. However, higher or lower rates may be used, depending upon the response you desire.

Method of Application: Best results have been achieved with high volume sprays of 100 gallons or more of finished spray per acre. However, lower volume sprays can be equally effective, but extreme care must be exercised to avoid an overdose as spray volume is decreased.

HIGH VOLUME SPRAY GUIDE (100 or more gallons per acre)

TREE AGE	6-10 YRS.	10-15 YRS.	16-20 YRS.	20+ YRS.
CONCENTRATION (PPM)	10 PPM	15 PPM	20 PPM	25 PPM
GRAMS ACTIVE INGREDIENT PER 100 GALLONS	4 GRAMS	6 GRAMS	8 GRAMS	10 GRAMS
RECOMMENDED WATER VOLUME (GALLONS/ACRE)	150	150	150	150

GRAMS ACTIVE INGREDIENT PER ACRE	6 GRAMS	9 GRAMS	12 GRAMS	15 GRAMS
----------------------------------	---------	---------	----------	----------

LOW VOLUME SPRAY GUIDE (50-100 gallons per acre)

APPROXIMATE TREE AGE	GRAMS OF ACTIVE INGREDIENT PER ACRE	
	NORMAL VIGOR	LOW VIGOR
6-10 YEARS	4	6
10-15 YEARS	8	10
15-20 YEARS	10	14
20 + YEARS	14	18

NOTE: Use a minimum of 50 gallons/acre for a low volume spray application and obtain uniform coverage of the whole tree. Rates of GibGro in the above chart are based on expected tree vigor at various ages in a normal orchard. Each orchard presents a different situation. Adjust GibGro rate to complement vigor of trees. If trees are vigorous, use lowest recommended rates. Use higher rate for trees low in vigor and weak in shoot and spur production. Excessive application rates on any tree will increase vegetative growth at the expense of fruit production the following year.

NOTE: Lowest rates of GibGro should be used on trees that have been heavily pruned or hedged. The use of additional wetting or spreading agents is not recommended.

GibGro will not improve growth of trees under stress (nutritional, moisture, winter injury) or other factors inhibiting normal growth and development resulting from physical damage or unsound orchard practices. Best results from GibGro will be obtained when combined with good cultural practices.

SPRAY GUIDELINES FOR NON-BEARING FRUIT TREES

YOUNG TART AND SWEET CHERRY TREES

(All states except California) To reduce flowering and fruiting in young tart and sweet cherry trees to minimize the competitive effect of early fruiting on tree development: apply GibGro two to four weeks after bloom. Use 20 to 40 grams a.i. in 100 gallons of water. Apply a foliar spray of 25 to 50 gallons per acre, assuming a tree density of 100 trees per acre equivalent, or apply about one quart of spray volume per tree. Under conditions of low vigor, two applications are recommended. If two spray applications are made, allow at least a seven-day interval between sprays.

NOTE: DO NOT SPRAY TREES IN THE FIRST YEAR. Treat in the second season for reduction of flowering in the third season and again in the third season if reduction of flowering and fruiting is desired in the fourth season.

NON-BEARING PEACHES

(North Carolina, South Carolina, Georgia, Florida, Alabama, Tennessee, Mississippi) *To reduce flowering and fruiting in young non-bearing peaches to minimize the competitive effect of early fruiting on tree development:* Apply a single spray in the fall after flower buds have been initiated. This corresponds to the period immediately before and at the onset of early leaf drop, typically late September to early October. Apply GibGro at the rate of 200-400 PPM in 10-50 gallons of water per acre. Best results are obtained when applied with a handgun and tree canopy is wetted thoroughly to the point of run-off. The addition of a non-ionic surfactant will improve efficacy. Refer to the table for mixing instructions.

	200 PPM	400 PPM
GRAMS OF GIBGRO 10% POWDER/LL:		
10 Gallons of Water	80	160
50 Gallons of Water	400	800

Treat only trees that are in good physiological condition. Trees should have completed their first leaf before commencing treatments. Discontinue treatment the year before desired harvest.

SPRAY GUIDELINES FOR OTHER FRUIT

OLYMPUS STRAWBERRIES

(N.W. US Only; propagation stock) *To increase runner production of mother plants of the Olympus cultivar:* apply a single spray to mother plants 10 to 30 days after planting. At the time of spraying, plants should have 1 to 6 leaves. Apply 100 gallons/A to thoroughly wet new foliage to the point of run-off. Use 20 grams a.i. per acre.

NOTE: Not for use on fruiting plants. Treatments may not be effective on plantings set out after mid-May.

FORCING RHUBARB

To increase yield of marketable forced rhubarb and to break dormancy on plants receiving insufficient chilling: apply 2 fluid ounces (60 ml) of a solution containing 20 grams a.i. in 10 gallons to each cleaned crown, when the rest period is not completely broken. When the rest period is broken by cold weather, apply 2 fluid ounces (60 ml) of a solution containing 10 grams a.i. in 10 gallons.

NOTE: Keep forcing house temperatures at 40° to 50°F for 24 hours after application. If house is warmer than 50°F, the crowns should be covered with plastic. Temperatures in the forcing house above 50°F will result in lower yields and poor stalk color.

SPRAY GUIDELINES FOR VEGETABLE CROPS

ARTICHOKES

(California) *To accelerate maturity of artichokes and to shift the harvest to an earlier date:* apply spray at bud initiation time, normally six weeks prior to anticipated harvest. Be sure the entire plant (leaves, stems and buds) are covered to point of run-off. Use 10 grams a.i. in 100 to 125 gallons/A.

NOTE: Do not apply within seven days of harvest.

CELERY

To increase plant height and yield, and overcome stress due to cold weather conditions, or saline soils and to obtain earlier maturity: apply spray one to four weeks prior to harvest. Lower concentrations are applied at the three to four-week interval. Higher concentrations at the one to two-week interval. Use 2.5 to 10 grams a.i. in 25 to 50 gallons/A.

NOTE: Do not apply earlier than four weeks before harvest as Gibberellic Acid may induce bolting (seed stalk formation).

Applications made less than one week preharvest may result in residues.

Celery plants must be harvested when mature to ensure quality.

LETTUCE FOR SEED

To obtain uniform bolting and increase seed production: apply the following spray schedule:

Growth Stage	ppm	g a.i./A	Gallon/Acre
4 leaf stage	10	0.4	10
8 leaf stage	10	1.6	40
12 leaf stage	10	4	100

NOTE: Do not feed crop wastes to livestock.

MELONS AND CUCUMBERS (Except California)

To stimulate fruit set during periods of extended cool temperatures, apply 2 grams a.i. per acre. Make one application prior to periods of extended cool temperatures. Make one application prior to bloom and two additional applications at 10-14 day intervals following fruit set on cantaloupes and watermelons. For cucumbers, as many as 3 to 4 applications may be required after fruit set.

An adequate spray volume should be used to insure thorough coverage of the exposed foliage. In order to obtain maximum benefit from GibGro, the vines must be in good condition except for a reduced growth rate due to cool temperatures.

SEED POTATOES

To stimulate uniform sprouting—for maximum production, more uniform development, fewer late maturing plants, and to break dormancy of newly harvested potatoes that have not had a full rest period: Dip freshly dug seed pieces in a solution containing 0.2 to 0.4 gram a.i. in 100 gallons prior to planting.

NOTE: If soil temperature is very high, avoid treating rested seed and use the minimum concentration for dormant seed.

SPINACH

(All States except California) *To facilitate harvest, increase yield and improve quality of fall and over-winter spinach:* apply a single spray 10 to 14 days before each anticipated harvest on fall or over-winter spinach, ideally

when daytime temperatures are 40° to 70°F and during early morning hours when dew is present on crop.

Use 6 to 8 grams a.i. per acre in 10 to 50 gallons/A by ground sprayer or in a minimum of 5 to 10 gallons/A by air.

Maximum benefit from GibGro is obtained when below normal temperatures predominate following application and growth would be otherwise slowed in untreated spinach.

NOTE: Since Gibberellic Acid can promote bolting, do not apply to spinach after the mid-winter period or if temperatures may be expected to exceed 75°F within several days of application. Do not apply on spring-planted spinach.

SPRAY GUIDELINES FOR FLORICULTURE CROPS

POMPOM CHRYSANTHEMUMS

(Florida) *For elongating peduncles on pompom chrysanthemums:* apply a single spray 4 to 5 weeks after initiation of short day conditions.

Use 1/2 to 1 gram a.i. in 12 gallons for application to 1,000 sq. ft. of bed (equivalent to 20 to 40 grams a.i. in 500 gallons/A).

Apply with overhead nozzles directing the spray to the flower buds.

NOTE: Overuse or incorrect timing may cause long, spindly and weak stems.

STATICE

(Florida) *To promote earlier flowering and to increase flower yield:* apply a single drench spray when plants are more than 10 inches in diameter (approximately 90 to 110 days after normal seeding time). Use 40 to 50 grams a.i. in 25 gallons to provide 10 ml (5mg a.i.) solution per plant.

NOTE: Do not exceed specified rates. Do not apply repeated sprays. Accelerated flowering is influenced by extended photoperiod, adequate nutrition and reduced night temperature. Treatment with gibberellins lessens the requirement for the cold requirement and/or the long photoperiod.

SPRAY GUIDELINES FOR ADDITIONAL CROPS

BERMUDAGRASS GOLF TURF

(Florida) *To initiate or maintain growth and prevent color change during periods of cold stress and light frosts on golf course Bermudagrass (e.g., Tidwarf, Tigreen, etc.):* apply 10 grams a.i. weekly or 25 grams a.i. biweekly in 25 to 100 gallons/A.

Use 1/4 to 2/3 gram a.i. in approximately 6 gallons appropriate for the spray equipment for application to 1,000 sq. ft. (equivalent to 10 to 25 grams a.i. per acre in 25 to 100 gallons/A).

NOTE: Do not exceed specified rates.

Do not apply during extended warm periods where night temperatures exceed 65°F.

Maintain adequate moisture and proper fertilization programs recommended in local area.

Discontinue treatments if thinning is observed.

Do not apply the high rate more frequently than every two weeks. More frequent mowing may be necessary.

Do not use on dormant turf.

GRAIN SORGHUM SEED TREATMENT

(EXCEPT CALIFORNIA)

For use as a seed treatment to break dormancy and allow germination under cold soil conditions. Use 0.25 to 1.00 grams a.i. per 100 pounds of seed. GibGro can be applied to dry seed with standard mist-treating equipment. Make certain the seed is completely and uniformly covered with GibGro. Fill the seed treatment tank with water to one-half the final tank mix volume. Add the required amount of GibGro, mixing thoroughly while adding water and other seed treatment products to the desired final volume.

Do not use treated seed for food, feed or oil purposes. An approved dye must be added to distinguish GibGro treated seed and prevent inadvertent use for food, feed or oil purposes. Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the Federal and State seed laws. GibGro is compatible with most commonly used fungicide seed treatments such as VITAVAX and DITHANE, standard dyes and sticker-binding agents. When preparing tank mixes, the user should ensure adequate physical compatibility and mixing characteristics.

HOPS

(For seeded and seedless Fuggle hops and similar varieties adapted to Oregon and the Northwest) *To increase yield and pickability:* apply spray when vine growth is five to eight feet in length. Use 4 to 6 grams a.i. in 100 to 150 gallons/A.

NOTE: Do not apply within three weeks of harvest.

RICE SEED TREATMENT (EXCEPT CALIFORNIA)

For use as a seed treatment on both semi-dwarf and tall rice varieties to promote germination, emergence and final stand densities when planted at greater depths where soil moisture levels are more adequate for germination. GibGro is particularly effective on semi-dwarf varieties such as 'L3030', 'Gulfmont', and 'Texmont'. This will also result in more uniform emergence thus allowing more accurate and efficient herbicide, fertilizer, fungicide and insecticide applications and may maximize yield and improve grain quality.

Apply only to rice seed intended for drill seeded or dry broadcast systems. Do not apply to rice used in a 24 hour presoak prior to broadcast. Do not use more than 2.0 grams a.i. per 100 pounds of seed. DO NOT USE TREATED SEED FOR FOOD, FEED, OR OIL PURPOSES.

Use 1.00 to 2.00 grams a.i. in 8 to 20 ozs water per 100 pounds of rice seed. GibGro can be applied to dry seed with standard mist-treating equipment. Best results are obtained using a higher treatment volume (12 to 20 ozs/cwt of seed) to insure the seed is completely and uniformly covered with GibGro. Fill the seed treatment tank with water to one-half the final tank mix volume.

An approved dye must be added to distinguish GibGro treated seed and prevent inadvertent use for food, feed or oil purposes. Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the Federal and State seed laws. GibGro is compatible with most commonly used fungicide seed treatments such as VITAVAX® and DITHANE®, standard dyes and sticker-binding agents. When preparing tank mixes, the user should ensure adequate physical compatibility and mixing characteristics.

RICE POST-EMERGENCE SEEDLING TREATMENT (EXCEPT CALIFORNIA)

For use as a postemergence seedling application on semi-dwarf rice varieties grown in the United States to promote more uniform and vigorous growth prior to permanent flooding. This will allow earlier flooding of drill or dry broadcast-seeded varieties and may reduce additional flushing costs associated with a delay in permanent flooding, weed infestations and the number of herbicide applications as well as promote earlier and more uniform grain maturity. GibGro application may result in a temporary lighter green foliage color due to accelerated growth rates.

Apply GibGro only to fields which have been drained of flood water taking precautions to avoid drift or accidental application to other crops. Use should be limited to a single application prior to permanent flood and should not be applied when rice is subject to stress conditions. GibGro can be tank mixed with Propanil when these commonly used rice herbicides are used according to their manufacturer's label recommendations. Use the more restricted labeling instructions.

GibGro may be applied at a rate of 1 to 3 grams a.i. per acre to semi-dwarf rice between the 1 to 4 leaf growth stage. Timing and dosage is based on environmental conditions and preferred permanent flood practice in relation to tiller development.

For best results, apply GibGro at a rate of 2 to 3 grams a.i. per acre in 1 to 2 leaf stage rice when permanent flood is desired before tiller development. Apply 1-3 grams a.i. per acre in 3 to 4 (4th leaf showing) stage rice when flooding will be made following initial tillering. Use higher rates when temperatures are likely to average 75°F or less during the 14 days after application. Either application will permit establishment of a permanent flood 7 to 10 days earlier.

Apply GibGro by fixed wing aircraft equipped with spray systems capable of producing a uniform medium to fine spray droplet pattern. Do not apply less than 10 gallons total spray volume per acre. Low pressure ground sprayers equipped with boom and flat fan nozzles and applying 10 to 15 gallons total spray volume per acre may be used.

GRAMS OF ACTUAL GIBBERELIC ACID/ACRE	TO	AMOUNT OF GIBGRO POWDER FORMULATION PER ACRE
Desired Actual Gibberellic Acid Concentration (grams a.i.) in Finished Spray (Per Acre)		GibGro 10% Powder Contains 1.0 gram a.i./10 Grams Formulated Product
1.0		10 grams
2.0		20 grams
4.0		40 grams
8.0		80 grams
10.0		100 grams
16.0		160 grams
20.0		200 grams
24.0		240 grams
32.0		320 grams
40.0		400 grams
50.0		500 grams
64.0		640 grams
80.0		800 grams
96.0		960 grams
100.0		1000 grams

GibGro 10% Powder

(160 grams/bottle)

Active Ingredient:

Gibberellic Acid 10% W/W

Equivalent to 16 grams a.i. of Gibberellic Acid per bottle.

EPA Reg. No. 55146-66

NOTICE TO USER

Seller makes no warranty, express or implied, of merchantability, fitness or otherwise concerning use of this product other than as indicated on the label. User assumes all risks of use, storage or handling not in strict accordance with accompanying directions.

AGTROL CHEMICAL PRODUCTS

7322 Southwest Freeway

Suite 1400

Houston, Texas 77074

713/995-0111