

55146-73

09-30-2009

1
39



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Mary Beth Endres
Nufarm Americas Inc.
150 Harvester Drive Suite 200
Burr Ridge, IL 60527

SEP 30 2009

Dear Ms. Endres:

SUBJECT: Ultra Flourish Agricultural Fungicide
EPA Registration Number 55146-73
Your submission dated July 16, 2009
OPPIN Decision Number 408360

The amendment referred to above, submitted in connection with registration under section (3) of the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) is acceptable provided you make the following changes:

Page 2 – First Aid:

In the "Note to Physician", delete "See Label for Additional Precautions and Directions for Use."

Page 2 and page 34:

Change "Engineering control statement" to "Engineering controls statement."

Page 4 – Directions for Use – Resistance Management:

Delete the third sentence: "Therefore, Nufarm cannot assume liability for crop damage resulting from resistant strains of fungi."

Page 4 – Directions for Use – Rotation (Plantback) Restrictions – Replanting:

Change: "If replanting is necessary, additional applications of this product may be made, provided that the total amount of mefenoxam applied does not exceed the maximum allowed for the specific crop." to "If replanting is necessary, additional applications of this product may be made, provided that the total amount of mefenoxam applied does not exceed the maximum application rate or seasonal amount allowed for the specific crop."

Page 5 – Directions for Use – Rotational Crop table:

Change the first column heading from "Rotation Crop" to "Rotational Crop."
Change the second column heading from "Planting Time From Last Application (Days)" to "Planting Time From Last Application."

Change the row entry "Leafy Vegetables (e.g., lettuce, spinach, celery)" to "Leafy Vegetables except Brassica (e.g., lettuce, spinach, celery)"
Add the text "0 days" to the big box immediately under the heading "Planting Time From Last Application."
In the "Planting Time From Last Application." column for the "Crops Not Intended for Food or Feed" change "0" to "0 days."

Page 5 – Directions for Use – Mixing Procedures:

Delete the second sentence, namely "Any liability . . . specifically disclaimed by manufacturer."

Page 6 – Directions for Use – Mixing Procedures - Compatibility:

Delete the second sentence, namely "Any liability . . . specifically disclaimed by manufacturer."

Page 6 – Directions for Use – General Application Instructions:

Change the sentence "Refer to the specific crop directions for use for application recommendations." to "Refer to the specific crop application directions."

Page 6 – Directions for Use – General Application Instructions:

Change the beginning of the second sentence in the Banded Applications subsection from "Some row-crop recommendations are based . . ." to "Some row-crop directions are based . . ."

In the first sentence in the In-Furrow applications subsection change "prouduct" to "product."

In the Fluid Ounces per Acre table, change the first (left-hand-most) column header from "Use Rate . . ." to "Product Use Rate . . ."

Change the beginning of the second sentence in the Instructions for Moving Ultra Flourish . . . Seed Zone" subsection from "Some crop directions recommend . . ." to "Some crop directions specify . . ."

Page 7 – Directions for Use – Application Through Irrigation Systems:

Change the beginning of the first sentence in second paragraph from "This product should be diluted with water . . ." to "Dilute this product with water . . ."

Change the first sentence under "Application Instructions" from "This product must be applied on the schedule specified in the specific crop use recommendations, not according to the irrigation schedule." to "This product must be applied on the schedule specified in the specific crop use directions, not according to the irrigation schedule."

Page 10 – Directions for Use – "Crops" Table - Apples:

In the "Soil Spray " and "Soil Drench" entries in the "Use Instructions" column, change "Application should be made . . ." and "Applications should be made . . ." to "Apply . . ."

In the second bullet of the bottom row change "Applications of this product should be made . . ." to "Apply this product . . ."

Page 10 – Directions for Use – "Crops" Table:

Add "Avocados - continued" to the empty box in the "Crop" column of the top row.

Page 14 – Directions for Use – "Crops" Table:

Add "Citrus - continued" to the empty box in the "Crop" column of the top row.

Page 14 – Directions for Use – “Crops” Table - Cole crops:

In the Irrigation (drip) entry under Use Instructions change “. . . at the recommended rates.” to “. . . at the labeled rates.”

Page 15 – Directions for Use – “Crops” Table:

Add “Cole crops - continued” to the empty box in the “Crop” column of the top row.

Page 15 – Directions for Use – “Crops” Table - Cotton:

In the In-furrow entry under Use Instructions change “. . . 3 to 6 pints of Terraclor 2E . . .” to “. . . labeled rates of Terraclor 2E . . .”

Page 16 – Directions for Use – “Crops” Table:

Add “Cucurbit vegetables - continued” immediately above “Muskmelon (cantaloupe, . . .” in the “Crop” column of the top row.

Page 17 – Directions for Use – “Crops” Table:

Add “Cucurbit vegetables - continued” to the empty box in the “Crop” column of the top row.

Page 18 – Directions for Use – “Crops” Table:

Add “Herbs - continued” immediately above “Curry, leaf . . .” in the “Crop” column of the top row.

Page 18 – Directions for Use – “Crops” Table – Cucurbit Vegetables:

Under Disease change “Damping-off” to “Damping-off”

Page 19 – Directions for Use – “Crops” Table:

Add “Leafy Vegetables - continued” immediately above “Cardoon . . .” in the “Crop” column of the top row.

Add “Damping-off (Pythium spp)” to the empty box under the “Diseases” heading.

Add “2 to 4” to the empty box under the “Rate (Pints per Acre)” heading.

Add “Follow specific application rate and directions for spinach” under the “Use Instructions” heading.

Page 20 – Directions for Use – “Crops” Table:

Add “Legumes (Succulent or Dried) - continued” immediately above “(grain, lupiin, . . .” in the “Crop” column of the top row.

Add “Damping-off (Pythium spp)” to the empty box under the “Diseases” heading.

Add “1 to 2” to the empty box under the “Rate (Pints per Acre)” heading.

Page 32 – Limitation of Liability:

Change the sentence: “To the extent consistent with applicable law, in no event shall manufacturer or seller be liable for special, incidental, or consequential damages, or for damages in their nature of penalties relating to the goods sold, including use, application, handling, and disposal.” to : “To the extent consistent with applicable law, manufacturer or seller shall not be liable for special, incidental, or consequential damages, or for damages in their nature of penalties relating to the goods sold, including use, application, handling, and disposal.”

Page 34 – Environmental Hazards statement:

Add the following heading and text immediately below “Do not contaminate water when disposing of equipment washwater or rinsate.”

“Groundwater Advisory Statement

This chemical is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.”

Page 34 – Directions for Use:

After the end of the “Physical or Chemical Hazards” section, add the heading “Directions for Use followed by the text:

“It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.”

Add the heading “Agricultural Use Requirements “ and the text “Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to label booklet under “Agricultural Use Requirements “ in the Directions for Use section for information about this standard. Place the heading and text in a box.

Please submit one copy of the final printed labeling before the product is released for shipment. A stamped copy of the label is enclosed for your records

If you have any questions, please contact Lisa Jones on my team by phone at (703) 308-9424 or by e-mail at jones.lisa@epa.gov.

Sincerely,



Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505P)

Enclosure: Stamped label

5
39

[BOOKLET COVER]

ULTRA FLOURISH®

AGRICULTURAL FUNGICIDE

For the control of certain diseases in various crops caused by the Oomycete class of fungi

ACTIVE INGREDIENT:

Mefenoxam: D-Alanine, N-(2,6-dimethylphenyl)-N-(methoxyacetyl)-, methyl ester 25.1%

OTHER INGREDIENTS: 74.9%

TOTAL: 100.0%

Contains Petroleum Distillates

Contains 2 pounds mefenoxam per gallon

KEEP OUT OF REACH OF CHILDREN

WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

SEP 30 2009

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.**

55146-73

EPA REG. NO. 55146-73
EPA EST. NO.

NET CONTENTS GALLONS

055146-00073.20090716.Add New Uses. Revised Per EPA

Manufactured For:
NUFARM AMERICAS INC. AGT DIVISION
150 Harvester Drive
Burr Ridge, IL 60527



[Page 1 of booklet to end]

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p> <p>You may also contact 1-877-325-1840 for emergency medical treatment information.</p>	
NOTE TO PHYSICIAN	
<p>If this product is ingested, lavage stomach. A slurry of activated charcoal in water can be left in the stomach. Contains petroleum distillate – vomiting may cause aspiration pneumonia. See Label for Additional Precautions and Directions for Use.</p>	

**PRECAUTIONARY STATEMENT
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING - AVISO**

Causes substantial but temporary eye injury. Harmful if swallowed or absorbed through skin. Wear protective eyewear such as goggles or face shield. Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long -sleeved shirt and long pants
- Chemical-resistant gloves such as barrier laminate or viton
- Shoes plus socks
- Protective eyewear

See engineering controls for additional requirements.

USER SAFETY REQUIREMENTS

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statement:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Apply only as specified on this label. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

GROUNDWATER ADVISORY STATEMENT

This chemical is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statement of this label about personal protective equipment (PPE) and restricted-entry interval (REI). The requirements in this box only apply to users of this product that are covered by the WPS.

Do not enter or allow worker entry into treated areas during the REI of 48 hours.

Exception: If the product is soil-injected or soil-incorporated, the WPS, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: Coveralls, chemical-resistant gloves such as barrier laminate or Viton, shoes plus socks and protective eyewear

Failure to follow the directions for use and precautions on this label may result in crop injury, poor disease control, or illegal residues.

GENERAL USE PRECAUTIONS

This product is a systemic fungicide for use on selected crops to control certain disease caused by members of the Oomycete class of fungi. Other fungicides must be used to control diseases incited by other classes of fungi.

- This product is not to be used in foliar applications, unless specified on this label.

8
39

- Where rate ranges are specified on this label, use the higher rate when heavy disease pressure is expected and the lower rate when disease pressure is expected to be light unless otherwise noted.
- Under conditions of severe disease pressure when more than the label-specified number of applications of this product may be required, make additional fungicide treatments with other fungicides registered for the crop/disease found on this label.
- To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, as crop injury may result.
- Do not use this product in greenhouses, lath houses, float houses, and hydroponic facilities.
- Do not use this product for disease control in bedding plants, transplant trays or nurseries.
- Do not dip plants or roots, spray bare roots, or use a transplant water treatment with solutions of this product.

RESISTANCE MANAGEMENT

This product is a systemic fungicide having a specific mode of action and could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, Nufarm cannot assume liability for crop damage resulting from resistant strains of fungi. If treatment is not effective following the use of this product as recommended, a resistant strain of fungi may be present. If the treatment is ineffective due to the presence of a mefenoxam resistant strain of fungi, neither this product nor any other fungicide with similar action will effectively control that disease. Consideration should then be given to the prompt use of other types of suitable fungicides. Do not make foliar applications to field grown tobacco, or other crops, unless specified since this practice may encourage more rapid development of resistance. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance on your particular crop and disease control situation.

ROTATION (PLANTBACK) RESTRICTIONS

REPLANTING

If replanting is necessary, additional applications of this product may be made, provided that the total amount of mefenoxam applied does not exceed the maximum allowed for the specific crop.

ROTATION (PLANTBACK) RESTRICTIONS

Do not plant any crop that is not registered for use with mefenoxam in mefenoxam-treated soil for a period of 12 months unless a shorter interval is specified on the following list.

9
39

ROTATION CROP	PLANTING TIME FROM LAST APPLICATION (Days)
Alfalfa (including birdsfoot trefoil)	0
Asparagus	
Brassica (Cole Crop) Vegetables (e.g., broccoli, cabbage, cauliflower)	
Clover	
Cotton	
Cucurbit Vegetables	
Fruiting Vegetables	
Garlic	
Globe Artichoke	
Herbs (fresh and dried)	
Leafy Vegetables (e.g., lettuce, spinach, celery)	
Legume Vegetables (e.g., beans and peas, succulent and dried)	
Onions (dry bulb, green)	
Peanuts	
Pineapples	
Root and Tuber Vegetables	
Soybeans	
Strawberries	
Sunflower	
Cereal Grains (other than Corn)	14 days
Corn	9 months
Crops Not Intended for Food or Feed	0
All Other Crops Intended for Food or Feed	12 months

MIXING PROCEDURES

Add 1/4 to 1/2 of the required amount of water to the spray tank, add the proper amount of this product, then add the rest of the water. When tank mixing other products with this product, follow the proper sequence of adding products to the spray tank. Wettable powders or water dispersible granules should be added to the water in the tank first, followed by flowable products, with emusifiable concentrates, such as this product, added last. Provide sufficient agitation during mixing and application to maintain a uniform emulsion.

Unless otherwise prohibited on this label or the label of an intended tank mix product, this product may be applied in combination with any pesticide registered for the same crop, timing, and method of application. Observe the most restrictive label statements of various tank mix products used.

IMPORTANT: PESTICIDE TANK MIXES MAY INCREASE THE RISK OF MIXING INCOMPATIBILITIES, REDUCED EFFECTIVENESS AND/OR CAUSE CROP INJURY OR LOSS. ANY LIABILITY FOR LOSS, INJURY OR DAMAGE RESULTING FROM A TANK MIXTURE NOT SPECIFIED ON THIS LABEL OR IN MANUFACTURER'S SUPPLEMENTAL LABELING DISTRIBUTED FOR THIS PRODUCT IS SPECIFICALLY DISCLAIMED BY MANUFACTURER.

COMPATIBILITY

Before full-scale mixing of this product with other pesticides, fertilizers, secondary plant nutrients, adjuvants, surfactants or oils, you must determine the compatibility of the proposed mixture. Use proportionate quantities of each ingredient and mix in a small container. Always mix one product thoroughly with the diluent before adding another product. If no incompatibility is evident after 30 minutes, the mixture is generally compatible for spraying. To evaluate potential short term effects of applying the mixture, test the tank mix combination on a few plants or a small area before larger-scale treatments. Wait at least 2 to 3 days for problems to become apparent.

IMPORTANT: MIXING WITH OTHER SUBSTANCES MAY INCREASE THE RISK OF MIXING INCOMPATIBILITIES, REDUCED EFFECTIVENESS AND/OR CAUSE CROP INJURY OR LOSS. ANY LIABILITY FOR LOSS, INJURY OR DAMAGE RESULTING FROM A MIXTURE NOT SPECIFIED ON THIS LABEL OR IN MANUFACTURER'S SUPPLEMENTAL LABELING DISTRIBUTED FOR THIS PRODUCT IS SPECIFICALLY DISCLAIMED BY MANUFACTURER.

GENERAL APPLICATION INSTRUCTIONS

Apply this product by ground or air in sufficient water or liquid fertilizer to provide uniform coverage of the soil surface. Apply in a minimum of 20 gallons per acre for ground applications and 5 gallons per acre by air. Refer to the specific crop directions for use for application recommendations.

Banded Applications: For banded applications, the area actually is the area covered by the band, not total cropland area planted. Some row-crop recommendations are based on treating in-the-row and these rates generally are specified as amounts (fluid ounces) of product per certain row length (often 1,000 feet). Others express rates as amount per treated acre, which means the total area treated with the pesticide. If rates are expressed as amount per treated acre and banded applications are used, the amount of pesticide used per acre will be proportionately less. The following formula can be used to calculate the amount of this product needed per acre of crop when banded applications are made.

Calculate the amount of this product needed for band treatment by the formula:

$$\frac{\text{band width in inches}}{\text{row spacing in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$$

In-Furrow Applications: For in-furrow applications, apply this product as an in-furrow spray in 3 to 7 gallons of water per acre at planting. Mount the spray nozzle so the spray is directed into the furrow just before the seeds are covered. Following is a table of the common row spacings and the amount of this product applied per acre.

Use Rate fl. oz./1,000 row feet (oz. AI/1,000 row feet)	ULTRA FLOURISH (Fluid Ounces per Acre)								
	20 inch rows	22 inch rows	24 inch rows	30 inch rows	32 inch rows	34 inch rows	36 inch rows	38 inch rows	40 inch rows
0.15 (0.038)	4.0	3.6	3.2	2.6	2.5	2.3	2.2	2.1	2.0
0.3 (0.075)	7.8	7.0	6.5	5.2	5.0	4.6	4.3	4.1	4.0
0.56 (0.14)	14.5	13.3	12.2	9.8	9.1	8.5	8.1	7.7	7.3
0.84 (0.21)	22.0	20.0	18.3	14.5	13.7	13.0	12.2	11.6	11.0
20" = 26,136 row ft./acre			22" = 23,760 row ft./acre			24" = 21,780 row ft./acre			
30" = 17,424 row ft./acre			32" = 16,315 row ft./acre			34" = 15,374 row ft./acre			
36" = 14,520 row ft./acre			38" = 13,754 row ft./acre			40" = 13,068 row ft./acre			

Instructions for Moving Ultra Flourish into the Root or Seed Zone

To ensure maximum activity on soilborne pathogens, this product must be moved into the seed or root zone of the plant. Some crop directions recommend incorporating the product to move it to the seed or root zone (preplant incorporated application, soil drenches, shank applications) while others place the fungicide into the seed or root zone (in-furrow sprays, soil injections, crown dips). For applications made to the soil surface, rainfall will move the fungicide to the seed or root zone. However, if rainfall is not expected within 24 hours after application, mechanically incorporate before planting or sprinkler-irrigate after planting with 1/2 to 1 inch of water.

APPLICATION THROUGH IRRIGATION SYSTEMS

This product alone or in combination with other pesticides, which are registered for application through irrigation systems, may be applied through irrigation systems. Apply this product only through center pivot, solid set, hand move, moving wheel, micro-sprinkler, pressurized drench (flood) or drip (trickle), micro-irrigation such as a spaghetti-tube or individual tube irrigation, hand-held calibrated irrigation equipment such as the hand-held wand with injector, calibrated overhead watering booms, ebb and flow or bench flooding sub-irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

This product should be diluted with water on a 1/10 basis prior to injection into an irrigation system, proper tank-mix agitation is required during this mixing procedure.

If you have any questions about calibration, you should contact the State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide applications to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

OPERATING INSTRUCTIONS

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back-flow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

APPLICATION INSTRUCTIONS

This product must be applied on the schedule specified in the specific crop use recommendations, not according to the irrigation schedule.

With the exception of citrus, this product has not been sufficiently tested when applied through irrigation systems to assure consistent product performance for all labeled uses. The following calibration and application techniques are provided for user reference, but do not constitute a warranty of fitness for application through sprinkler or drip irrigation equipment. Users must check with state and local regulatory agencies for potential use restrictions before applying any agricultural chemical through sprinkler or drip irrigation equipment.

NOTE: Do not inject this product at full strength or deterioration of valves and seals may occur. Use a dilution ratio of at least 10 parts water to 1 part of this product in the mix tank. This product is corrosive to many seal materials. Leather seals are best. EPDM or silicone rubber seals can be used, but should be replaced once a year. Do not use Viton, Buna-N, Neoprene or PVC seals.

12
39

CENTER PIVOT IRRIGATION EQUIPMENT Use only with drive systems, which provide uniform water distribution

Determine the size of the area to be treated. Determine the time required to apply 1/2 to 1 inch of water over the area to be treated when the system and injection equipment are operated at normal pressures recommended by the equipment manufacturer. Run the system at 80 to 95% of the manufacturer's rated capacity. Using water, determine the injection pump output when operated at normal line pressure. Determine the amount of this product required to treat the area covered by the irrigation system. Add the required amount of this product and sufficient water to the solution (mix) tank to meet the injection time requirements. Maintain constant solution tank agitation during the injection period. Continue to operate the system until all of the product solution has cleared the most distant sprinkler head.

SOLID SET, HAND MOVE, AND MOVING WHEEL IRRIGATION EQUIPMENT

Determine the acreage covered by the sprinklers. Fill injector solution tank with water and adjust flow rate to use the contents over a 20 to 30 minute interval. Determine the amount of this product required to treat the area covered by the irrigation system. Add the required amount of this product into the same quantity of water used to calibrate the injection period. Operate the system at the same pressure and time interval established during the calibration. Inject this product at the end of the irrigation cycle in 1/2 to 1 inch of water or as separate application to maximize the effectiveness of the fungicide. Stop injection equipment after treatment is completed. Continue to operate the system until all of the this product solution has cleared the last sprinkler head.

MICRO SPRINKLER, DRIP IRRIGATION SYSTEMS, OVERHEAD WATERING BOOMS

General Instructions

Each run of the irrigation system must be calibrated separately to determine the time it takes water to move through the system and to make sure all emitters in the system are putting out the same amount of water. Only pressure injection or venturi equipment is recommended. Determine the area to be treated in each irrigation run. Measure the output of each of the emitters or drip tubes closest to and farthest from the injector site. For calibration, substitute a concentrated detergent (such as Wisk) or a soluble fertilizer for the this product in the injector tank. The detergent will bubble as it leaves the emitters. The time period over which bubbles occur should be checked for both the closest and farthest emitters. If these times are not within 2 minutes of each other, adjust the dilution ratio and/or the injection rate. If a soluble fertilizer is used, measure the time intervals with a salt bridge. If a drip system is being calibrated, substitute soluble fertilizer for this product in the injector and measure the time intervals with a salt bridge. (Step by step instructions) Before starting to calibrate, operate the system until all the emitters are putting out equal flow rates or until the system is operating at full pressure. Make up an indicator solution of detergent or fertilizer, using the same ratio to be used when mixing this product. Set the injector to apply the indicator solution at the injection rate to be used in the actual product application. Attach a 5 inch length of flexible tubing over the emitter closest to the injection point, another length over the emitter farthest away. Both emitters should be monitored to determine the time intervals that the indicator solutions are observed. Begin injecting the indicator solution. Direct the flow from the tubes at the emitters into a small container. Begin timing when the indicator solution is first detected, stop timing when the indicator solutions are no longer detected. If the period of detection of the indicator solution between the 2 emitters is within 2 minutes of each other, comparable coverage will be obtained. If they are not, make adjustments by increasing the dilution ratio, using more water per part of product, or adjust the injector to a slower flow rate. Once the system is calibrated, dilute the needed amount of this product with water using a minimum of 10 part of water to 1 part this product. Do not begin to inject this product into the system until all emitters are producing equal flow rates, or until the system is at full pressure. Inject this product into the system at the end of the irrigation set in 1/2 to 1 inch of irrigation water.

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days of the year.

Chemigation systems connected to public water systems must contain a functional, reduced pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the

point of pesticide introduction. As an option to the RPZ, the water from the public water systems should be discharged into a reservoir tank prior to pesticide introduction.

There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where the pesticide distribution is adversely affected.

Systems must use a metering pump, as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

CROPS

Use on the crops listed below is for field use only and is not intended for use on transplant trays, dips, greenhouses, lath houses, float houses, hydroponic production, or in bedding plant structures.

Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation.

For banded applications, use the formula in the General Instructions section of this label to calculate the amount of this product needed per acre for the band width actually used.

Equivalents: 1 quart = 2 pints.

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS						
ALFALFA (Including birdsfoot trefoil)	Damping-of (<i>Pythium</i> spp.)	0.5 to 1	Apply as a broadcast surface spray at planting in a sufficient amount of water for thorough coverage. If inter-seeding alfalfa into existing stands for renovation, apply 0.5 pint per acre as a broadcast spray at planting. If alfalfa seed was previously treated with a metalaxyl or mefenoxam seed treatment product, use this product at 0.5 pint per acre at planting.						
	Root rot (<i>Phytophthora</i> spp.)								
<ul style="list-style-type: none"> Do not feed green forage or cut hay for 60 days following application. 									
APPLES (Bearing and Nonbearing Trees)	Crown, Collar and Root rot (<i>Phytophthora</i> spp.)	8	Soil Spray (Broadcast or Banded): Apply as a broadcast spray in sufficient water to obtain thorough coverage. Treated area is based on the area under the tree canopy or the area of the sprayed row. Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation. Application should be made in early spring before growth starts and in the fall after harvest but before the ground freezes.						
		or 3 fluid ounces per 1,000 square feet							
		1	Soil Drench: Mix in 100 gallons of water. Apply around the trunk of each tree. Applications should be made in early spring before growth starts and in the fall after harvest but before the ground freezes. On new plantings, delay the first application until 2 weeks after planting.						
			<table border="1"> <thead> <tr> <th>Trunk Diameter 12 Inches Above Soil Line</th> <th>Pints of Diluted Mixture Per Tree</th> </tr> </thead> <tbody> <tr> <td><1</td> <td>2</td> </tr> <tr> <td>1 to 3</td> <td>6</td> </tr> <tr> <td>>5</td> <td>8</td> </tr> </tbody> </table>	Trunk Diameter 12 Inches Above Soil Line	Pints of Diluted Mixture Per Tree	<1	2	1 to 3	6
Trunk Diameter 12 Inches Above Soil Line	Pints of Diluted Mixture Per Tree								
<1	2								
1 to 3	6								
>5	8								
<ul style="list-style-type: none"> Use of this product will aid in control when used in conjunction with good cultural practices and with root stocks that are tolerant to the disease. Applications of this product should be made before symptoms appear especially in areas of the orchard favorable for disease development. This product will not revitalize trees showing moderate to severe disease symptoms. Do not graze or feed cover crops in treated orchards. 									

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
ARTICHOKE, GLOBE	Damping-of (<i>Pythium</i> spp.)	2 to 4	Apply as a broadcast soil spray at planting in a sufficient amount of water to provide thorough coverage.
	Root rot (<i>Phytophthora</i> spp.)		Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation.
<ul style="list-style-type: none"> Do not apply this product within 200 days of harvest. 			
ASPARAGUS	Crown rot Spear rot (<i>Phytophthora</i> spp.)	2	Apply as a broadcast or banded spray to soils in a minimum of 10 gallons of water over the beds. Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation.
			Cutting Beds: Apply 30 to 60 days before first cutting. For additional control, make another application just before the beginning of harvest.
			New Plantings: Apply after planting seedlings or after covering one year old crowns.
<ul style="list-style-type: none"> Do not apply this product within 1 day of harvest. 			
AVOCADOS	Root rot (<i>Phytophthora</i> spp.)	0.5 fluid ounces	Sleeve Drench: Mix with 18 gallons of water. At the time of transplanting, drench the roots inside the sleeve with 1 quart of diluted mixture per tree. The sleeve drench will not replace the soil surface applications for long-term control of root rot.
		4 to 8	Soil Spray: Apply to the soil under the drip emitter. Irrigate to incorporate residues into the soil. If there is more than one emitter per tree, distribute the total amount of product needed among the emitters.
		or 1 to 2 fl. oz. per 1,000 gallons (3.9 - 7.8 ppm ai)	Injection (Drip Irrigation): Inject this product into the irrigation water at the timing specified below.
			Sprinkler Irrigation: Apply the specified amount of product to the soil directly under the tree canopy. For trees with a canopy diameter of 2 ft, use 2 pints/A. As the canopy diameter increases, use a higher rate. For trees with a canopy diameter of 15 ft., use 8 pints/A. Time applications to occur at the start of the growing season, at transplanting or when <i>Phytophthora</i> is detected in soils. Up to two additional applications may be made every 3 months. No fungicide treatment is required during the months of November through February.

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
<ul style="list-style-type: none"> • For best results, use this product as soon as soil test indicate the presence of <i>Phytophthora</i>. • For new plantings, the use of <i>Phytophthora</i>-resistant rootstocks with this product is recommended. Mature trees in moderate to advanced stages of decline cannot be cured with this product. • Do not apply more than 24 pints (3 gallons) per acre per crop (6 lb. a.i./A) of soil-applied mefenoxam products. • Do not make an application within 28 days of harvest. 			
BLUEBERRIES	Root rot (<i>Phytophthora</i> spp.)	7.2 or 0.5 per 1,000 linear feet	Established Plantings: Apply 7.2 pints per acre or 0.5 pints per 1,000 linear ft. to soil in a 3 foot band over the row before plant growth starts in the spring. Make one additional application if needed to coincide with a period favorable for root rot. Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation.
		7.2	New Plantings: Apply 7.2 pints per acre as a band or broadcast application to the soil at planting. For banded applications, use an 18 inch band over the row. Make one additional application if needed to coincide with a period favorable for root rot. Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation.
<ul style="list-style-type: none"> • Use this product in conjunction with good cultural practices to minimize disease problems. • This product will not revitalize plants showing moderate to severe root rot symptoms. • This product may be applied on the day of harvest. • Do not soil-apply more than 7.2 pints per acre per crop (3.6 lb. a.i./A). 			
CARROTS	Damping-of Root Dieback (<i>Phythium</i> spp.)	2 to 4	Apply preplant as a broadcast spray or banded application to soil in a sufficient amount of water or liquid fertilizer and mechanically incorporate in the top 2 inches of soil. Apply to soils at planting or prior to emergence by broadcast or banded applications in water or liquid fertilizers. Move into soil as recommended in General Application Instructions. For banded applications, a 7 inch band is recommended.
			<ul style="list-style-type: none"> • Do not apply more than 4 pints per acre per crop (1.0 lb. a.i./A) if soil-applied and more than 0.2 pint per acre per crop (0.4 lb. a.i./A) if foliar-applied.

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
CITRUS Including: Calamodin Chironja Citrus citron Grapefruits Kumquats Lemons Limes Mandarin Oranges Pummelo Tangelos Tangerines Tangor and Hybrids of these	Brown Rot Citrus foot rot Root rot Trunk Cankers Gummosis (<i>Phytophthora</i> spp.)	4 to 8	RESETS OR NEW PLANTINGS Apply at planting in sufficient water to obtain uniform coverage of the soil. Apply spray to the soil surface beneath the tree canopy or apply through irrigation water (micro-sprinkler or drip). Up to two more applications may be made every 3 months when root growth flushes occur. Florida, Texas and Puerto Rico: Apply 4 pints/A as a soil spray or 1 pint per grove acre via injection. Make a total of 2 to 3 applications per year at one of these schedules: spring/summer, summer/fall or spring/summer/fall.
		2 to 3 fluid ounces	Water Ring Drench: Mix in 100 gallons of water. Apply 5 gallons of the mix around the base of each tree within the watering ring of resets or new plantings.
		2 to 12	Established Plantings Soil Application: Apply in sufficient water to obtain uniform coverage of the soil. Apply spray to the soil surface beneath the tree canopy or apply through irrigation water (micro-sprinkler or drip). Time applications to begin in the spring at root flush. Up to two more applications may be made when root growth flushes occur using 2 to 4 pints per acre (when a total of 3 applications are made) or 6 pints per acre (when a total of 2 applications are made). Florida, Texas and Puerto Rico: Apply 2 pints per acre as a soil spray or 1/2 pint per grove acre via injection to groves that have a Phytophthora count of 10 to 20 propagules per cubic centimeter of soil. If more than 20 propagules per cubic centimeter are present, use 2 pints per acre as a spray or 2 pints per grove acre through injection. Make a total of 2 to 3 applications per year at one of these schedules: spring/summer, summer/fall or spring/summer/fall.
	Gummosis	4	Trunk Spray: Mix in 3 gallons of water and spray the surface of the trunks using enough spray to thoroughly wet the cankers. Apply up to 3 times per year.
		2	Trunk Spray (FL only): Mix in 10 gallons of water and spray the surface of the trunks using enough spray to thoroughly wet the cankers. Apply up to 3 times per year.

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
			<ul style="list-style-type: none"> • Additional directions for use are available from local extension agents. • Where nematodes are a problem, best results can be achieved if effective EPA registered nematicides are used. Nematicides can be used in combination or in sequence with applications of this product. • For best Phytophthora control, a combination of cultural practices and resistant varieties is recommended. The use of this product is not recommended in Florida for use on the highly susceptible sweet orange rootstock. • In Florida, Texas and Puerto Rico, phytotoxicity to citrus resets or new plantings (plants less than 5 years old) may be prevented by using rates lower than 4 pints per acre. • Do not use on highly susceptible sweet orange rootstock in Florida. • Do not apply to bare roots. • Do not make trunk gummosis sprays and soil applications to the same tree in the same crop season. • For trees less than 3 years old, do not tank mix herbicides with this product. First apply the herbicide and after 3 to 4 weeks apply this product. • Do not apply more than 24 pints (3 gallons) of this product per acre per crop (6 lb. a.i. /A).
CLOVER	Damping off (<i>Pythium</i> spp.) Root rots (<i>Phytophthora</i> spp.)	0.5 to 1	Soil Spray (broadcast): Apply as a broadcast surface spray at planting. If seed was previously treated with mefenoxam seed treatment product, use 0.5 pint per acre rate at planting. Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation.
	<ul style="list-style-type: none"> • Do not feed green forage or cut hay for 90 days following application. 		
COLE CROPS Including: Broccoli Broccoli, Chinese (gai lon) Broccoli Raab (rapini) Brussels sprouts Cabbage Cabbage, Chinese (bok choy and napa) Cauliflower Cavalo Broccoli Collards Kale Kohlrabi Mizuna	Damping off (<i>Pythium</i> spp.)	0.5 to 1	Preplant Incorporated Application: Apply as a broadcast or banded soil application in sufficient water or liquid fertilizer to provide uniform coverage and incorporate into the top 2 inches of soil. For banded applications, a 7 inch band is recommended.
	Basal stem rot (<i>Phytophthora</i> spp.)	2 to 4	Preplant Incorporated Application: Apply as a broadcast or banded soil application in sufficient water or liquid fertilizer to provide uniform coverage and incorporate into the top 2 inches of soil. For banded applications, a 7 inch band is recommended. Surface Application: Apply at planting in sufficient water or liquid fertilizer to provide uniform coverage. For banded application, a 7 inch band is recommended. Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation. Irrigation (drip): Inject this product into the irrigation water at the recommended rates.

19
39

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
Mustard Greens Mustard Spinach Rape Greens All hybrids and varieties of these	<ul style="list-style-type: none"> Do not apply more than 4 pints per acre per crop (1 lb a.i./A) for soil applications and 2 pints per acre per crop (0.5 lb a.i./A) as a foliar spray. 		
COTTON	Seed and Root Rot (<i>Pythium</i> spp.) Seedling Blight (<i>Pythium</i> spp.)	0.15 to 0.30 fl. oz. per 1,000 row ft.	In-furrow: Apply in 5 to 15 gallons of water or liquid fertilizer at planting. Mount the spray nozzle so the spray is directed into the furrow over the seed just before the seeds are covered. For broader spectrum seedling disease, tank-mix this product with 3 to 6 pints of Terraclor 2E® per acre, or as a unit pack alternative, use labeled rates of Terraclor Super X EC or Terraclor Super X® 18.8 Granule.
CRANBERRIES	Phytophthora Root Rot (<i>Phytophthora</i> spp.)	2 to 3.5	Soil Spray (Broadcast): Apply 3 times using ground or chemigation equipment. Time applications to occur first in the fall after harvest, then in the spring with a final application up to, but not later than, 45 days prior to harvest. Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation.
	<ul style="list-style-type: none"> Do not apply this product to cranberries by air. Do not apply within 45 days before harvest Do not apply more than 10.5 pints per acre per crop (2.65 lb. a.i./A). 		
CUCURBIT VEGETABLES Including Chayote Chinese waxgourd Citron melon Cucumber Gherkin, Gourd, edible <i>Momordica</i> spp. (balsam apple, balsam pear, bitter melon Chinese cucumber)	Damping-off Cottony leak (<i>Pythium</i> spp.)	2 to 4	Pre-plant Incorporated Application: Apply as a broadcast or banded soil application in sufficient water or liquid fertilizer to provide uniform coverage and incorporate in the top 2 inches of soil. For banded application, a 7 inch band is recommended. Surface Application: Apply at planting in sufficient water or liquid fertilizer to provide uniform coverage. Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation. For banded application, a 7 inch band is recommended. Injection (drip): This product may also be injected into the irrigation water at the appropriate rates. Two additional applications are permitted every 30 days after the initial at-planting application.

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
<p>Muskmelon (cantaloupe, casaba, Crenshaw, Honeydew, Honey balls, Mango, Persian, pineapple, Santa Claus, snake melons)</p> <p>Pumpkin Squash, summer Squash, winter Watermelon All hybrids and varieties of these</p>	<p>Root rot (<i>Pythium</i> spp.)</p>	<p>0.5 to 0.8</p>	<p>Soil Directed Spray: If soil application was made at planting, two additional applications may be made at 20 to 30 day intervals during the season. Direct the spray to the base of the plants and cover 6-8 inches of the soil on either side of the plants. Incorporate using either mechanical means or through irrigation to move the product into the root zone.</p> <p>Injection (drip): This product may also be injected into the irrigation water at the appropriate rates.</p> <ul style="list-style-type: none"> • Do not apply within 5 days of harvest. • Do not exceed 4 pints per acre per crop (1 lb. a.i./A) from all methods of application combined.
<p>FRUITING VEGETABLES (Except Cucurbits) Including Eggplant Groundcherry Pepper (bell, chili, cooking pepper, pimento, sweet) Tomatillo Tomato (see Tomato Section below)</p>	<p>Damping-off (<i>Pythium</i> spp.)</p> <p>Crown rot (<i>Phytophthora capsici</i>)</p>	<p>2</p>	<p>To control Crown Rot, apply before the plants are infected to obtain satisfactory control. Plants already infected with <i>Phytophthora capsici</i> cannot be cured with this product.</p> <p>Soil Spray (broadcast or band): Apply in water or liquid fertilizer preplant or at planting. For direct seeded peppers, apply preplant or prior to emergence. Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation. For banded applications, use 12 to 16 inch band.</p> <p>Injection (drip irrigation): Inject this product into the irrigation water. Make up to 2 additional applications on a 30 day schedule following initial application at planting.</p> <p>Banded Spray: For Crown Rot, make 2 post directed applications at 30 day intervals after transplanting. Direct the spray to the base of the plants and cover 6 to 8 inches of the soil on either side of the plants. Incorporate mechanically or sprinkler-irrigate to move the product into the root zone.</p> <p>Shank Application: For Crown Rot, apply in liquid fertilizer, shanked in as a banded treatment to either side of the plant.</p>

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
			<ul style="list-style-type: none"> • This product may cause some yellowing of the pepper leaves. • Plants already infected with <i>Phytophthora</i> cannot be cured with this product. • The foliar blight phase of <i>Phytophthora</i> cannot be controlled with foliar applications of this product. • To avoid possible illegal residues do not apply within 7 days of harvest. • Do not soil-apply more than 6 pints of this product per acre per crop (1.5 lb a.i./A) or foliar apply more than 2 pints per acre per crop (0.5 lb a.i./A).
GINSENG	Phytophthora root rot (<i>Phytophthora cactorum</i>)	1.5	<p>Drench: Apply in 100 to 400 gallons of water uniformly to the soil surface in the spring before the plants begin to grow.</p> <ul style="list-style-type: none"> • Do not make more than 1 application of this product.
GRASSES, FORAGE, FODDER, HAY*	Seeding diseases (<i>Pythium</i> spp.)	0.5 to 2	<p>Soil Spray (broadcast): Apply as a broadcast soil spray at planting. Use 0.5 to 1 pints per acre if grass seed was previously treated with a mefenoxam seed treatment product. Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation.</p> <p>*Includes any enclosed pasture grasses or grasses grown for hay or silage such as Bermudagrass, bluegrass, bromegrass, or fescue.</p> <ul style="list-style-type: none"> • Do not graze, feed green forage, or cut for hay for 60 days following applications. • Do not apply to range grasses.
HERBS (Fresh and Dried) Including: Angelica Balm Basil Borage Burnet Chamomile Catnip Chervil (dried) Chinese Chive Chive Cilantro (leaf) Clary Coriander (leaf) Costmary	Damping-off (<i>Pythium</i> spp.)	2 to 4	<p>Pre-plant Incorporated Application: Apply as a broadcast or banded soil application in sufficient water or liquid fertilizer to provide uniform coverage and mechanically incorporate in the top 2 inches of soil. For banded application, a 7 inch band is recommended.</p> <p>Surface Application (broadcast or band): Apply in sufficient water or liquid fertilizer to provide uniform coverage. Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation. For banded application, a 7 inch band is recommended.</p> <p>Banded Spray: Direct sprays to baseline of plants covering 6-8 inches on each side of plants (12 to 16 inch band width/row). Apply 28 days after planting or after first cutting.</p>

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
Curry (leaf) Dillweed Horehound Hyssop Lavender Lemongrass Lovage (leaf) Marigold Marjoram Nasturtium Parsley (dried) Pennyroyal Rosemary Rue Sage Savory, summer and winter Sweet Bay Tansy Tarragon Thyme Wintergreen Woodruff Wormwood	<ul style="list-style-type: none"> • Do not apply within 21 days of harvest. • Do not exceed 8 pints per acre (2 lb a.i./A) per crop from all soil-applied applications. 		
HOPS	Downy mildew <i>(Pseudoperonospora humulii)</i>	1	<p>Soil Drench: Apply in a sufficient amount of water or liquid fertilizer for thorough coverage. Apply to the soil surface over the crowns after pruning, but before training. Early application before shoots are 6 inches long provides best protection.</p> <p>Foliar Spray: Apply using ground equipment in a minimum of 50 gallons water per acre. Treat when primary infection (spikes) persists after a soil drench treatment and/or there is evidence of secondary (foliar) infection and apply this product in combination with a contact copper fungicide.</p> <ul style="list-style-type: none"> • Do not soil-apply more than 1 pint per acre per crop (0.25 lb a.i./A) or foliar-apply more than 2 pints per acre per crop (0.5 lb a.i./A). • Do not make the last application within 45 days of harvest. • Do not apply foliar sprays of this product without a copper fungicide registered for use on hops.
LEAFY VEGETABLES (Except Brassica) Including Amaranth Arugula	Damping-of <i>(Pythium spp.)</i>	2 to 4	Preplant Incorporated (broadcast or band): Apply as a soil application in sufficient water or liquid fertilizer to provide uniform coverage and mechanically incorporate in the top 2 inches of soil. For banded application, a 7 inch band is recommended.

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
Cardoon Celery Celery, Chinese Celtuce Chervil Chrysanthemum, edible-leaved Chrysanthemum, garland Corn salad Cress, garden Cress, upland Dandelion Dock (Sorrel) Endive (Escarole) Fennel, Florence (finochio) Lettuce - head and leaf Orach Parsley Purslane, garden Purslane, winter Radicchio (red chicory) Rhubarb Spinach (see below) Spinach, New Zealand Spinach, vine Swiss Chard			Surface Application (broadcast or band): Apply at planting to soil surface in sufficient water or liquid fertilizer to provide uniform coverage. For banded application, a 7 inch band is recommended. Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation.
<ul style="list-style-type: none"> • Do not harvest within 7 days of application. • Do not soil-apply more than 4 pints per acre (1 lbs. a.i./A) or foliar-apply more than 1.6 pints per acre (0.4 lb a.i./A) per crop. 			
LEGUME VEGETABLES (Succulent or Dried) Including Bean (<i>Lupinus spp.</i>)	Damping off Root rot (<i>Pythium spp.</i>)	1 to 2	Preplant Incorporation (broadcast or band): Apply as a broadcast soil application in sufficient water or liquid fertilizer to provide uniform coverage and mechanically incorporate in the top 2 inches of soil. For banded application, a 7 inch band is recommended.

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
<p>(grain, lupin, sweet lupin, white lupin, white sweet lupin)</p> <p>Bean (<i>Phaseolus</i> spp.) (field, kidney, lima, navy, pinto, runner, snap, tepary, wax)</p> <p>Bean (<i>Vigna</i> spp.) (adzuki, asparagus, blackeyed pea, catjang, Chinese longbean, cowpea, crowder pea, moth, mung, rice, southern pea, urd, yardlong)</p> <p>Broad Bean (fava bean)</p> <p>Chickpea (garbanzo bean)</p> <p>Guar</p> <p>Jackbean</p> <p>Lablab bean (hyacinth bean)</p> <p>Lentil</p> <p>Pea (<i>Pisum</i> spp.) (dwarf, edible-pod, English, field, garden, green, snow, sugar snap)</p> <p>Pigeon pea</p> <p>Soybean (immature seed) (Also see Soybean section for specific use directions)</p> <p>Sword bean</p>			<p>Surface Application (broadcast or band): Apply to soil at planting in sufficient water or liquid fertilizer to provide uniform coverage. For banded application, a 7 inch band is recommended. Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation.</p>

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
LINGONBERRY	Root Rot (<i>Phytophthora</i> spp.)	7.2 or 0.5 per 1,000 row ft.	Surface Application (broadcast or band): Apply in sufficient water or liquid fertilizer to provide uniform coverage. For new plantings, make application at or after transplant. For banded application in new plantings, use an 18 inch band over the row. For established plants, use a 3 foot band over the row. Apply in the spring before plants start growth. If a second application is needed, time application to occur when root development is most likely to develop. Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation.
<ul style="list-style-type: none"> • This product may be applied on the day of harvest. • Do not soil-apply more than 14.4 pints per acre per crop (3.6 lbs. a.i./A). 			
ONIONS Including Dry Bulb Garlic Onions (dry bulb) Shallots (dry bulb). Green Green eschalots. Green onions Green shallots Japanese bunching onions Leeks Scallions Spring onions	Damping-off (<i>Pythium</i> spp.)	1 to 2	Preplant Incorporated: Apply as a broadcast soil application in sufficient water or liquid fertilizer to provide uniform coverage and mechanically incorporate in the top 2 inches of soil. For banded application, a 7 inch band is recommended. Surface Application: Apply at planting in sufficient water or liquid fertilizer to provide uniform coverage. For banded application, a 7 inch band is recommended. Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation.
<ul style="list-style-type: none"> • Dry bulb onions: Do not soil-apply more than 4 pints per acre (1 lb a.i./A) and foliar-apply more than 2 pints per acre (0.5 lb. a.i./A) per crop. • Green onions: Do not soil-apply more than 4 pints per acre (1 lb. a.i./A) and foliar-apply more than 1.2 pints per acre (0.3 lb. a.i./A) per crop. 			
PEANUTS	<i>Pythium</i> Root Rot Pod Rot (<i>Pythium</i> spp.)	0.5	Seedling Disease: Apply as an application to the seed in-furrow or in a 7 inch band at the time of planting. For the in-furrow applications, position the spray so the fungicide is mixed with the soil covering the seed. Avoid spraying the seed directly with the spray solution or crop injury may occur. Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation.
		1 to 2	Pod Rot: Apply at early pod set or pegging through overhead irrigation systems. See the General Information section of this label for further instructions and precautions when making applications through irrigation systems.

26
39

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
PINEAPPLE	Heart rot disease (<i>Phytophthora</i> spp.)	1 to 2	Crown Dip: Apply in 100 gallons of water as a crown dip before planting. The amount of dip solution per acre will depend upon crown size, planting density, and dipping techniques. For dipping, use 75 to 100 gallons of mixture per acre.
	<ul style="list-style-type: none"> • If there is a crop failure within 1 year of planting treated crowns, do not harvest plant material for animal feed. 		
POTATOES	Pythium leak (<i>Pythium</i> spp.)	0.84 fluid ounces per 1,000 row feet	<p>In-Furrow: Apply to row on a 6 to 8 inch band at planting in a minimum of 3 gallons of water per acre. Make application directly over the seed piece(s) prior to row closure. Use 11.6 fluid ounces per acre for 38 inch row spacing (13,756 linear feet per acre), 12.2 fluid ounces per acre for 36 inch row spacing (14,520 linear feet per acre), and 13 fluid ounces per acre for 34 inch row spacing (15,374 linear feet per acre). You may need to follow this in-furrow application with a foliar applied fungicide at tuber initiation:</p> <ul style="list-style-type: none"> • When conditions are conducive for disease development. • When variety planted is susceptible or moderately susceptible (pink rot/<i>Pythium</i> leak). • In areas with a long growing season. <p>This product may be impregnated on dry fertilizer or applied in combination with liquid fertilizers.</p>
	Pink rot (<i>Phytophthora erythroseptica</i>)		6.4 fluid ounces per acre.
	Seedling disease (<i>Pythium</i> spp.)	<ul style="list-style-type: none"> • Do not use the "dribble" application method. • Do not exceed 3/4 pints per acre per crop (0.188 lb a.i./A) of soil-applied mefenoxam and 1.6 pints per acre per crop (0.4 lb. a.i./A) of foliar-applied mefenoxam products per year. 	
RASPBERRIES	Phytophthora root rot (<i>Phytophthora</i> spp.)	0.5 per 1,000 row feet	Apply to the soil surface in a 3 foot band over the row. Make 1 application in the spring and another in the fall after harvest.
	<ul style="list-style-type: none"> • Do not apply within 45 days of harvest. 		

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
ROOT AND TUBER VEGETABLES Arracacha Arrowroot Artichoke (Chinese and Jerusalem) Beet, garden Beet, sugar (see Sugar Beet section) Burdock, edible Canna, edible Cassava (bitter and sweet) Carrots (see Carrot section below) Celeriac (celery root) Chayote (root) Chervil, turnip-rooted Chicory Chufa Dasheen (taro) Ginger Ginseng (see Ginseng section below) Horseradish Leren Parsley, turnip-rooted Parsnip Potato (see Potato section below) Radish Radish, oriental (daikon) Rutabaga Salsify (oyster plant, black, Spanish) Skirret Sweet potato Tanier (cocoyam) Turmeric Turnip Yam bean (jicama, manioc pea) Yam, true	<i>Pythium</i> Root Rot (<i>Pythium</i> spp.)	2 to 4	Preplant Incorporated: Apply as a broadcast soil application in sufficient water or liquid fertilizer to provide uniform coverage and incorporate in the top 2 inches of soil.
	Phytophthora Root Rot (<i>Phytophthora</i> spp.)		Surface Application: Apply at planting in sufficient water or liquid fertilizer to provide uniform coverage.
<ul style="list-style-type: none"> • For banded application, a 7 inch band is recommended. General Instructions section of this label to calculate the amount of this product needed per acre for the band width actually used. • Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation. 			

28
39

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
SOYBEANS	Phytophthora root and stem rot (<i>Phytophthora megasprema</i>)	0.16 to 0.56 fluid ounces per 1,000 row feet	In-Furrow: Apply in-furrow with water or liquid fertilizer at planting. Position the spray so the fungicide is mixed with the soil covering the seed. Avoid spraying the seed directly with the spray solution or crop injury may occur. Use the high rate for full season control and 0.16 to 0.3 fluid ounce rate for early to mid-season control.
	Pythium damping off (<i>Pythium</i> spp.)	0.74 to 2.5	Soil Spray: For full season control, apply 2.5 pints per treated acre in sufficient water or liquid fertilizer to provide uniform coverage at the time of plantings. For banded application, a 7 inch band is recommended. Use the formula in the General Instructions section of this label to calculate the amount of this product needed per acre for the band width actually used. For early to mid season control, apply 0.74 to 1.5 pints per acre. Move product into the seed zone after planting with 0.5 to 1 inch sprinkler irrigation.
<ul style="list-style-type: none"> • Under heavy late season Phytophthora pressure, this product may not provide complete control. • Use the higher rate in areas with a history of heavy Phytophthora damage. • For best results, use soybean varieties that have some degree of resistance to the races of Phytophthora present in the field. 			

29
39

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
SPINACH	White rust (<i>Albugo occidentalis</i>) Downy mildew (<i>Peronospora effuse</i>)	0.5	<p>Shank Application: In addition to the preplant incorporated or surface application described above, shank-in this product around 21 days after planting or after the first cutting. A second application may be shanked in after the next cutting, but at least 21 days after the first application. If less than the full bed is treated, use the formula in the GENERAL INSTRUCTIONS section to determine the amount of this product needed per acre. If this product is injected into the beds with liquid fertilizer, base calculations on a 7 inch band. Use sufficient mechanical or bypass agitation to keep this product mixed with the water or fertilizer.</p> <p>The additional applications of this product noted above and made after each cutting by shanking with fungicide into the beds along with liquid fertilizer provide continuing control of white rust. However white rust can only be controlled in a preventative disease control program that begins with an application of this product to the soil at planting. If this product is not used at planting, do not use this product at any other time throughout the season. Do not apply this product in foliar applications or in situations where white rust infections are already established. The use of this product in curative applications greatly increases the risk of the fungus developing insensitivity to mefenoxam. The development of insensitivity will destroy the effectiveness of this product in controlling white rust.</p>
<ul style="list-style-type: none"> • Do not harvest spinach within 21 days of application. • In spinach, do not apply more than 4 pints per acre to soil at planting (1 lb. a.i./A) and either apply no more than 1 pint per acre (0.25 lb a.i./A) to soil post-planting, shanked-in or 1.6 pints per acre (0.4 lb a.i./A) foliar-applied per crop. 			

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
STONE FRUITS Including Apricot Cherries (sweet and tart) Nectarine Peach Plum (Chickasaw, Damson, Japanese) Plumcot Prune (fresh) Hybrids or cultivars of these	Crown, Collar and Root rots (<i>Phytophthora</i> spp.)	8 or 3 fluid ounces per 1,000 square feet	Apply to soil in sufficient water to obtain thorough coverage of the soil under the canopy of the trees. Sufficient surface area should be treated to cover the root zone of the plants 2 weeks after planting (new plantings) or in the spring before plant growth (established plantings). Additional applications should be made at 2 to 3 month intervals depending on disease pressure. Up to 3 applications can be made per year. For intense plantings (2 to 3 times the normal planting rate) apply on a per area basis (1,000 square feet). Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation.
<ul style="list-style-type: none"> • Apply before symptoms appear. This product will not revitalize trees showing moderate to severe disease symptoms. • Do not graze livestock in treated areas. • Do not graze or feed cover crops in treated orchards. • Do not concentrate the treatment around the tree trunks or injury may occur. • Do not apply to trees under stress. • In CA, do not apply this product newly planted trees within 45 days of planting. Some varieties may exhibit chlorosis on leaf margins. • Do not soil-apply more than 24 pints (3 gallons) per acre per crop (6 lb. a.i./A). 			
STRAWBERRIES	Red stele (<i>Phytophthora fragariae</i>) Vascular collapse (<i>P. cactorum</i>) Leather rot (<i>P. cactorum</i>)	2	New Plantings: Apply in banded sprays or through drip irrigation in sufficient water to move the fungicide into the root zone of the plants. Make first application after transplanting. Make second application 30 days before the beginning of harvest or at fruit set. Apply the third application during the harvest season, depending on environmental conditions and disease pressure. Established Plantings: Apply in banded sprays or through drip irrigation in sufficient water to move the fungicide into the root zone of the plants. Make first application in the spring after the ground thaws and before first bloom. Make final application after harvest in the fall. For supplemental control of leather rot, make another application during the growing season at fruit set.
<ul style="list-style-type: none"> • For drip irrigation, calculate rates as a band application with a band width equal to the root zone width and inject this product into the irrigation water. • Application of this product may be made on the day of harvest. • Do not soil-apply more than 6 pints per acre per crop (1.5 lb. a.i./A). • For low annual rainfall areas, a surface application of this product needs to be moved into the root zone by rainfall, overhead irrigation, or mechanical incorporation. 			

31
39

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
SUGAR BEETS	<i>Pythium</i> Root Rot (<i>Pythium</i> spp.)	2 to 4	<p>Preplant Incorporated: Apply as a broadcast soil application in sufficient water or liquid fertilizer and mechanically incorporate in the top two inches of soil. For banded application, a 7 inch band is recommended. Use the formula in the General Instructions section of this label to calculate the amount of this product needed per acre for the band width actually used.</p> <p>Surface Application: Apply at planting in sufficient water or liquid fertilizer to provide uniform coverage. For banded application, a 7 inch band is recommended. Use the formula in the General Instructions section of this label to calculate the amount of this product needed per acre for the band width actually used. Soil surface sprays of this product will not be effective until, the fungicide is moved into the root zone by rainfall or irrigation.</p>
TOBACCO	Damping off (<i>Pythium</i> spp.)	1 to 2 or 0.5 to 1 fl. oz. per 150 sq. yds.	Soil spray (broadcast): Apply in 50 gallons of water (2 gallons water per 150 square yards) as a preplant soil application before or at time of planting. Use the higher application rate on broadleaf tobacco. Move the product into the seed zone after planting as recommended in the Application Instructions.

32
39

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
TOBACCO (Continued)	Black Shank <i>(Phytophthora parastica var. nicotiana)</i>	<p>For conventional till: 2 to 6</p> <p>For no-till tobacco: 1 to 2</p>	<p>Pre-plant/Pre-transplant Soil spray (broadcast or band): Apply in sufficient water or liquid fertilizer to provide uniform coverage at the time of planting. If rainfall is not expected before the seeds start to germinate, this product should be incorporated mechanically before planting, during the planting operation, or be moved into the seed zone after planting with 1/2 to 1 inch sprinkler irrigation.</p> <p>For banded applications, a 7 inch band is recommended. Use the formula in the General Instructions section of this label to calculate the amount of this product needed per acre for the band width actually used.</p> <p>Post-plant/Post-transplant Soil spray (broadcast): Apply to the soil within one week of planting. Incorporate in the top 2 to 4 inches of soil. Use the high rate if the disease epidemic is expected to be severe. In FL and GA, use 6 pints per acre where black shank is severe. Move into soil as recommended in the Application Instructions. Apply preventively for effective black shank control. If black shank is expected early in the season, apply as near as possible to transplanting followed by sequential applications.</p>
	Blue mold <i>(Peronospora tabacina)</i>	1 to 2	<p>Pre-plant/Pre-transplant Soil Spray (broadcast): Apply as a broadcast soil application prior to transplanting and incorporate in the top 2 to 4 inches of soil before forming beds. Under low disease pressure or for early season control, use the low rate. Use the high rate for high disease pressure, extended control and burley and other tobacco types other than flue-cured.</p> <p>Post-plant/Post-transplant Soil spray (band): If a pre-transplant application was made, make a supplemental application at lay-by or the last cultivation. Position the nozzles so that the spray is deposited under the plants and is covered by soil by the cultivator. Move into soil as recommended in Application Instructions.</p>

33
 39

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
TOBACCO (Continued)	<ul style="list-style-type: none"> • Consult local extension bulletins for additional use directions. • For best results against black shank, use this product with tobacco varieties that have high resistance to black shank and use crop rotation. • In fields where there is a history of severe black shank, use the highest rate and plant a variety that is resistant to the race of <i>Phytophthora</i> present in the field. (Burley L8 hybrids are only resistant to <i>Phytophthora</i> Race 0. • Do not use in high black shank areas on highly susceptible flue-cured varieties. • Failure to adequately control nematodes in fields treated with this product may result in poor control of black shank. • Do not use this product for black shank control in PA. • Do not exceed the equivalent of 1.5 lb A/A per crop of mefenoxam containing products. 		
TOMATO	Damping-off (<i>Pythium</i> spp.)	2 to 4	Soil Spray (Broadcast or Band): Apply at planting in water or liquid fertilizer. Move into soil as recommended in the General Application Instructions section. For banded applications, a 7 inch band is recommended.
	Root and Fruit Rot (<i>Phytophthora</i> spp.) (<i>Pythium</i> spp.)	2	Soil spray (broadcast or band) or soil injection: Apply as a directed soil surface spray under the vines or injected into the beds with water or liquid fertilizer. Move into the soil as recommended in the General Application Instructions. Make application 4 to 6 weeks after planting. If needed, make a second application up to 4 weeks before harvest, but before the last irrigation. Injection (drip irrigation): Initiate control of Root and Fruit Rot with a soil application as described above. Make subsequent applications through drip irrigation. Make the first drip application 4 to 6 weeks after planting. Apply the second rip application as needed up to 4 weeks before harvest, but before the last irrigation. For injection applications, base rate calculation on a 7 inch band.
	<ul style="list-style-type: none"> • Do not exceed 6 pints per acre per crop (1.5 lb. a.i./A) of soil-applied mefenoxam containing products and 2 pints per acre per crop (0.5 lb. a.i./A) of foliar-applied mefenoxam products per year. 		

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
TREE NUTS Almonds Walnuts	Crown, Collar and Root rots <i>(Phytophthora spp.)</i>	8 or 3 fluid ounces per 1,000 square feet	<p>Soil spray (broadcast or band): Apply under the tree canopy to cover the root zone two weeks after planting (new plantings) or in the spring before growth begins (established plantings). Move into soil as recommended in General Application Instructions.</p> <p>Additional applications may be made at 2 to 3 month intervals depending on disease pressure. Make up to 3 applications per year.</p> <p>Apply before symptoms appear. This product will not revitalize trees showing moderate to severe disease symptoms.</p> <p>For intense plantings (2 to 3 times the normal planting rate), apply on a per area basis (1,000 square feet).</p> <ul style="list-style-type: none"> • Apply before symptoms appear. This product will not revitalize trees showing moderate to severe disease symptoms. • Do not graze livestock in treated areas. • Do not graze or feed cover crops in treated orchards. • Do not concentrate the treatment around the tree trunks or injury may occur. • Do not apply to trees under stress. • In CA, do not apply this product newly planted trees within 45 days of planting. Some varieties may exhibit chlorosis on leaf margins. • Do not soil-apply more than 24 pints (3 gallons) per acre per crop (6 lb. a.i./A) of mfenoxam containing products.
TROPICAL FRUIT Kiwifruit	Root and Crown rot <i>(Pythium and Phytophthora spp.)</i>	11.2 to 22.4 fluid ounces	<p>Apply 1 quart of Ultra Flourish solution as a soil drench in a one square foot area around the base of each vine. At the recommended rate of 11.2 to 22.4 fluid ounces, this will apply 0.175 to 0.350 lb. a.i./A if the planting density is 160 vines per acre.</p> <p>Make the first application in the fall after harvest or in February or early March. Make a second application in the spring or approximately 60 days after the February or March application. Move into soil as recommended in the General Application Instructions.</p> <ul style="list-style-type: none"> • Do not apply more than 1.4 pints per acre (0.7 lb. a.i. /A) per growing season. • Do not apply within 200 days of harvest.

CROP	DISEASES	RATE (Pints per Acre)	USE INSTRUCTIONS
TROPICAL FRUIT Black Sapote Canistel Mamey Sapote Mango Papaya Sapodilla Star Apple	Root rot <i>(Phytophthora spp.)</i> Damping-off <i>(Pythium spp.)</i>	3 to 6	Soil Drench: Apply to the soil surface in sufficient water or liquid fertilizer to obtain thorough coverage of the soil under the canopy of trees. To determine the amount of solution required, measure the amount of water needed to drench an area of 1 square foot around one plant. Multiply that volume by the total number of plants in an acre. Add this product to this amount of water and drench plants. Two applications may be made per growing season. Make the first application at transplanting or in the spring at root growth flush. Apply the second application up to 30 days prior to harvest.
<ul style="list-style-type: none"> • Do not apply within 30 days of harvest. • Do not use this product if a metalaxyl-copper product is used as a trunk and foliage spray. 			
TROPICAL FRUIT Atemoya Chemimoya Custard Apple Ilama Soursop Birida Sugar Apple	Root rot <i>(Phytophthora spp.)</i> Damping-off <i>(Pythium spp.)</i>	3 to 6	Soil Drench: Apply to the soil surface spray in sufficient water or liquid fertilizer to obtain thorough coverage of the soil under the canopy of trees. Make one application in the spring at root growth flush and a second application in the fall.
<ul style="list-style-type: none"> • Do not apply within 30 days of harvest. 			
TROPICAL FRUIT Starfruit	<i>Phytophthora</i> Root and Crown rot <i>(Phytophthora spp.)</i> <i>Pythium</i> Root and Crown rot <i>(Pythium spp.)</i>	3 to 6	Soil Drench: Apply in sufficient water or liquid fertilizer to provide an adequate soil drench. Make one application in the spring when root growth begins and a second application in the fall. Direct applications to the soil surface under the canopy of the trees.
<ul style="list-style-type: none"> • Do not apply within 30 days of harvest. 			

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store at temperatures above 40°F. Crystals may form at lower storage temperatures. If this occurs, place the product in a warm room (68°F or above) and roll or shake the container at frequent intervals until all crystals are dissolved.

PESTICIDE DISPOSAL: Open dumping is prohibited. Wastes resulting from the use of this product are acutely toxic. Improper disposal of unused pesticide, spray mixture or rinsate is a violation of Federal law. Pesticide, spray mixture or rinsate that cannot be used according to the label instruction must be disposed of according to Federal, state or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

CONTAINER DISPOSAL:

Nonrefillable Containers 5 Gallons or Less: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty

the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

Nonrefillable containers larger than 5 gallons: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers larger than 5 gallons: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV071609)

Ultra Flourish is a registered trademark of Nufarm Americas, Inc.
All other trademarks are the property of their respective owners.

37
39

[BASE LABEL]

ULTRA FLOURISH®

AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENT:

Mefenoxam: D-Alanine, N-(2,6-dimethylphenyl)-N-(methoxyacetyl)-, methyl ester 25.1%

OTHER INGREDIENTS: 74.9%

TOTAL: 100.0%

Contains Petroleum Distillates

Contains 2 pounds mefenoxam per gallon

**KEEP OUT OF REACH OF CHILDREN
WARNING / AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

REFER TO BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE

**For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840**

FIRST AID

IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

NOTE TO PHYSICIAN

If this product is ingested, lavage stomach. A slurry of activated charcoal in water can be left in the stomach. Contains petroleum distillate – vomiting may cause aspiration pneumonia. See Label for Additional Precautions and Directions for Use.

**PRECAUTIONARY STATEMENT
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING - AVISO**

Causes substantial but temporary eye injury. Harmful if swallowed or absorbed through skin. Wear protective eyewear such as goggles or face shield. Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

38
39

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long -sleeved shirt and long pants
- Chemical-resistant gloves such as barrier laminate or viton
- Shoes plus socks
- Protective eyewear

See engineering controls for additional requirements.

USER SAFETY REQUIREMENTS

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statement:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Apply only as specified on this label. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store at temperatures above 40°F. Crystals may form at lower storage temperatures. If this occurs, place the product in a warm room (68°F or above) and roll or shake the container at frequent intervals until all crystals are dissolved.

PESTICIDE DISPOSAL: Open dumping is prohibited. Wastes resulting from the use of this product are acutely toxic. Improper disposal of unused pesticide, spray mixture or rinsate is a violation of Federal law. Pesticide, spray mixture or rinsate that cannot be used according to the label instruction must be disposed of according to Federal, state or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

CONTAINER DISPOSAL:

[Nonrefillable Containers 5 Gallons or Less: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.]

39
39

[Nonrefillable containers larger than 5 gallons: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.]

[Refillable containers larger than 5 gallons: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.]

EPA Reg. No. 55146-73
EPA Est. No.

Manufactured for:
Nufarm Americas Inc.
AGT Division
150 Harvester Drive
Burr Ridge, IL 60527

NET CONTENTS GALLONS