

55146-75

11/29/1999

1/13

Please read instructions on reverse before completing form.

NOTIFICATION

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

☐ Registration
☐ Amendment
☒ Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 55146-75	2. EPA Product Manager Waller	3. Proposed Classification <input type="checkbox"/> None <input checked="" type="checkbox"/> Restricted
4. Company/Product (Name) Micro Sulf	PM# 21	
5. Name and Address of Applicant (Include ZIP Code) Agtrol International 7322 Southwest Freeway, #1400 Houston, TX 77049 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION

NOV 29 1999

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

To add the crop, peppers. This crop was omitted by printing error from our currently accepted label but was previously accepted.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	
				<input type="checkbox"/> Plastic	
				<input type="checkbox"/> Glass	
				<input type="checkbox"/> Paper	
				<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Nancy R. Robey	Title Regulatory Specialist	Telephone No. (Include Area Code) (713) 995-0111
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Regulatory Specialist	
4. Typed Name Nancy R. Robey	5. Date November 12, 1999	



NOTIFICATION

NOV 29 1999

Micronized Wettable Sulfur

Fungicide - Miticide

For Use on Fruits - Nuts - Citrus - Vegetables - Hops

Sugar Beets - Cotton - Seed Alfalfa - Cereal Grains

Peanuts - Roses

ACTIVE INGREDIENT:

Sulfur..... 80%

INERT INGREDIENTS:..... 20%

Total..... 100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Give one or two glasses of water and induce vomiting by sticking finger down throat. Never give anything by mouth to an unconscious person.

IF ON SKIN: Wash with soap and water. Get medical attention if irritation persists.

IF IN EYES: Immediately flush eyes with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 55146-75

EPA Est. No. 44420-FR-1



PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

May cause irritation of eyes. Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes, or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants waterproof gloves, shoes plus socks, and protective eyewear. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE INSTRUCTIONS: Store product in a closed package in a cool, dry place. Do not store near flammable materials. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling, sweep up spillage and dispose of in accordance with the Pesticide Disposal Instructions below.

PESTICIDE DISPOSAL INSTRUCTIONS: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL INSTRUCTIONS: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment, then dispose of bag in a sanitary landfill, by incineration, or if allowed by State and local authorities, by burning. If burned stay out of smoke.



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

OBSERVE SPECIFIC INSTRUCTIONS FOR USE IN CALIFORNIA

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling, and with the Worker Protection Standard (WPS), 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about PPE, notification to workers, and restricted-entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the WPS.

Do not enter or allow worker entry into treated areas during the REI of 24 hours. PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, shoes plus socks, and protective eyewear.

CHEMICAL OR PHYSICAL HAZARD

Keep away from heat, sparks, or flame. Do not smoke while applying this product.

IMPORTANT

In order to obtain effective mite and disease control, both the upper & lower sides of plant foliage and fruit must be thoroughly covered with Micro Sulf™ spray solution. Poor spray coverage will result in poor mite and disease control. When growing crops for "processing", consult the processor before applying sulfur. Do not use on any crop unless sulfur has been shown to be safe for that crop in your locality. Sulfur may burn foliage when temperatures exceed 90°F. Do not apply within 14 days of an oil spray for all crops except citrus. For citrus do not apply within 21 days of an oil spray.

APPLICATION INSTRUCTIONS

Apply through any type of handheld, knapsack, mechanical or pneumatic spraying equipment.



MIXING: Pour the required amount of MICRO SULF into the indicated amount of water while stirring. MICRO SULF disperses instantly.

Dilute Application:

Field: Specified rate in 20 to 60 gallons of water per acre.

Orchard: Specified rate in 100 to 800 gallons of water per acre.

Concentrated Application:

Field: Specified rate in 5 to 10 gallons of water per acre.

Orchard: Specified rate in 20 to 100 gallons of water per acre.

Aerial Application:

Field: Specified rate in 3 to 20 gallons of water per acre.

Orchard: Specified rate in 10 to 20 gallons of water per acre.

COMPATIBILITY

MICRO SULF is completely compatible with Bordeaux mixture, copper fungicides, liquid fertilizers and all organic fungicides, insecticides, acaricides and herbicides.

USE INSTRUCTIONS

Some crops may be damaged by sulfur under certain climatic conditions, such as when temperature is high. AGTROL INTERNATIONAL does not recommend application if temperature will exceed 90°F within the three days following spraying, due to the risk of crop injury. Crops grown in greenhouses may be much more sensitive to sulfur injury, so the lowest labeled rate (or below labeled rates) should be tried initially. Do not use on any crop unless sulfur has been shown to be safe in your locality. Do not allow spray to drift into sulfur-sensitive crops such as apricots, cranberries, spinach and d'Anjou pears.

Do not use within two weeks of an oil spray treatment, except for Dormant, Delayed Dormant, or Postharvest applications with a suitable oil. For citrus, do not apply within 21 days of an oil spray. Spreaders/stickers are not required for use with MICRO SULF. If spreaders/stickers are used with MICRO SULF, foliar injury may be enhanced under high temperatures. When growing crops for processing, consult the processor before applying sulfur.

NOTE: The application rates specified for each crop are the maximum that should be applied. These amounts may be reduced if experience or local practice indicate that lesser amounts may be effective. Consult State Agriculture Experiment Station or State Extension Service Specialist for rates recommended in your local area.



VEGETABLE CROPS

BEANS

Leaf Spot, Powdery Mildew, Red Spider Mites: apply at a rate of 7 pounds per acre. Apply at early leaf stage and repeat at 14 day intervals or as needed.

BROCCOLI, BRUSSELS SPROUT, CABBAGE, CAULIFLOWER

Powdery Mildew, Red Spider Mite, Septoria Leaf Spot: apply at a rate of 3 to 10 pounds per acre. Begin in early leaf stage. Repeat every 10 to 14 days or as needed.

CARROTS*

Powdery Mildew, Petrobia Mite: apply at a rate of 3 to 10 pounds per acre. Apply at early leaf stage and repeat every 14 days or as needed.

***Not For This Use In California**

CUCUMBERS

Powdery Mildew: apply at a rate of 2 to 4 pounds per acre. Apply when disease first appears and repeat as necessary. Do not apply more than 2 pounds MICRO SULF per acre per application during periods of high temperatures.

GREENS

(COLLARDS, KALE, MUSTARD*, TURNIP*)

Powdery Mildew, Red Spider Mite, Septoria Leaf Spot: apply at a rate of 3 to 10 pounds per acre. Apply at early leaf stage and repeat every 14 days or as needed.

***Not For This Use In California**

LETTUCE* (HEAD AND LEAF), *Powdery Mildew, Rust, Red Spider Mite.* Apply at a rate of 5-6 pounds per acre. Apply at early leaf stage and repeat every 14 days or as needed. Thorough coverage is required.

***Not For This Use In California**

ONIONS*, GARLIC*, DRY ONIONS*, DRY SHALLOTS*

Powdery Mildew, Petrobia Mite: apply at a rate of 3 to 10 pounds per acre. Apply when disease first appears and repeat as needed.

***Not For This Use In California**

PEAS

Powdery Mildew, Septoria Leaf Spot, Red Spider Mite: apply at a rate of 3 to 5 pounds per acre. Start application at early leaf stage. Repeat application every 10-14 days or immediately after rain.



PEPPERS

Powdery Mildew, Broad Mite, Red Spider Mite:* apply at a rate of 3 to 10 pounds per acre. Apply at early leaf stage and repeat every 14 days or as needed.

***Not For This Use In California**

POTATOES

Powdery Mildew, Red Spider Mite, Septoria Leaf Spot: apply at a rate of 5 pounds per acre. Apply at early leaf stage and repeat every 14 days or as needed.

RUTABAGAS

Powdery Mildew, Red Spider Mite:* apply at a rate of 3 to 10 pounds per acre. Apply at early leaf stage and repeat every 14 days or as needed.

***Not For This Use In California**

SQUASH

Powdery Mildew: apply at a rate of 5 to 10 pounds per acre. Apply when disease first appears and repeat as needed. Sulfur can injure plants, especially when temperature reaches 95°F. Do not use on sulfur sensitive varieties.

TOMATOES

Two-Spotted Mite, Russet Mite, Powdery Mildew:* apply at a rate of 5 to 10 pounds per acre. Begin application when symptoms first appear. Thorough coverage is required.

***Not For This Use In California**

FRUIT AND NUT CROPS**ALMONDS**

Brown Rot, Blossom Blight, Powdery Mildew, Leaf Spot, Silver Mite, Flat Mite, Red Spider Mite: apply at a rate of 10 to 20 pounds per acre. Apply at bloom or early petal fall. Repeat as necessary, usually 10 to 14 days after a period of wet weather.

APPLES, PEARS

Powdery Mildew, Scab, Blister & Silver Mites, Two-Spotted Mite, Red Spider Mite:* apply at a rate of 10 to 20 pounds per acre. Apply at pre-bloom and petal fall. Apply with cover sprays throughout the season. Do not apply to d'Anjou pears.

***Not For This Use In California**

APPLES, PEARS (DORMANT, DELAYED DORMANT, OR POSTHARVEST)

Pear Psylla, Blister Mite, Two-Spotted Mite, Red Spider Mite:* apply at a rate of 10 to 20 pounds per acre. Apply with a suitable spray oil postharvest and during the dormant periods of bud development. Do not apply to sensitive varieties of pears except at these stages.



Not For This Use In California*AVOCADOS**

Brown Mite: apply at a rate of 10 to 20 pounds per acre. Apply as needed.

CITRUS

*Broad Mite**, *Citrus Rust Mite**, *Flat Mite*, *Silver Mite*, *Bud Mite**, *Red Spider Mite*, *Six Spotted Mite**, *Thrips**, *Yuma Spider Mite**: apply at a rate of 10 to 15 pounds per acre. Apply as needed. Do not apply within 21 days of an oil spray.

Not For This Use In California*FIGS**

Mites, including Fig Mite, Eriophyid Mite: apply at a rate of 5 pounds per acre. Apply at a minimum of 100 gallons per acre. Begin applications in April, and continue on a 3 to 4 week schedule through June.

GRAPES

Powdery Mildew, Bud Mite, Blister Mite, Red Spider Mite, Phomopsis: apply at a rate of 3 to 10 pounds per acre. Apply throughout the season at 7 to 14 day intervals or as needed. Thorough coverage is required.

NOTE: Use with caution on Concord or Labrusca type grapes as injury can occur after bloom.

MANGOS*

Powdery Mildew: apply at a rate of 10 to 30 pounds per acre. Begin application before bloom. Repeat at bloom, after fruit set, and three weeks later or as needed.

Not For This Use In California*PECANS**

Powdery Mildew, Sooty Mold, Two-Spotted Mite, Red Spider Mite, Eriophyid Mite (including Pecan and Hickory): apply at a rate of 5 to 10 pounds per acre. Apply throughout the season as needed.

NOTE: Some varieties of pecans are sensitive to sulfur sprays under certain conditions. Do not apply unless varieties are known to be tolerant of sulfur.

PISTACHIOS*

Mites: apply at a rate of 10 to 20 pounds per acre. Begin applications in June-July. Make additional applications as necessary. When temperatures exceed 95°F, lower rates and more frequent applications are advised in order to avoid crop injury.

***Not For This Use In California**

POMEGRANATES*

Mites: apply at a rate of 3 to 10 pounds per acre. Begin applications in May or June. Make additional applications on a 3 to 4 week schedule, or as necessary. Use higher rates if past mite damage has been high. If temperatures exceed 95°F, lower rates and more frequent applications advised in order to avoid crop injury.

***Not For This Use In California**

STONE FRUITS**(PEACHES, PLUMS, CHERRIES, NECTARINES, PRUNES)**

Powdery Mildew, Blossom Blight, Brown Rot, Leaf Spot, Silver Mite, Flat Mite Red Spider Mite: apply at a rate of 10 to 20 pounds per acre. Follow directions for almonds, above. For Brown Rot of fruit, apply when fruit starts to ripen. Do not apply to apricots.

WALNUTS*, MACADAMIAS*

Pacific Mite, Almond Mite, Two-Spotted Mite, Red Spider Mite, Broad Mite: apply at a rate of 10 to 20 pounds per acre. Apply throughout the season as needed.

***Not For This Use In California**

NOTE: Some varieties of walnuts are sensitive to sulfur sprays under certain conditions. Do not apply unless varieties are known to be tolerant of sulfur.

FIELD CROPS**SEED ALFALFA**

Atlantic Mite, Red Spider Mite*: apply at a rate of 8 to 10 pounds per acre. Apply throughout the season as necessary.

***Not For Use In California**

CEREALS**(CORN, WHEAT, BARLEY, OATS, RYE, SORGHUM)**

Red Spider Mite, Two-Spotted Mite, Pacific Mite, Atlantic Mite: apply at a rate of 8 to 10 pounds per acre. Apply when symptoms first appear and repeat applications as necessary.

COTTON

Atlantic Mite, Two-Spotted Mite*, Red Spider Mite*: apply at a rate of 5 to 10 pounds per acre. Apply throughout the season as necessary.

***Not For This Use In California**

PEANUTS

Powdery Mildew, Leaf Spot, Two-Spotted Mite*, Red Spider Mite, Rust**: apply at a rate of 5 to 8 pounds per acre. Apply at early leaf stage and repeat application as necessary.



Not For This Use In California*SOYBEANS**

Leaf Spot, Powdery Mildew:* apply at a rate of 10 to 15 pounds per acre. Apply at early leaf stage and repeat at 14 day intervals or as needed.

Not For This Use In California*SUGAR BEETS**

Powdery Mildew, Red Spider Mite: apply at a rate of 5 to 10 pounds per acre. Apply as soon as symptoms appear and repeat every 2 to 3 weeks or as needed.

SMALL FRUITS AND BERRIES**BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, BLUEBERRIES, GOOSEBERRIES, HUCKLEBERRIES, CURRANTS**

Powdery Mildew: apply at a rate of 6 to 15 pounds per acre. Begin application before blossoms open and continue at 10 day intervals or as needed.

RASPBERRIES

Powdery Mildew, Red Berry Mites: Apply at a rate of 6 to 15 pounds per acre. Apply each week from first bloom to fruit set. Repeat as the weather requires.

STRAWBERRIES

Powdery Mildew, Two-Spotted Mite, Red Spider Mite:* apply at a rate of 5 to 10 pounds per acre. Begin application at early leaf stage and repeat as needed. Do not use on sulfur-sensitive varieties.

Not For This Use In California*MISCELLANEOUS CROPS****MELONS**

Powdery Mildew: apply at a rate of 5 to 10 pounds per acre. Apply when disease first appears and repeat as needed. Sulfur can injure plants, especially when temperature reaches 95°F. Do not use on sulfur sensitive varieties.

SPEARMINT, PEPPERMINT

Powdery Mildew: apply at a rate of 4 to 6 pounds per acre. Apply when mint is 5" - 6" tall or when disease appears. Repeat twice, at 30 day intervals. Do not apply within 30 days of harvest.

HOPS*

Red Spider Mite: apply at a rate of 2 to 6 pounds per acre. Begin applications at early leaf stage, and repeat as needed.

***Not For This Use In California**

ORNAMENTALS, ROSES

*Powdery Mildew, Leaf Spot, Two-Spotted Mite**: apply at a rate of 3 to 10 pounds per acre by ground or 3-20 pounds per acre by air. Apply to ensure complete plant coverage. Begin when disease first appears and repeat at 5-10 day intervals.

***Not For This Use In California**

DIRECTIONS FOR USE THROUGH CHEMIGATION SYSTEMS

NOT FOR THIS USE IN CALIFORNIA

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Observe all applicable precautions and limitations on the Federal label. Label must be in the possession of the user at the time of pesticide application.

GENERAL INSTRUCTIONS

Apply this product through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have any questions about calibration, you should contact your State Extension Service Specialists, equipment manufacturers or other experts.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Do not connect the chemigation system to any public water system. Public water system means a system for the provision of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

MICRO SULF may be applied in combination with chemically neutral liquid fertilizers. Application in conjunction with highly alkaline fertilizers, such as aqueous ammonia, may cause a degradation of the pesticide, resulting in reduced performance, and should be avoided.

SYSTEM REQUIREMENTS

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.



The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functioning interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

APPLICATION INSTRUCTIONS

Observe the requirements in the System Requirements section above.

Apply MICRO SULF only through systems containing anti-syphon and check valves designed to prevent water source contamination or overflow of the mix tank and containing interlocking controls between the metering device and the water pump to insure simultaneous shutoff.

Maintain a gentle continuous agitation in mix tank during mixing and application to assure a uniform suspension.

Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute suspension per unit time.

Application of more than recommended quantities of irrigation water per acre may result in decreased product performance.

Do not apply when wind speed favors drift, when system connection or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product cannot be flushed and must be dismantled and drained. In a center pivot system, block the nozzle set nearest the well/pivot/injection unit to prevent spray being applied to this area.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water.

SPRAY PREPARATION

Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system. Flush with clean water. Prepare a suspension of MICRO SULF in a mix tank. Fill the tank



with 1/2 or 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Slowly add the required amount of MICRO SULF and then the remaining volume of water.

SPRINKLER IRRIGATION NOTES:

Observe all System Requirements and Application Instructions above.

Set sprinkler system to deliver 0.1 to 1.25 inches of water per acre. Volumes of water higher than this may reduce product efficacy. Start sprinkler and then uniformly inject the suspension of MICRO SULF into the irrigation waterline so as to deliver the desired rate per acre. The suspension of MICRO SULF should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing. When treatment with MICRO SULF has been completed, do not irrigate the treated area for 24 to 48 hours to prevent washing the chemical off the crop.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Where sprinkler distributed patterns do not overlap sufficiently, unacceptable pest control may result.

Check local restriction and requirements regarding sprinkler irrigation applications, as they may vary from state to state.

Micro Sulf is a trademark of Agtrol International.

© 1999 Agtrol International.

