

55146-107

8/2/2012

1/12



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF  
CHEMICAL SAFETY AND  
POLLUTION PREVENTION

Nathan P Ehresman  
Director Regulatory Affairs  
Nufarm Americas Inc - AGT Division  
% Nufarm Americas Inc  
4020 Aerial Center Parkway Suite 101  
Morrisville NC 27560

AUG - 2 2012

Product Name SEBRING 318 FS  
EPA Reg No 55146 107  
Subject Your amendment dated April 27 2012  
OPP Decision Number 463987

Dear Mr Ehresman

The amended label referred to above submitted in connection with registration under the Federal Insecticide Fungicide and Rodenticide Act as amended is acceptable provided the following change is made

- 1 On page 3 delete the word GENERAL from the header GENERAL PRECAUTIONS AND RESTRICTIONS

One copy of the label stamped Accepted with comments is enclosed for your records This label supersedes all labels previously accepted for this product Please submit one copy of the final printed label before the product is released for shipment

If you have any questions please contact Robert Westin by phone at (703) 305 5721 or via email at westin.robert@epa.gov

Sincerely

A handwritten signature in cursive script that reads "Mary L Waller".

Mary L Waller  
Product Manager (21)  
Fungicide Branch  
Registration Division (7504P)

Enclosure

# SEBRING™ 318 FS FUNGICIDE

FOR USE AS A SEED TREATMENT FOR CONTROL OF SYSTEMIC DOWNY MILDEW,  
PYTHIUM SEED ROT PYTHIUM DAMPING-OFF, AND EARLY SEASON PHYTOPHTHORA  
DISEASES OF CERTAIN CROPS

**ACTIVE INGREDIENT**

Metalaxyl N (2,6-dimethylphenyl) N-(methoxyacetyl) alanine methyl ester 30.14%

**OTHER INGREDIENTS**

**TOTAL**

69.86%  
100.00%

Contains 2.65 pounds Metalaxyl per gallon (318 grams per liter)

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

**SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS**

For Chemical Spill, Leak, Fire or Exposure, Call CHEMTREC (800) 424 9300  
For Medical Emergencies Only Call (877) 325 1840

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:**

8/2/2012

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended for the pesticides  
registered under EPA Reg. No.**

55146-107

EPA REG NO 55146 107  
EPA EST NO \_\_\_\_\_

Manufactured For  
NUFARM AMERICAS INC  
AGT Division  
150 Harvester Drive  
Burr Ridge IL 60527



NET CONTENTS \_\_\_\_\_ GAL (\_\_\_\_ Liters)  
[Designation as NONREFILLABLE or REFILLABLE for containers > 5 GAL]

<b>FIRST AID</b>	
<b>IF SWALLOWED</b>	Call a poison control center or doctor immediately for treatment advice Have person sip a glass of water if able to swallow <b>DO NOT</b> induce vomiting unless told to do so by the poison control center or doctor <b>DO NOT</b> give anything by mouth to an unconscious person
<b>IF IN EYES</b>	Hold eye open and rinse slowly and gently with water for 15 to 20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing eye Call a poison control center or doctor for treatment advice
<b>IF ON SKIN OR CLOTHING</b>	Take off contaminated clothing Rinse skin immediately with plenty of water for 15 to 20 minutes Call a poison control center or doctor for treatment advice
<b>IF INHALED</b>	Move person to fresh air If person is not breathing call 911 or an ambulance then give artificial respiration preferably by mouth to mouth if possible Call a poison control center or doctor for further treatment advice
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor or going for treatment You may also contact 1 877 325 1840 for emergency medical treatment information	

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed Causes moderate eye irritation Avoid contact with eyes skin or clothing Wash thoroughly with soap and water after handling Remove contaminated clothing and wash before reuse

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical resistant to this product are listed below If you want more options follow the instructions for category A on an EPA chemical resistance selection chart

**Applicators and other handlers must wear**

- Long sleeved shirt and long pants
- Shoes and socks
- Chemical resistant gloves such as barrier laminate or nitrile rubber or neoprene rubber or Viton

Follow manufacturer's instructions for cleaning/maintaining PPE If no such instructions for washables exist use detergent and hot water Keep and wash PPE separately from other laundry

<b>USER SAFETY RECOMMENDATIONS</b>
<p><b>Users Should</b></p> <p>Wash hands before eating drinking chewing gum using tobacco or using the toilet            Remove clothing/PPE immediately if pesticide gets inside Then wash thoroughly and put on clean clothing            Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing</p>

4/12

### ENVIRONMENTAL HAZARDS

**DO NOT** apply directly to water to areas where surface water is present or to intertidal areas below the mean high water mark **DO NOT** apply when weather conditions favor drift from treated areas **DO NOT** contaminate water when disposing of equipment washwater or rinsate

Groundwater Advisory Statement This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use Use of this chemical in areas where soils are permeable particularly where the water table is shallow may result in ground water contamination

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

**DO NOT** apply this product in a way that will contact workers or other persons either directly or through drift Only protected handlers may be in the area during application For any requirements specific to your State or Tribe consult the agency responsible for pesticide regulation

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170 This standard contains requirements for the protection of agricultural workers on farms forests nurseries and greenhouses and handlers of agricultural pesticides It contains requirements for training decontamination notification and emergency assistance It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry intervals The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard

**DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours

Exception If the seed is treated with the product and the treated seed is soil injected or soil incorporated the Worker Protection Standard under certain circumstances allows workers to enter the treated area if there will be no contact with anything that has been treated

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants soil or water is coveralls shoes plus socks and chemical resistant gloves such as barrier laminate or nitrile rubber or neoprene rubber or Viton

### GENERAL PRECAUTIONS AND RESTRICTIONS

This product is a systemic fungicide seed dressing specifically for control of systemic downy mildews *Pythium* and *Phytophthora spp* For control of other soilborne diseases combination of captan thiram and carboxin registered formulations are compatible **DO NOT** use with other seed treatment products unless previous experience assures compatibility

**Note** This product is a systemic fungicide having a specific mode of action This product could be subject to development of resistant strains of fungi Development of resistance cannot be predicted Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible metalaxyl resistant strains of fungi which may occur **FAILURE TO FOLLOW DIRECTIONS OF THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL**

**In California** Only for use in commercial seed treating establishments Not allowed for on farm use

**On Farm Seed Treatment** **DO NOT** apply this product through any type of irrigation system **DO NOT** use treated seed for food or feed purposes or process for oil Treat only those seeds

needed for immediate use minimizing the interval between treatment and planting **DO NOT** store excess treated seeds beyond planting time

Seed treatment on agricultural establishment at or immediately before planting is within the scope of WPS while commercial treatment of seeds is not within the scope

**Planter Box Treatment** Apply at the specified rate and premix with the seed directly in the planter box at planting Place half the amount of seed into the planter box and apply half the required amount of pesticide to the seed Stir the seed thoroughly with a paddle until the seed is uniformly coated Place the remaining half of the seed into the planter box and add the remaining half of the pesticide Again stir the seed with the paddle until all seed is thoroughly coated

**For Commercial Seed Treatment** Seed treated commercially or for sale must be dyed an unnatural color using an EPA approved dye before sale Federal law requires that bags containing treated seed shall be labeled with the following information

This seed has been treated with Sebring 318 FS Fungicide which contains the active ingredient metalaxyl **DO NOT** use treated seed for food feed or oil purposes Store away from feed and foodstuffs **DO NOT** harvest mustard greens Rape greens grown and harvested from Sebring 318 FS Fungicide treated seed must not be used for human and feed consumption Rapeseed grown and harvested from Sebring 318 FS Fungicide treated seed is only for industrial uses and cannot be used for edible oil or any other human/feed consumption **DO NOT** use Metalaxyl or mefenoxam for control of Pythium in water seeded rice

**Tank Mixing** When tank mixing follow the most restrictive label instructions for any tank mix partner **DO NOT** tank mix with any product that contains a prohibition against tank mixing

**SEED TREATMENT**

This product may be applied as a water based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist type commercial seed treatment equipment

**DOMESTIC USE**

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)
<b>Brassica Leafy Vegetables*</b> Broccoli (including Chinese and raab) Brussels sprout Cabbage (including Chinese bok choy Chinese napa and mustard) Cauliflower Cavalo broccoli Collards Kale Kohlrabi Mizuna Mustard greens Mustard Spinach and Rape greens)	<i>Pythium</i> damping off control	0.75
<b>Canola*</b> <b>Mustard Seed*</b> <b>Rapeseed*</b>	<i>Pythium</i> damping off control  <b>NOTE DO NOT</b> harvest mustard greens Rape greens grown and harvested from Sebring 318 FS Fungicide treated seed must not be used for human and feed consumption Rapeseed grown and harvested from Sebring 318 FS Fungicide treated seed is only for industrial uses and cannot be used for edible oil or any other human/feed consumption	0.25 - 0.50
<b>Cotton</b>	<i>Pythium</i> seed rot and damping off control	0.75 - 1.50

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)	
<b>Cucurbit Vegetables*</b> Chayote (fruit) Chinese waxgourd (Chinese preserving melon) Citron melon Cucumber Gherkin Gourd (edible) (includes hyotan cucuzza hechima Chinese okra) <i>Momordica</i> spp (includes balsam apple balsam pear bittermelon Chinese cucumber) Muskmelon (including hybrids and/or varieties of <i>Cucumis melo</i> (including true cantaloupe cantaloupe casaba Santa Claus melon crenshaw melon honeydew melon honey balls Persian melon golden pershaw melon mango melon pineapple melon snake melon) Watermelon (including hybrids and/or varieties of <i>Citrullus</i> spp) Pumpkin Squash (summer winter including butternut squash calabaza hubbard squash acorn squash spaghetti squash)	<i>Pythium</i> damping off control	0.75	
<b>Forage Grasses</b> Grasses grown for hay grazing or silage Corn (fodder or silage) Sorghum (hay or silage) and Small grain grown for hay grazing or silage	<i>Pythium</i> damping off control	0.75	
<b>Forage Legumes</b> Alfalfa Beans (for forage) Clover Cowpeas (for forage) Cowpea hay Lespedeza Peanuts (for forage) Peanut hay Peas (for forage) Pea vine hay Soybeans (for forage) Soybean hay Trefoil Velvet beans (for forage) and Vetch	<i>Pythium</i> damping off and early season <i>Phytophthora</i> control	0.75 1.50	
<b>Fruiting Vegetables*</b> Eggplant Ground cherry ( <i>Physalis</i> spp) Pepino Pepper (includes bell pepper chili pepper cooking pepper pimento sweet pepper) Tomatillo Tomato	<i>Pythium</i> damping off control	0.75	
<b>Grain Crops</b> Barley Buckwheat Corn (field corn sweet corn and popcorn) Millet Milo Oats Rye Triticale and Wheat	<i>Pythium</i> damping off control	0.75	
	Systemic downy mildew control of sweet corn	1.50	
<b>Use Rate Conversions for Field Corn</b>			
<b>PRODUCT FLUID OUNCES /</b>			
<b>100 lb seed</b>	<b>1000 seeds</b>	<b>80 000 seed count</b>	<b>MG A / SEED</b>
0.75	0.0042	0.334	0.040
1.50	0.0084	0.672	0.080
This product contains 9.4045 mg metalaxyl (AI) / fluid ounce Based on standard number of 1786 seeds / lb (100 000 seeds / 56 lb bu) <b>NOTE</b> To maintain mg ai / seed adjust fluid ounces / 100 lb in proportion to the actual seeds / lb compared to the standard number of seeds / lb for that crop For example if there are 1600 corn seeds / lb and the targeted rate is 0.040 mg ai / seed at a non adjusted rate of 0.75 fl oz / 100 lb of seed use $(1600 - 1786) \times 0.75 = 0.67$ fl oz / 100 lb of seed			

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)
<b>Leafy Vegetables*</b> Amaranth (Chinese spinach) Cardoon Celery Celery (Chinese) Celeriac Celtuce Chervil Chrysanthemum (edible leaved garland) Corn salad Cress (garden upland) Dandelion Dock (sorrel) Endive (escarole) Fennel (Florence) Lettuce (head and leaf) Orach Parsley Purslane (garden winter) Radicchio (red chicory) Rhubarb Spinach (including New Zealand and vine) Swiss chard	<i>Pythium</i> damping off control	0 75
<b>Onions (dry bulb and green)*</b>	<i>Pythium</i> damping off control	0 75
<b>Peanuts*</b>	<i>Pythium</i> seed rot and damping off control	0 75
<b>Rice*</b>	<i>Pythium</i> damping off control	0 75 1 50
<b>Root and Tuber Vegetables* (except potato)</b> Arracacha Arrowroot Artichoke (Chinese Jerusalem) Beet (garden sugar) Burdock (edible) Canna (edible) Carrot Cassava (bitter and sweet) Chayote (root) Chervil (turnip rooted) Chicory Chufa Dasheen (taro) Ginger Ginseng Horseradish Leren Parsley (turnip rooted) Parsnip Radish (including oriental) Rutabaga Salsify (including black and Spanish) Skirret Sweet potato Tanier Turmeric Turnip Yam bean Yam (true)	<i>Pythium</i> damping off control	0 75
<b>Seed and Pod Vegetables</b> Black eyed peas Chickpea (Garbanzos) Cowpeas Edible soybeans Field beans Field peas Garden peas Green beans Kidney beans Lentils Lima beans Lupines Navy beans Okra Peas Pole beans Snap beans String beans and Wax beans	<i>Pythium</i> damping off control	0 75
	Early season <i>Phytophthora</i> control	0 75
	Control of <i>Pythium</i> damping off and systemic downy mildew of peas	1 50

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)		
Sorghum	<i>Pythium</i> damping off control	0.375 - 0.75		
	Systemic downy mildew (races susceptible to chemical control)	0.75 - 3.00		
	<b>Note</b> Apply 0.75–1.50 fluid ounces of this product on cultivars which possess sorghum downy mildew resistance. To those cultivars which do not possess resistance to a particular race of sorghum downy mildew in heavily infected areas with recent severe history, apply 1.50 – 3.00 fluid ounces of this product. Only high quality sorghum seed (90% germination and above) should be treated with this product.			
Soybeans	<i>Pythium</i> damping off control and early season <i>Phytophthora</i> control	0.75 - 1.50		
<b>Use Rate Conversions for Soybeans</b>				
<b>PRODUCT FLUID OUNCES /</b>				
100 lb seed	1000 seeds	80 000 seed count	<b>MG AI / SEED</b>	
0.75	0.0025	0.20		0.0235
1.50	0.005	0.40		0.047
This product contains 9.4045 mg metalaxyl (AI) / fluid ounce				
Based on standard number of 3000 seeds / lb (150 000 seeds / 50 lb bu)				
<b>NOTE</b> To maintain mg ai / seed, adjust fluid ounces / 100 lb in proportion to the actual seeds / lb compared to the standard number of seeds / lb for that crop. For example, if there are 2800 soybean seeds / lb and the targeted rate is 0.0235 mg ai / seed at a non-adjusted rate of 0.75 fl oz / 100 lb of seed, use $(2800 - 3000) \times 0.75 = 0.70$ fl oz / 100 lb of seed.				
Sunflowers	Systemic downy mildew control	1.50 - 3.00		
Turf Grasses Golf Courses Home Lawns Reclamation and Erosion Control Projects	<i>Pythium</i> damping off control	1.50		

\* Not registered for use in California



**FOR EXPORT USE ONLY**

For treatment of the following seed for export to countries requiring treatment with metalaxyl before importation

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)
Brassica (Cole) Leafy Vegetables	<i>Pythium</i> damping off control	0 75 1 50
Canola	<i>Pythium</i> damping off control	0 50 1 70 (31 100 ml per 100 Kg of seed)
Corn	<i>Pythium</i> damping off control	0 75 1 50
	Systemic downy mildew control	3 00
Cucurbit Vegetables	<i>Pythium</i> damping off control	0 75
Fruiting Vegetables	<i>Pythium</i> damping off control	0 75 1 50
Leafy Vegetables	<i>Pythium</i> damping off control	0 75 1 50
Millet	<i>Pythium</i> damping off control	0 75 1 50
	Systemic downy mildew control	2 50
Onions	<i>Pythium</i> damping off control	0 75 1 50
Peas	<i>Pythium</i> damping off control and early season <i>Phytophthora</i> control	3 50
Root and Tuber Vegetables	<i>Pythium</i> damping off control	0 75 1 50
	<b>NOTE DO NOT</b> apply this product to potatoes	
Sorghum	<i>Pythium</i> damping off control	0 75 1 50
	Systemic downy mildew control	5 25
Sunflower	Systemic downy mildew control	9 00

Seed treated in accordance with the export use pattern must be labeled as follows

**FOR EXPORT ONLY  
NOT FOR DOMESTIC SALES OR USE**

**DIRECTIONS FOR USE AT REDUCED RATES IN COMBINATION WITH OTHER FUNGICIDES**

This product is a systemic fungicide seed dressing which may be applied in combination with other EPA registered fungicides to aid in control of *Pythium*. This product may be applied as a water based slurry through standard slurry of mist type commercial seed treatment equipment

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)
Barley, Corn (field), Oats, Rye and Wheat Beans (field kidney, navy and pinto), Lentils Peas and Soybeans Peanuts* Rice* Sunflowers	Aid in control of seed decay and damping off caused by <i>Pythium</i>	0 10 0 375

\* Not registered for use in California

**NOTE** Apply only in combination with EPA registered rates of broad spectrum seed treatment fungicides

**STORAGE AND DISPOSAL**

**DO NOT** contaminate water food or feed by storage or disposal Open dumping is prohibited

**PESTICIDE STORAGE** Store in cool dry location **DO NOT** store in direct hot sunlight **DO NOT** store above 90 F for prolonged periods of time After prolonged storage in intense cold allow product to return to approximately 75 F before using

**PESTICIDE DISPOSAL** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility

**CONTAINER DISPOSAL**

[Note to Reviewer The following statement will be included on all Final Printed Labels bearing multiple Container Disposal (Container Handling) statements] **NOTE** This product is available in multiple containers Refer to the Net Contents section of this products labeling for the applicable Nonrefillable or Refillable designation Follow the container disposal [handling] instructions below that apply to your container type / size

**[Nonrefillable Containers 5 Gallons or Less]**

**Nonrefillable container DO NOT** reuse or refill this container Triple rinse container (or equivalent) promptly after emptying

**Triple rinse as follows** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container 1/4 full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities Plastic containers are also disposable by incineration or if allowed by State and local authorities by burning If burned stay out of smoke

**[Nonrefillable containers larger than 5 gallons]**

**Nonrefillable container DO NOT** reuse or refill this container Offer for recycling if available Triple rinse or pressure rinse container (or equivalent) promptly after emptying

**Triple rinse as follows** Empty the remaining contents into application equipment or a mix tank Fill the container 1/4 full with water Replace and tighten closures Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds Stand the container on its end and tip it back and forth several times Turn the container over onto its other end and tip it back and forth several times Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal Repeat this procedure two more times Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities Plastic containers are also disposable by incineration or if allowed by State and local authorities by burning If burned stay out of smoke

**Pressure rinse as follows** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal Insert pressure rinsing nozzle in the side of the container and rinse at about 40 psi for at least 30 seconds Drain for 10 seconds after the flow begins to drip

**[Refillable containers]**

**Refillable container** Refill this container with pesticide only **DO NOT** reuse this container for any other purpose Cleaning the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller To clean the container before final disposal empty the remaining contents from this container into application equipment or a mix tank Fill the container about 10% full with water and if possible spray all sides while adding water Agitate vigorously or recirculate water with the pump for two minutes Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times

**[Refillable For return to Nufarm ]**

**Refillable Container** Refill this container with pesticide only **DO NOT** reuse this container for any other purpose Close all openings and replace all caps Contact Nufarm s Customer Service Department at 1 800 345 3330 to arrange for return of the empty refillable container

**WARRANTY DISCLAIMER**

The directions for use of this product must be followed carefully TO THE EXTENT CONSISTENT WITH APPLICABLE LAW (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE UNINTENDED CONSEQUENCES INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS OR THE MANNER OF USE OR APPLICATION INCLUDING WEATHER ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER MANUFACTURER AND SELLER AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS

**LIMITATION OF LIABILITY**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL INCIDENTAL OR CONSEQUENTIAL DAMAGES OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD INCLUDING USE APPLICATION HANDLING AND DISPOSAL TO THE EXTENT CONSISTENT WITH APPLICABLE LAW MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER IF ANY OR FOR ANY DAMAGES OR SUMS OF MONEY CLAIMS OR DEMANDS WHATSOEVER RESULTING FROM OR BY REASON OF OR ARISING OUT OF THE MISUSE OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER USER OR ITS CUSTOMERS TO THE EXTENT CONSISTENT WITH APPLICABLE LAW BUYER'S OR USER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT

**If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability do not use the product, and return it unopened to the Seller, and the purchase price will be refunded**

(RV042712)

Sebring is a trademark of Nufarm Americas Inc

**LABEL HISTORY**

<u>File Name</u>	<u>Revision Mark</u>	<u>Comment</u>
055146 00107 20100818 Revised_Use_Table	RV081810	Revised Use Tables
055146 00107 20101108 MASTER	RV110810	EPA Approval
055146 00107 20120427 Revised_Label	RV042712	Addition of Use Rate conversions for Corn & Soybeans Update of Container Disposal Statement to add flexibility for single label on multiple container types