

55146-117

8/28/2012

1/10

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460



AUG 28 2012

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Nathan Ehresman
Nufarm Americas Inc.
150 Harvester Drive
Burr Ridge, IL 60527

Subject: Agency-initiated amendment to revise the plant-back interval by deleting oats under "30 days" and keeping under "immediate"
Spirato IM 413 FS
EPA Reg. No. 55146-117

Dear Mr. Ehresman:

The labeling referred to above submitted in connection with the Federal Insecticide, Fungicide and Rodenticide Act, as amended is acceptable.

A stamped copy of the label is enclosed for your records. Please submit one final printed copy of the labeling before releasing the product for shipment. If you have any questions regarding this label, please contact Autumn Metzger at (703) 305-5314 or metzger.autumn@epa.gov.

Sincerely,

A handwritten signature in cursive script that reads "Venus Eagle".

Venus Eagle
Product Manager 01
Insecticide-Rodenticide Branch
Registration Division (7505P)

SPIRATO™ IM 413 FS

For use as a seed treatment to protect seed and seedlings of listed legume vegetables (including soybeans) against listed insect and crop diseases.

ACTIVE INGREDIENT:

Imidacloprid.....	27.41%
Metalaxyl.....	6.85%
Fludioxonil.....	1.11%

OTHER INGREDIENTS:	64.63%
TOTAL:	100.00%

Contains 2.67 lbs imidacloprid, 0.67 lbs metalaxyl, and 0.11 lbs fludioxonil per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 55146 - RRT
EPA EST. NO. _____

MANUFACTURED FOR
NUFARM AMERICAS INC.
AGT DIVISION
150 HARVESTER DRIVE
BURR RIDGE, IL 60527



ACCEPTED
AUG 28 2012
Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under:

EPA. Reg. No: 55146-117

NET CONTENTS _____ GAL. (_____ Liters)
[Designation as "NONREFILLABLE" or "REFILLABLE" for containers > 5 GAL]

FIRST AID

IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- **DO NOT** induce vomiting unless told to by a poison control center or doctor.
- **DO NOT** give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement

When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to birds, fish and aquatic invertebrates. For terrestrial use, **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment wash water or rinsate. Cover or incorporate spilled treated seeds.

Groundwater Advisory: Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Pollinator Precautions: Imidacloprid is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Ensure that planting equipment is functioning properly in accordance with manufacturer specification to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use in commercial seed treatment facilities, or as an end-use seed treatment on agricultural establishments at, or immediately before, planting. This product is to be used in liquid or slurry treaters only.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the **restricted-entry interval (REI)** of 12 hours.

Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Shoes plus socks

PRODUCT INFORMATION

SPIRATO IM 413 FS is a seed treatment product that protects against certain early season insects, soil-borne and seed-borne diseases of listed crop plants. SPIRATO IM 413 FS contains the active ingredients imidacloprid, metalaxyl and fludioxonil.

Imidacloprid is a systemic seed treatment insecticide in the neonicotinoid class. Imidacloprid protects against listed chewing and sucking insect pests.

Metalaxyl is a fungicide which protects against *Pythium*, *Phytophthora* and systemic downy mildew.

Fludioxonil is a fungicide which protects against *Fusarium*, *Rhizoctonia*, and suppresses seed-borne *Sclerotinia* and *Phomopsis* spp.

When using this product on-farm for seed treatment prior to planting, you must comply with the following USE RESTRICTIONS. If treated seed will not be immediately planted, but will be stored for planting at a later date, you must label the treated seed according to the "SEED BAG LABELING REQUIREMENTS" specified on this label.

USE RESTRICTIONS

DO NOT use as a planter (hopper) box treatment.

CALIFORNIA: Only for use in commercial seed treatment facilities, not allowed for "on farm" use.

ON FARM TREATMENT: DO NOT apply this product through any type of irrigation system.

DO NOT use for feed, food, or oil purpose.

Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.

Store away from feeds and foodstuffs.

Treated seed must be planted into the soil at a depth greater than 1 inch.

Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds that are spilled during handling.

Dispose of all excess treated seed. Leftover seed may be doublesown around the headland or buried away from water sources in accordance with local requirements.

DO NOT contaminate water bodies when disposing of planting equipment wash water.

In the event of a crop failure or harvest of a crop grown from SPIRATO IM 413 FS treated seed the field may be replanted immediately to artichoke barley Brassica (cole) leafy vegetables canola cilantro corn (field or sweet) cotton cranberry cucurbits eggplant groundcherry leafy petiole vegetables leafy vegetables legume vegetables (succulent or dried) mustard seed oats okra pepinos pepper popcorn potato rapeseed sorghum soybean strawberry sugarbeets tomatillo tomato triticale turnip greens root and tuber vegetables watercress and wheat For cereals including buckwheat millet rice and rye and for safflower the minimum plant back interval is 30 days from the date SPIRATO IM 413 FS treated seed was planted Cover crops for soil building or erosion control may be planted at any time but do not graze or harvest for food or feed For all other crops not listed on an imidacloprid metalaxyl or fludioxonil label or for crops for which no imidacloprid metalaxyl or fludioxonil tolerance for the active ingredient has been established a 12 month plant back interval must be observed

DO NOT allow children pets or livestock to have access to treated seed

The maximum application rate (all imidacloprid uses seed treatment foliar application soil application) per acre per year for imidacloprid is 0.5 lbs **DO NOT** use more than 0.5 lb (226.8 mg) per acre per year

DO NOT apply a neonicotinoid insecticide within 45 days of planting seed treated with SPIRATO IM 413 FS

MIXING INSTRUCTIONS

Mix thoroughly before use or use entire container at one time All tank mixes should be pre tested to determine physical compatibility between formulations Follow the most restrictive precautions and limitations on labeling of all products used in mixtures

Apply SPIRATO IM 413 FS as a water based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage Uneven or incomplete seed coverage may not give the desired level of disease control Thoroughly mix the specified amount of SPIRATO IM 413 FS into the required amount of water or liquid inoculant for the slurry treater and dilution rate to be used

IMPORTANT Always recirculate SPIRATO IM 413 FS thoroughly before using The typical density of SPIRATO IM 413 FS is 9.7 lbs per gallon Consult the manufacturer of the application equipment you plan to use for suitability for this application and for instructions on operation and calibration of the equipment Follow the manufacturer application instructions for the seed treatment equipment being used

Certain crops require addition of inoculants when the seed is treated or planted SPIRATO IM 413 FS is compatible with several liquid inoculant products Consult the maker of the inoculant product and a Nufarm Americas representative for directions before applying SPIRATO IM 413 FS with inoculants

Under certain disease conditions additional amounts of fungicides may be required When needed apply additional SEBRING 318 FS (metalaxyl) and SPIRATO 480 FS (fludioxonil) according to the label **USE SPECIFICATIONS** Other tank mix partners may be used with SPIRATO IM 413 FS however the user must consider the use rate formulation seed and crop safety factors and compatibility of each product to be mixed when determining the total application volume

The total application volume must be sufficient to provide desired level of coverage Dilution is typically done with water or liquid inoculants The minimum slurry volume to achieve adequate coverage is 4.0 fluid ounces per 100 pounds of seed More diluent may be required to obtain complete coverage

Continuous agitation or mixing of the slurry mixture is necessary to prevent settling out of the solution

Allow seed to dry before bagging

Follow planter manufacture specifications for use of talc or other hopper box additives at planting

Seed must be completely dry before adding to planter

SPIRATO IM 413 FS contains an EPA approved dye/colorant that imparts an unnatural color to the seed as required by the Federal Seed Act

Treated seed must not be used for or mixed with food or animal feed or processed for oil Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the Federal Seed Act

6/10

USE INFORMATION

TARGET INSECTS (All Uses)

Early season protection against injury by aphids, bean leaf beetle, grape colaspis, leaf miners, leaf hoppers, Mexican bean beetle, seed corn maggot, three-cornered alfalfa hopper, thrips, white grubs, and wireworm.

SPIRATO IM 413 FS also protects against pea leaf weevil in peas.

When treated according to the following directions for post-planting protection against listed pests, SPIRATO IM 413 FS will also provide protection during post treatment storage of the seed against damage from the following stored grain insects: Indian Meal Moth (*Plodia interpunctella*), Rice Weevil (*Sitophilus oryza*), Red Flour Beetle (*Tribolium castaneum*), and Lesser Grain Borer (*Rhizopertha dominica*).

If seed to be treated has existing infestations of stored grain insects, fumigate the seed prior to treating with SPIRATO IM 413 FS and bagging.

TARGET DISEASES (All Uses)

Protection against damping-off and seed-borne rots due to *Pythium*, *Phytophthora*, *Fusarium*, *Rhizoctonia* species and *Phytophthora* root rot.

SPIRATO IM 413 FS also suppresses seed-borne *Sclerotinia* and *Phomopsis* species.

APPLICATION DIRECTIONS

SEED AND POD VEGETABLES (Black-eyed peas, Chickpeas, Cowpeas, Edible soybeans, Garbanzos, Green beans, Kidney beans, Lentils, Lima beans, Lupines, Navy beans, Pole beans, Snap beans, String beans, Wax beans):

RATE OF SPIRATO IM 413 FS		OPTIONAL TREATMENT	
		See Remarks section below WHEN TO ADD ADDITIONAL SEBRING 318 FS	
		ADDITIONAL SEBRING 318 FS (metalaxyl)	
Fl. Oz. / 100 lb. of seed	Grams AI / 100 kg of seed	Fl. Oz. / 100 lb. of seed	Grams AI / 100 kg of seed
3.0	62.4	0.375	7.8
	15.6	to	to
	2.5	0.75	15.6
	imidacloprid		
	metalaxyl		
	fludioxonil		
REMARKS			
WHEN TO ADD ADDITIONAL SEBRING 318 FS (metalaxyl)			
Add additional SEBRING 318 FS if target fields have a history of high <i>Phytophthora</i> pressure, add additional SEBRING 318 FS as directed in the label use specifications. The additional SEBRING 318 FS may reduce compatibility with some rhizobia inoculants. Consult with the maker of rhizobia inoculants before adding the additional SEBRING 318 FS.			
RESTRICTIONS			
CA RESTRICTION: Seed and Pod Vegetables seed treated in California must be destined for planting in states other than California and is not to be planted in California.			

PEAS (field and garden)

RATE OF SPIRATO IM 413 FS				OPTIONAL TREATMENT See Remarks section below <u>WHEN TO ADD ADDITIONAL SEBRING 318 FS</u> See Remarks section below <u>WHEN TO ADD ADDITIONAL SPIRATO 480 FS</u>			
				ADDITIONAL SEBRING 318 FS (metalaxyl)		ADDITIONAL SPIRATO 480 FS (fludioxonil)	
Fl Oz / 100 lb of seed		Grams AI / 100 kg of seed		Fl Oz / 100 lb of seed	Grams AI / 100 kg of seed		
1.5		31.2	imidacloprid metalaxyl fludioxonil	0.375 to 1.125	7.8 to 23.2	0.04	1.25
REMARKS							
WHEN TO ADD ADDITIONAL SEBRING 318 FS (metalaxyl) Add additional SEBRING 318 FS if target fields have a history of high <i>Phytophthora</i> pressure add additional SEBRING 318 FS as directed in the label use specifications The additional SEBRING 318 FS may reduce compatibility with some rhizobia inoculants Consult with the maker of rhizobia inoculants before adding the additional SEBRING 318 FS Use the higher listed rate (1.125 Fl Oz / 100 pounds of seed) for systemic downy mildew protection							
WHEN TO ADD ADDITIONAL SPIRATO 480 FS (fludioxonil) Add additional SPIRATO 480 FS if planting location has a history of high <i>Fusarium</i> or <i>Rhizoctonia</i> pressure add additional SPIRATO 480 FS as directed in the label use specifications							
RESTRICTIONS							
CA RESTRICTION Peas (field and garden) seed treated in California must be destined for planting in states other than California and is not to be planted in California							

SOYBEANS

RATE OF SPIRATO IM 413 FS Use Rate based on 3000 seeds / lb (150 000 seeds / 50 lb bushel)						OPTIONAL TREATMENT See Remarks section below <u>WHEN TO ADD ADDITIONAL SEBRING 318 FS</u>		
						ADDITIONAL SEBRING 318 FS (metalaxyl)		
Fl Oz /		ml / 1 000 seeds	Grams AI / 100 kg of seed	mg AI / Seed	Fl Oz / 100 lb of seed	Grams AI / 100 kg of seed		
100 lbs seed	1 000 Seeds						140 000 seeds	
3.0	0.01	1.4	0.2957	62.4 15.6 2.5	0.0946 0.0238 0.0038	imidacloprid metalaxyl fludioxonil	0.375 to 0.75	7.8 to 15.6
REMARKS								
WHEN TO ADD ADDITIONAL SEBRING 318 FS (metalaxyl) Add additional SEBRING 318 FS if target fields have a history of high <i>Phytophthora</i> pressure add additional SEBRING 318 FS as directed in the label use specifications The additional SEBRING 318 FS may reduce compatibility with some rhizobia inoculants Consult with the maker of rhizobia inoculants before adding the additional SEBRING 318 FS IMPORTANT To maintain mg ai / seed adjust fluid ounces / 100 lb of seed in proportion to the actual seed count / lb compared to the standard number of seeds / lb for that crop fluid ounces per number of seeds (i.e. 1000 80 000 140 000) does not need to be adjusted For example if there are actually 2800 soybean seeds / lb use $2800 \div 3000 \times 3.0 = 2.8$ fl oz / 100 lb of seed								

SEED BAG LABELING REQUIREMENTS

Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the federal seed act. The user of this product is responsible for ensuring that the seed bag meets all requirements under the federal seed act.

THE FEDERAL SEED ACT REQUIRES THAT BAGS CONTAINING TREATED SEEDS SHALL BE LABELED WITH THE FOLLOWING STATEMENTS

- This seed has been treated with imidacloprid insecticide, metalaxyl and fludioxonil fungicide
- **DO NOT** use for feed, food, or oil purpose

THE U.S. ENVIRONMENTAL PROTECTION AGENCY REQUIRES THE FOLLOWING STATEMENTS ON BAGS CONTAINING SEEDS TREATED WITH SPIRATO IM 413 FS

- **Groundwater Advisory** Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Pollinator Precautions Imidacloprid is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Ensure that planting equipment is functioning properly in accordance with manufacturer specification to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.

Store away from feeds and foodstuffs

- Wear long sleeved shirt, long pants and chemical resistant gloves when handling treated seed. Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds that are spilled during handling. Treated seed must be planted into the soil at a depth greater than 1 inch. Dispose of all excess treated seed. Leftover seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. **DO NOT** contaminate water bodies when disposing of planting equipment wash water.
- Dispose of seed packaging in accordance with local requirements. In the event of a crop failure or harvest of a crop grown from SPIRATO IM 413 FS treated seed, the field may be replanted immediately to artichoke, barley, Brassica (cole), leafy vegetables, canola, clantro, corn (field or sweet), cotton, cranberry, cucurbits, eggplant, groundcherry, leafy petiole vegetables, leafy vegetables, legume vegetables (succulent or dried), mustard seed, oats, okra, pepinos, pepper, popcorn, potato, rapeseed, sorghum, soybean, strawberry, sugarbeets, tomatillo, tomato, triticale, turnip greens, root and tuber vegetables, watercress, and wheat. For cereals including buckwheat, millet, rice, and rye, and for safflower, the minimum plant back interval is 30 days from the date SPIRATO IM 413 FS treated seed was planted. Cover crops for soil building or erosion control may be planted at any time, but do not graze or harvest for food or feed. For all other crops not listed on an imidacloprid, metalaxyl, or fludioxonil label, or for crops for which no imidacloprid, metalaxyl, or fludioxonil tolerance for the active ingredient has been established, a 12-month plant back interval must be observed. **DO NOT** allow children, pets, or livestock to have access to treated seed. The maximum application rate (all imidacloprid uses: seed treatment, foliar application, soil application) per acre per year for imidacloprid is 0.5 lbs. **DO NOT** use more than 0.5 lb (226.8 mg) per acre per year.
- This seed has been treated with X mg imidacloprid, X mg metalaxyl, and X mg fludioxonil per seed. **DO NOT** apply a neonicotinoid insecticide within 45 days of planting seed treated with SPIRATO IM 413 FS.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool place, **DO NOT** store in direct sunlight. Protect from freezing temperatures.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL [HANDLING]:

[Note to Reviewer: The following statement will be included on all Final Printed Labels bearing multiple Container Disposal (Container Handling) statements] "This product is available in multiple containers. Refer to the Net Contents section of this products labeling for the applicable "Nonrefillable" or "Refillable" designation. Follow the container disposal [handling] instructions below that apply to your container type / size."

[Nonrefillable Containers 5 Gallons or Less]

Nonrefillable container. DO NOT reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

[Nonrefillable containers larger than 5 gallons]

Nonrefillable container. DO NOT reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

[Refillable containers larger than 5 gallons]

Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

[Refillable containers for return to Nufarm]

Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Close all openings and replace all caps. Contact Nufarm's Customer Service Department at 1-800-345-3330 to arrange for return of the empty refillable container. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1)THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULARTRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BYWAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALL SUCH RISKS SHALL BE ASSUMED BYTHE BUYER, USER, OR ITS CUSTOMERS .TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV062012)

Spirato and Sebring are registered trademark of Nufarm Americas Inc.

All other trademarks are the property of their respective owners.