

FINAL LABEL COLDCIDE-0.25% DISINFECTANT DEODORIZING SPRAY PAGE 1 09-26-1997

COLDCIDE - 0.25% DISINFECTANT/DEODORIZING SOLUTION & PUMP SPRAY

Glutaraldehyde / Compatible Dual Phenolics

Kills Herpes simplex virus Type 1 & 2, Influenza A₂ (Hong Kong) & HIV-1 (AIDS) virus** in 30 seconds on hard non-porous surfaces.

Virucidal*	ACTIVE INGREDIENTS:	
Fungicidal	Glutaraldehyde	0.27600%
Bactericidal	O-Phenylphenol	0.01375%
Tuberculocidal	P-Tertiary Amylphenol	0.00275%
Pseudomonacidal	INERT INGREDIENTS	<u>99.70850%</u>
	TOTAL	100.00000%

ACCEPTED
MAR 27 1998
55195-3

REFILL AVAILABLE

CAUTION: KEEP OUT OF REACH OF CHILDREN
See side panels for additional precautionary statements.

ColCide, Inc.
11500 W. Hill Dr.
Rockville, MD 20852

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

CONTENTS: 20 FL. OZ. (591 ml) or 1 quart (.9463 l)

U.S. Patent #4,469,614 + 4,654,374 / Additional Patent Pending

EPA Reg. No. 55195-3
EPA Est. Nos Dymon, Inc. 11694-KS-01 Metrex Research, Inc. 46781-CO-1

Expiration Date: 2 Years After Date Manufactured
Production Lot No. is stamped on bottom of container.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For use in non-critical care areas as a general disinfectant spray on hard surfaces.

Disinfection: This product is to be used for the disinfection of hard, inanimate surfaces such as counter tops, tables, bed springs, and frames, bedside and office furniture, chart racks, equipment, cabinets, door knobs, office machinery, sinks, mirrors, light fixtures, switches, trash carts, telephones, waste cans, laundry hampers, wheel chairs, walkers, urinals, exam tables, cages, or other similar, hard surfaces. This product is to be used as is, undiluted. Contact time is 10 minutes at room temperature (20°C/68°F) or higher) for bactericidal activity, 2 minutes for virucidal activity, and 15 minutes at room temperature for tuberculocidal activity. Thoroughly wet pre-cleaned surface(s) and allow to air dry.

ColdCide-0.25% Disinfectant Solution & Pump Spray kills Salmonella choleraesuis, Staphylococcus aureus, and Pseudomonas aeruginosa, and Trichophyton mentagrophytes in 10 minutes at 20°C/68°F or higher. Mycobacterium tuberculosis is completely inactivated in 15 minutes at room temperature (25°C or higher). Herpes simplex virus 1 & 2 and Influenza A₁HK are inactivated in 30 seconds, and the listed viruses* within 2 minutes at 20°C/68°F or higher: Coxsackie B5a, Adenovirus-2 virus, Poliovirus 1 & 2, Rotavirus SA-11, Rhinovirus, Vaccinia virus, Cytomegalovirus and Respiratory Syncytial Virus. ColdCide-0.25% Disinfectant Solution & Pump Spray inactivates the HIV-1 (AIDS virus) on hard, non-porous surfaces in 30 seconds at room temperature (20-25°C). See panel for complete HIV-1 (AIDS virus) information.

Deodorizing: For effective deodorizing, apply to hard, non-porous surfaces to eliminate generally offensive odors due to spillage from post-colostomy, post-ileostomy dressing and bag changes, and other spills such as urine or fecal matter; smoke or soot deposits, industrial plant processes, mildew and mold, and surfaces in musty closets.

PRECAUTIONARY STATEMENTS: Hazard to Humans and Domestic Animals.

CAUTION: May be irritating to eyes. Do not get in eyes, on skin or on clothing. Wear safety glasses and rubber gloves when handling. Repeated dermal contact may result in dermal sensitization in some individuals. Harmful if swallowed. Avoid contamination of food. Avoid reuse of unwashed, disinfectant soaked gloves. Avoid prolonged inhalation.

STATEMENT OF PRACTICAL TREATMENT: If in eyes, immediately flush with water for at least 15 minutes. Get medical attention. If on skin, wash thoroughly with water. Wash contaminated clothing before reuse.

WASH THOROUGHLY AFTER HANDLING.

STORAGE AND DISPOSAL: Store in cool, well ventilated area. Unused product may be discarded by pouring down drain and flushing with large quantity of water. Rinse empty container thoroughly and discard in trash.

Manufactured in U.S.A. by Dymon, Inc.
Olathe, KS 66061
1-800-224-4860

Or by Metrex Research, Inc.
Parker, CO 80134
1-800-841-1428

ACCEPTED

MAR 27 1998

551953

373

HIV-1 (AIDS VIRUS) INFORMATION
THE ACTIVITY OF COLDCIDE-0.25% DISINFECTANT DEODORIZING SPRAY
(EPA REG. #55195-3)
AGAINST THE HIV-1 (AIDS VIRUS)

ColdCide-0.25% Disinfectant Solution & Pump Spray completely inactivates the HIV-1 (AIDS virus) on hard, non-porous surfaces in the presence of a moderate amount of organic soil (5% blood serum) at a contact time of 30 seconds at room temperature (20-25°C). Although efficacy at a 30 second contact time has been shown to be adequate against HIV-1, this time would not be sufficient for other organisms. Use a 15-minute contact time for disinfection against all of the organisms claimed.

ColdCide-0.25% Disinfectant Solution & Pump Spray kills HIV-1 on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1)(associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

Personal Protection: Specific barrier protection items to be used when handling items soiled with blood or body fluids are disposable latex gloves, gowns, masks, and eye coverings.

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of ColdCide-0.25% Disinfectant Solution & Pump Spray.

Contact Time: Leave surfaces wet for 30 seconds. This contact time will not control other common types of viruses or bacteria.

Disposal of Infectious Materials: Materials that may contain blood or other body fluids should be autoclaved and disposed of according to Federal, State and Local regulations for infectious waste disposal.

ColdCide, Inc.
11500 W. Hill Dr.
Rockville, MD 20852
1-301-468-1668

