



SS 620-6

07-29-2009

1/11

<p align="center"><b>U.S. ENVIRONMENTAL PROTECTION AGENCY</b></p> <p align="center">Office of Pesticide Programs Registration Division (7505P) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460</p> <p>NOTICE OF PESTICIDE:  <input type="checkbox"/> Registration  <input checked="" type="checkbox"/> Reregistration          (under FIFRA, as amended)</p>	<p>EPA Reg. Number: <b>55620-6</b></p>	<p>Date of Issuance: <b>JUL 29 2009</b></p>
	<p>Terms of Issuance: <b>Conditional</b></p>	
	<p>Name of Pesticide Product: SYLLIT FL Fungicide</p>	
<p>Name and Address of Registrant (include ZIP Code):  <b>Agriphar S.A.</b>  <b>Rue De Renory, 26</b>  <b>B-4102 Ougree, Belgium</b></p>		
<p><b>Noté:</b> Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>Based on your response to the Reregistration Eligibility Document(s) for Dodine, EPA has reregistered the product listed above provided that the following changes are made to the label:</p> <ol style="list-style-type: none"> <li>1. On page 2, in the Precautionary Statements, move the heading, HAZARDS TO HUMANS AND DOMESTIC ANIMALS" above "DANGER."</li> <li>2. On page 2, in the Environmental Hazards section, revise the first sentence to read "This Pesticide is toxic to fish and aquatic invertebrates."</li> </ol>		
<p>Signature of Approving Official:            Mary L. Waller          Product Manager (Team 21)          Fungicide Branch          Registration Division</p>	<p>Date:  </p>	

- 3. On page 2, in the Surface Water and Erosion Control Statement section, add the following statements to the beginning of the paragraph:

“This product may contaminate water through runoff and has a potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product.”

- 4. On page 4, in the first sentence of the Compatibility section, change the word “recommended” to “specified.”
- 5. On page 4, in the Compatibility section, delete “Dinocap.”
- 6. On page 6, under “Apple scab – After-Infection Schedule, revise the third sentence to read “Fruit color and finish on most apple varieties sprayed with SYLLIT FL at labeled dosages have frequently been outstanding.”

This action is taken under the authority of section 4(g)(2)(c) of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain the registration of your product.

Please note that this amended label is in response to decision number 414525 (fast track amendment dated 5/18/09) and decision number 383339 (reregistration amendment).

Submit one (1) copy of final printed labeling incorporating the above changes before you release the product for shipment. Amended labeling will supersede all previously accepted labels. A copy of your label stamped “Accepted with Comments” is enclosed for your records.

If you have questions concerning this letter, please contact Rose Kearns of my staff by phone at 703-305-5611 or via email at [kearns.rosemary@epa.gov](mailto:kearns.rosemary@epa.gov) or myself at 703-308-9354 or via email at [waller.mary@epa.gov](mailto:waller.mary@epa.gov).

Sincerely,



Mary L. Waller  
Product Manager (21)  
Fungicide Branch  
Registration Division

3  
11

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

JUL 29 2009

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

55620-6



# SYLLIT<sup>®</sup> FL FUNGICIDE

FOR CONTROL OF CERTAIN DISEASES OF APPLES, BANANAS, CHERRIES,  
PEACHES, PEANUTS, PEARS, PECANS, AND WALNUTS.

**ACTIVE INGREDIENT:**

Dodine (n-dodecylguanidine acetate) **39.6%\***

**OTHER:** **60.4%**

**TOTAL:** **100.0%**

\*Equivalent to 3.4 pounds Dodine per gallon

EPA Reg. No. 55260-6

EPA Est. No. 55260-BL-001

**KEEP OUT OF REACH OF CHILDREN  
DANGER PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

### FIRST AID

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing and shoes. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

### HOT LINE NUMBER

For MEDICAL and TRANSPORTATION EMERGENCIES ONLY call **CHEMTREC 24 hours a day 800-424-9300**. For PRODUCT USE Information call your Distributor.

"Have the product container or label with you when calling a poison control center or doctor, or going to treatment."

### NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the usage of gastric lavage.

## PRECAUTIONARY STATEMENTS

### DANGER

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Corrosive. Causes irreversible eye damage. Causes skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Avoid breathing spray mist or vapor.

Some materials that are chemical-resistant to this product are barrier laminate or viton. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

All mixers, loaders, applicators and other handlers must wear:

- Coveralls over short sleeved shirt and short pants,
- Chemical-resistant footwear plus socks,
- Chemical-resistant gloves,
- Goggles or face shield,
- Chemical-resistant headgear for overhead exposure,
- Chemical-resistant apron for mixers and loaders

See engineering controls for additional requirements.

#### USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning /maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### ENGINEERING CONTROLS

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.249(d)(6)].

#### USER SAFETY RECOMMENDATIONS

User should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of the gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Surface Water and Erosion Control Statement: A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

### DIRECTIONS FOR USE

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.** Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is:

- coveralls worn over short-sleeve shirt and short pants,
- chemical-resistant gloves made of any waterproof material,
- chemical-resistant footwear plus socks,
- protective eyewear, and
- chemical-resistant headgear (if overhead exposure).

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Do not store in or around the home. Storage should be under lock and key and secure from access by unauthorized persons, children and animals. Do not use or store near heat, open flame or hot surfaces. Avoid storage near water supplies, food, feed and fertilizer to avoid contamination. Carefully open containers. After partial use, replace lid and keep container tightly sealed. Keep container tightly closed when not in use. Avoid contamination with acids and alkalies. Store in original containers only.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**CONTAINER HANDLING: Nonrefillable Containers:** Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying.

**Nonrefillable container equal to or less than 5 gallons:** Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

## **GENERAL USE PRECAUTIONS**

Do not apply this product through any type of irrigation system.  
Do not apply in greenhouses.  
Do not graze livestock in treated fields, orchards or treated groves.  
Do not feed hay or vines (peanuts) to livestock.  
Consult your state experiment station or state extension service for proper timing of sprays.

## **RESISTANCE MANAGEMENT**

It is known that certain plant pathogens might develop resistance to products used repeatedly for disease control. For good resistance management practices, it is recommended to follow the specific use directions on the label. It is essential to alternate SYLLIT FL with other labeled fungicides that have a different mode of action. This management program will prevent the buildup of resistance and will ensure that SYLLIT FL Fungicide can be maintained as a valuable fungicide. Where resistance to SYLLIT FL is known or suspected, use an alternate fungicide for control of the disease.

## **DILUTE AND CONCENTRATE SPRAYS**

SYLLIT FL may be used effectively in dilute and concentrate sprays when applied properly to deliver the labeled rate (pints per acre). Trees must be covered uniformly. Do not excessively drench fruit and foliage with dilute sprays. Avoid heavy deposits when using concentrate sprays.

## **INSTRUCTIONS FOR MIXING**

Add SYLLIT FL directly to the spray tank when half filled with water. Keep agitator running during filling and spraying.

## **COMPATIBILITY**

Experience indicates that SYLLIT FL can be used effectively at recommended dosages with most of the commonly used fungicides, insecticides and acaricides. However, we always recommend to test the compatibility of the products used in combination with SYLLIT FL at small scale before spraying or to ask your local distributor about potential incompatibility.

While SYLLIT FL has been used satisfactorily with emulsifiable concentrate formulations and flowable (water-based) emulsions, it is desirable to check the compatibility of this type of combination before materials are mixed in the spray tank. SYLLIT FL is not compatible with foliar fertilizer, Dinocap, Captan, Bordeaux mixtures, fixed copper, lime or lime sulfur. Observe all cautions and limitations on labeling of all products used in mixtures. Follow the most restrictive label limitations and precautions. No label dosage rates may be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

## **SYLLIT FL AND OIL**

Most of the tank-mix combinations of SYLLIT FL and oils have been used effectively as a delayed dormant application on apples. However, SYLLIT FL showed to be incompatible with some oil formulations especially when oils were used at a high rate per acre. Because uniform dispersibility and sprayability may be influenced by the particular oil formulation used or by the local water supply, it is recommended that compatibility be determined before adding pesticides to the spray tank.

## **PROCEDURE FOR TANK-MIXING SYLLIT FL WITH INSECTICIDES AND OIL**

1. SYLLIT FL should be added to the tank when 1/4 to 1/2 full with water and allowed to mix thoroughly.
2. If insecticide is used, add as tank is filling.
3. Add self-emulsifying oil last when tank is nearly 3/4 full.

Consult your state agricultural experiment station or state extension service for information on self-emulsifying oils and pesticide formulations for tank mixing with SYLLIT FL.

**SPRAY DRIFT MANAGEMENT**

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity), and method of application (e.g., ground, aerial airblast) can influence pesticide drift. The applicator and grower must evaluate all factors and make appropriate adjustments when applying this product

**WIND SPEED**

Do not apply at wind speeds greater than 10 mph at the application site.

**DROPLET SIZE**

Apply as a medium or coarser spray (ASAE standard 572).

**AIRBLAST**

Sprays must be directed into the crop canopy. Turn off outward pointing nozzles at row ends and when spraying outer rows.

**AERIAL APPLICATION**

**Release Height** – Do not release spray at a height greater than 10 feet above the ground or crop canopy.

**Boom Length** – The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

**Swath Adjustment** – When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for the displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind. Leave at least one swath unsprayed at the downwind edge of the treated field.

**GROUND APPLICATION**

**Release Height** – Do not release spray at a height greater than 4 feet above the ground or crop canopy.

**APPLES:**

**Restrictions:**

- Do not apply more than 9.1 lbs ai/acre/year (21 pints of Syllit FL per acre per year).
- Do not apply more than 2 applications per year at a rate of 1.95 lbs ai/acre/application (4.5 pints of Syllit FL per acre per year).
- Do not apply more than 4 applications per year at a rate of 1.3 lbs ai/acre/application (3 pints of Syllit FL per acre per year).
- Do not apply less than 7 days from the last application.
- Do not apply within 7 days of harvest.
- Do not use apple pomace for livestock feed.

**Scab - Protection Schedule**

**Ground Application:** Use 1.5 to 3 pints per acre of SYLLIT FL at 7 day intervals to maintain scab control pre-bloom through the first cover applications. Use 1.5 pint per acre of SYLLIT FL in subsequent cover sprays.

SYLLIT FL may be used in a delayed dormant oil schedule. Use 1.5 to 3 pints of SYLLIT FL with 8 gallons of oil of 60-75 viscosity per acre. For mixing instructions, refer to SYLLIT FL and OIL section on the label.

**Air Application:** Use 2.4 pints of SYLLIT FL per acre in 5 to 7 gallons of spray at 7 day intervals to maintain scab control from pre-bloom through the first cover sprays. Repeat application in subsequent cover sprays.

**Apple Scab - "After-Infection" Schedule**

After an infection period has occurred, use 4.5 pints per acre of SYLLIT FL during or following a rain from pre-bloom through first cover applications and in later cover sprays. While timing of the spray may vary considerably depending upon temperatures, it is recommended that application be made within 36 hours, from the beginning of the rain or wet period which caused the infection.

Fruit color and finish on most apple varieties sprayed with SYLLIT FL at recommended dosages have frequently been outstanding.

SYLLIT FL may cause russetting of Golden Delicious and Grimes Golden apple varieties if applied during cool, slow drying weather conditions. Do not use on these varieties under these conditions.

Freezing or near-freezing temperatures, particularly when accompanied by slow-drying conditions may injure apple fruit or predispose the fruit to chemical injury. Do not exceed 3 pints of SYLLIT FL per acre immediately before, during or immediately after occurrence of such conditions.

**BANANA: Sigatoka (Air application)**

Use 1.5 to 3 pints per acre of SYLLIT FL as needed to maintain Sigatoka control through the first cover applications. Repeat application thereafter at 7 to 15 day intervals using 1.5 pints per acre of SYLLIT FL in a mixture of 9 to 18 pints water/acre and 5 to 7 pints mineral oil/acre. It is recommended for subsequent cover sprays to use SYLLIT FL in rotation with other labeled fungicides that have a different mode of action. Do not exceed more than 5 applications at 1.66 pints of SYLLIT FL per acre for a seasonal total of 8.3 pints of product or 3.6 lbs ai/acre/season. SYLLIT FL may be applied up to the day of harvest.

**CHERRIES (SOUR AND SWEET):**

**Restrictions:**

- Do not apply more than 5.2 lbs ai/acre/year (12 pints of Syllit FL per acre per year).
- Do not apply more than 2 applications per year at a rate of 1.3 lbs ai/acre/application (3 pints of Syllit FL per acre per year).
- Do not apply more than 4 applications per year at a rate of 0.65 lbs ai/acre/application (1.5 pints of Syllit FL per acre per year).
- Do not apply less than 7 days from the last application.
- Do not apply within 7 days of harvest.

**Leaf Spot:** Use 1.5 to 3 pints of SYLLIT FL per acre at petal fall or when first leaves unfold. Repeat application thereafter at 7 to 10 day intervals. If conditions favor leaf spot, continue applications through harvest.

**Post-Harvest Sprays:** Apply 1.5 to 3 pints of SYLLIT FL per acre immediately after harvest and as needed.

**Blossom Brown Rot (Western States Only):** Use 3 pints of SYLLIT FL per acre at the time petals begin to open. Repeat application at full bloom and again at petal fall.

Applications of SYLLIT FL to cherries when temperatures and humidity are excessively high may cause spotting to foliage or fruit.

**PEACHES:**

**Restrictions:**

- Do not apply more than 7.8 lbs ai/acre/year (18 pints of Syllit FL per acre per year).
- Do not apply more than 2 applications per year at a rate of 1.95 lbs ai/acre/application (4.5 pints of Syllit FL per acre per year).
- Do not apply more than 3 applications per year at a rate of 1.3 lbs ai/acre/application (3 pints of Syllit FL per acre per year).



- Do not apply less than 7 days from the last application.
- Do not apply within 15 days of harvest.

**Bacterial Spot** - Under light to moderate disease conditions, use 3 pints of SYLLIT FL and 4 lbs. captan (50%) per acre (use 0.8 pint SYLLIT FL and 1 lb. captan (50%) per 100 gallons of spray) at shuck-split and repeat application at 7 to 10 day intervals through the cover sprays. Pay attention to physical compatibility of tank-mix with some captan formulations. Shorten intervals during wet periods as needed to maintain maximum protection against infection.

Do not combine SYLLIT FL and captan with solvent formulations of insecticides or sulfur. Spray injury may result if the combination is applied with, immediately before, or closely after an oil or zinc sulfate-lime spray.

**Leaf Curl (Western States Only)** - Use 4.5 pints of SYLLIT FL per acre just before buds swell in the spring. If leaf curl was moderate to severe the previous season, use 4.5 pints SYLLIT FL per acre 1 month before buds swell in the spring and again just prior to bud swelling.

**Blossom Brown Rot (Western States Only)** - Use 3 pints of SYLLIT FL at the time petals begin to open. Repeat applications at full bloom and again at petal fall. Do not apply after petal fall.

**PEANUTS: Early Leaf Spot (*Cercospora arachidicola*) and Late Leaf Spot (*Cercosporidium personatum*)** – Use a maximum of 1.5 pints SYLLIT FL per acre in sufficient water to achieve thorough plant coverage. Begin applications when disease first appears (roughly 35 days after planting), and repeat at 10 to 14-day intervals. Do not use more than 3 applications per season. Do not apply within 14 days of harvest.

#### PEARS:

##### Restrictions:

- Do not apply more than 9.1 lbs ai/acre/year (21 pints of Syllit FL per acre per year).
- Do not apply more than 2 applications per year at a rate of 1.95 lbs ai/acre/application (4.5 pints of Syllit FL per acre per year).
- Do not apply more than 4 applications per year at a rate of 1.3 lbs ai/acre/application (3 pints of Syllit FL per acre per year).
- Do not apply less than 7 days from the last application.
- Do not apply within 7 days of harvest.

**Scab (Western States Only)** Use 4.5 pints of SYLLIT FL per acre of spray at 7 day intervals to maintain scab control from pre-bloom through petal fall and again in first cover. Use 3 to 4.5 pints of SYLLIT FL in subsequent spray.

A maximum of two applications may be made in the final two months prior to harvest: one made no later than 14 days prior to harvest and the final application made no later than 7 days prior to harvest.

**(Air Application - Oregon Only)** - Use 3 to 4.5 pints of SYLLIT FL per acre in 7 day intervals to maintain scab control from pre-bloom through the first cover sprays. Repeat application in subsequent cover sprays.

A maximum of two applications may be made in the final two months prior to harvest: one made no later than 14 days prior to harvest and the final application made no later than 7 days prior to harvest.

#### PECANS:

##### Restrictions:

- Do not apply more than 7.8 lbs ai/acre/year (18 pints of Syllit FL per acre per year).
- Do not apply more than 2 applications per year at a rate of 1.95 lbs ai/acre/application (4.5 pints of Syllit FL per acre per year).

- Do not apply more than 3 applications per year at a rate of 1.3 lbs ai/acre/application (3 pints of Syllit FL per acre per year).
- Do not apply less than 10 days from the last application.
- Do not apply after shucks have started to open.

**Scab, liver spot, brown leaf spot, downy leaf spot, leaf blotch, downy mildew - (Ground Application)** - Use 3 to 4.5 pints of SYLLIT FL per acre of spray. Begin applications when first leaves are showing and repeat at 10 to 14 day intervals through first cover. Repeat applications at 2 to 3 week intervals to maintain disease control.

**(Air Application)**- Use 3 pints of SYLLIT FL per acre in a minimum of 20 gallons of spray. Begin applications when first leaves are showing and repeat at 10 to 14 day intervals through first cover. Repeat application at 2 to 3 week intervals to maintain disease control. Under severe scab conditions, use 4.5 pints of SYLLIT FL per acre in a minimum of 20 gallons of spray.

SYLLIT FL has been used effectively on most pecan varieties. Severe foliage injury has been experienced on Barton, Moore, Shawnee and Van Deman in the Southeast.

**WALNUTS:**

**Restrictions:**

- Do not apply more than 7.8 lbs ai/acre/year (18 pints of Syllit FL per acre per year).
- Do not apply more than 2 applications per year at a rate of 1.95 lbs ai/acre/application (4.5 pints of Syllit FL per acre per year).
- Do not apply more than 3 applications per year at a rate of 1.3 lbs ai/acre/application (3 pints of Syllit FL per acre per year).
- Do not apply less than 10 days from the last application.
- Do not apply within 7 days of harvest.

**Anthracoese (Leaf Blight)** - Use 4.5 pints of SYLLIT FL per acre when first leaves unfold. Repeat application at 10 to 14 day intervals.

**IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of AGRIPHAR S.A.. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** AGRIPHAR S.A. MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of AGRIPHAR is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, AGRIPHAR S.A. DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN

11  
11

CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT AGRIPHAR'S ELECTION, THE REPLACEMENT OF PRODUCT.

Agriphar S.A.,  
Rue De Renory,26  
B-4102 Ougree, Belgium

SYLLIT is a registered trademark of Agriphar S.A.

*SYLLIT FL (MASTER), consolidated label - approved by EPA/SRRD on 08/07/2008 and EPA/RD on 03/27/2009 with comments incorporated;  
Submission on 05-18-2009*