

JUN 1 1994

Dr. Alice Walker  
Continental Sulfur Co.  
7500 San Felipe  
Suite 410  
Houston, Texas 77063

Dear Dr. Walker:

Subject: CSC Wettable or Dusting Sulfur  
EPA Reg. No. 55429-1  
Your Submission of February 4, 1994

The amendment referred to above, which supersedes your submission of August 30, 1993, submitted in connection with registration under FIFRA sec. 3(c)(7)(A) is acceptable provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) or section 4(a) when the Agency requires all registrants of similar products to submit such data.

2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling.

a. Under Environmental Hazards, replace the first two sentences with the following: "Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters."

b. Certain entries need clarification as to the type of application intended. For example, the entry for blackberries, boysenberries, gooseberries and loganberries does not indicate whether this is a spray or dust application. Specify the type of application for this entry and the following entries: blueberries, currants, dewberries, huckleberries; dates; mangoes; pistachios; raspberries; walnuts; cucumbers, squash; table beets; corn, barley oats, rye, sorghum, soybeans, wheat; and spearmint, peppermint."

20411

c. Add the statement, "Do not apply this product through any type of irrigation system" or add labeling required by PR Notice 87-1.

d. In the WPS labeling, under personal protective equipment, add protective eyewear since your certification indicated Toxicity Category II for eye irritation potential and this is also a requirement imposed by the Reregistration Eligibility Document.

e. Reinstate the section "Physical Hazards" which was included on your WPS label amendment of August 3, 1993.


f. This label does not bear direction for any non-agricultural uses. Delete the section "Non-agricultural Use Requirements".

3. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

  
Steve Robbins  
Acting Product Manager (21)  
Fungicide-Herbicide Branch  
Registration Division (7505C)

Enclosure

7505C:C.Grable:cq:5/31/94

30411

# CSC WETTABLE OR DUSTING SULFUR

FUNGICIDE - MITICIDE  
FOR SPRAYING OR DUSTING

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

JUN 01 1994

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

55429-1

## ACTIVE INGREDIENT:

Sulfur.....90.0%

INERT INGREDIENTS: .....10.0%

Total 100.0%

KEEP OUT OF REACH OF CHILDREN

## CAUTION

### STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with soap and water. Get medical attention if irritation persists.

IF IN EYES: Flush eyes with flowing water for at least 15 minutes. Get medical attention if irritation persists.

See side/back panel for additional "Precautionary Statements."

EPA Reg. No. 55429-1

EPA Est. No. 55429-TX-1

Manufactured by:  
CONTINENTAL SULFUR COMPANY  
7500 San Felipe, Suite 410  
Houston, Texas 77063

Net Contents  
\_\_\_\_\_ Lbs.

BEST AVAILABLE COPY

## PRECAUTIONARY STATEMENTS

### CAUTION

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

May be harmful if swallowed, inhaled, or absorbed through the skin. Avoid inhalation of vapors, spray or dust. Avoid contact with eyes, skin and clothing.

#### PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

#### ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply directly to any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply where run-off is likely to occur.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**BEST AVAILABLE COPY**

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. Exception: If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is coveralls, waterproof gloves, and shoes plus socks.

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store product in its original labeled container in a cool, dry, locked place out of reach of children.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**BEST AVAILABLE COPY**

## TO USE AS A SPRAY

**HOW TO MIX** - Select a suitable hand sprayer or power unit which uses a ground rig or aerial applicator. Measure the recommended amount for the specific spray problem separately into water and stir to make a thin "slurry" paste. Fill the spray tank 1/2 full and then add the sulfur "slurry" to the tank before other materials are added. When using a suitable "approved" wetting agent, add it to the tank last to avoid possible foaming. Keep spray mixture agitated while spraying. Mix only the amount of spray needed for the job.

## TO USE AS A DUST

Choose a suitable Hand or Power Duster as needed, or select a Certified Ground or Aerial Applicator who uses specialized dusting equipment designed to deposit the dust on all plant surfaces. For best results apply when wind velocity is below 5 mph. Observe precautions concerning use on sulfur sensitive plants and temperatures at the time of application.

**WHEN TO USE** - Apply to crops as needed in season when mites or diseases first appear and when insect "buildup" and plant disease damage is imminent. Repeat as directed for continued control. Contact your State Agricultural Extension Service for crop spray or dusting schedules which give the proper timing for applications in your locality.

## HOW MUCH SPRAY OR DUST TO USE

### WHEN TO APPLY THE LOWER RECOMMENDED RATE

1. During periods of minimal to moderate disease or insect infestation.
2. For regular season Cover Sprays.
3. When temperatures are up in the 80's and low 90's.

### WHEN TO APPLY THE HIGHER RECOMMENDED RATE

1. When mites or the diseases are damaging the crop and when weather conditions are favorable for their continued damage.
2. When temperatures are below and will not exceed 90° F within the following 3 days after the application.

**AS A SPRAY** - The lower rates would be 2 or 3 lbs. of the 2 to 6 lbs/100 gallons rate recommendation. The higher rates would be 4 or 6 lbs/100 gallons.

**AS A DUST** - The lower rate would be 50 lbs. of the 50 to 80 lb. rate per acre. The higher rate would be 80 lbs.

## DIRECTIONS FOR DILUTION

### DILUTE APPLICATION

- Ground: Apply specified rate in 20 to 60 gallons of water per acre.  
 Orchard: Apply specified rate in 100 to 800 gallons of water per acre.

**BEST AVAILABLE COPY**

## CONCENTRATE APPLICATION

Ground: Apply specified rate in 5-10 gallons of water per acre.

Orchard: Apply specified rate in 20 to 100 gallons of water per acre.

## AERIAL APPLICATION

Ground: Apply specified rate in 3 to 20 gallons of water per acre.

Orchard: Apply specified rate in 10 to 20 gallons of water per acre.

## PRECAUTIONS

In order to obtain effective mite and disease control, both the upper and lower sides of plant foliage and fruit must be thoroughly covered with spray solution. Poor spray coverage will result in poor mite and disease control. When growing crops for "processing," consult the processor before applying sulfur. Do not use on any crop unless sulfur has been shown to be safe for that crop in your locality. Sulfur may burn foliage when temperatures exceed 90° F. Sulfur will cause severe fruit and leaf injury to sensitive plants. Use only on varieties known to be tolerant to sulfur. Do not apply within 14 days of oil spray for all crops except CITRUS. **FOR CITRUS do not apply within 21 days of an oil spray.**

## AS A SPRAY OR DUST FOR FRUIT AND NUTS

**ALMONDS, CHERRIES, PLUMS, PRUNES:** For BROWN ROT, POWDERY MILDEW, LEAF SPOTS, SILVER MITES, FLAT MITES, SPIDER MITES. **AS A SPRAY** use 4 to 6 pounds per 100 gallons of water for ground applications. Use 10 to 30 lbs. in 15 to 20 gallons water per acre for aerial application. Apply at bloom or early petal fall, repeat as necessary, usually 10 to 14 day intervals or after periods of wet weather for disease control. **AS A DUST** for BROWN ROT on Cherries use 40 to 50 lbs. per acre applied 2 to 3 weeks before harvest. **FOR BROWN ROT and PRUNE ROT** on Plums and Prunes use 30 to 35 lbs. per acre 5 weeks before harvest. Repeat each 7 days if wet weather prevails and disease threatens during the harvest period.

**APPLES AND PEARS:** For POWDERY MILDEW, SCAB, BLISTER MITES, SILVER MITES & SPIDER MITES. **AS A SPRAY** use 4 to 6 lbs. per 100 gallons water for ground applications. Use 10 to 30 lbs. in 15 to 20 gallons water per acre for aerial application. Apply Pre-Bloom and Petal Fall. Apply with cover sprays as needed throughout the season. **AS A DUST** for SCAB use 30 to 35 lbs. per acre at "Green Tip" and again at "Pink Bud." Pink Bud treatments will also control POWDERY MILDEW. Do not apply during or shortly before frost conditions. **PRECAUTION:** Certain varieties of apples and pears are sensitive to sulfur sprays under certain conditions. Do not apply unless the varieties are known to be tolerant of sulfur.

**AVOCADO:** **AS A SPRAY** for BROWN MITES apply 2 to 6 lbs. per 100 gallons of water for ground applications. Use 10 to 30 lbs. in 15 to 20 gallons water per acre for aerial application. Apply at first sign of mite build up and repeat for control as needed or at 10 to 14 day intervals. See Re-Entry Statement.

**BLACKBERRY:** (Rubus spp.) For RED BERRY MITE (Acalitus essigi) **AS A SPRAY** apply 3 to 5 pounds per 100 gallons of water. Start applications in early "bud swell"

**BEST AVAILABLE COPY**

and repeat applications as necessary for mite control. **AS A DUST** use 15 to 20 lbs./acre and apply as for spray.

**BLACKBERRIES, BOYSENBERRIES, GOOSEBERRIES, LOGANBERRIES:** To control POWDERY MILDEW. Apply 5 1/3 to 13 1/3 lbs. per acre. Begin when new shoots are 6 inches long and before blossoms open or when disease first appears. Repeat at 7 to 10 day intervals or as necessary.

**BLUEBERRIES, CURRANTS, DEWBERRIES, HUCKLEBERRIES:** For control of POWDERY MILDEW. Apply 4 1/2 to 9 lbs. per acre. Begin when new shoots are 6 inches long and before blossoms open or when disease first appears. Repeat at 7 to 10 day intervals or as necessary.

**CITRUS: (GRAPEFRUIT, LEMONS, LIMES, ORANGES, TANGELOS, TANGERINES, CITRUS CITRON, KUMQUATS, and hybrids of these).** For FLAT MITES, SILVER MITES, & SPIDER MITES. **AS A SPRAY** use 5 to 10 pounds per 100 gallons of water for ground applications. Use 10 to 30 lbs. in 15 to 20 gallons water per acre for aerial application. For Florida CITRUS RUST MITE use 25 lbs. per 500 gallons water equal to 40 to 50 lbs. sulfur per acre for ground applications. Use 40 to 50 lbs. in 15 to 20 gallons water per acre for aerial application. Limit to one application per season where supplemental rust mite control is needed between main sprays. Otherwise, apply in any spray NOT CONTAINING OIL as needed for control during season or within 3 weeks of an oil spray as fruit burn may occur. Apply when mites are present in damaging numbers and repeat as needed or at 10 to 14 day intervals. See Re-Entry Statement.

**CITRUS: AS A DUST** for SIX SPOTTED MITE and CITRUS RUST MITE use 50 to 80 lbs. per acre. For CITRUS FLAT MITE use 80 to 100 lbs. per acre as needed. For YUMA MITE use 100 lbs. per acre. For CITRUS THRIP use 100 lbs. per acre in Mid March, Petal Fall and again 2 to 4 weeks later. Do not apply within 2 months of an oil spray. See Re-Entry Statement.

**DATES:** To control BANKS GRASS MITES. Use 44 1/2 to 111 1/4 lbs. per acre, applying by ground or air. Use the higher rates by air. Thorough coverage is essential for good control.

**FIG: AS A DUST** for FIG RUST MITE use 40 lbs. per acre in spring and again in the early summer.

**GRAPES:** For POWDERY MILDEW, BUD MITES, BLISTER MITES, & SPIDER MITES. **AS A SPRAY** use 2 to 6 pounds per 100 gallons of water. Apply in late dormant or with 1.) "cutworm spray;" 2.) in early shoot period; 3.) then through season at 14 day "cover spray" intervals or with or without pesticides. **AS A DUST** for POWDERY MILDEW use 15 to 20 lbs. per acre when shoots are 6, 12, & 18 inches long. Thereafter, repeat each 10 to 14 days to harvest. For RUST apply 15 to 20 lbs. per acre. Begin when new shoots are 6 inches long. Repeat before blossoms open and continue as necessary. NOTE: Concord and Labrusca type grapes may be injured by sulfur. Do not apply if temperatures during or shortly after treatment are expected to exceed 100° F. See Re-Entry Statements.



**MANGO:** For POWDERY MILDEW. Apply 9 to 26 1/2 lbs. per acre. Begin before bloom. Repeat in bloom, after fruit set and 3 weeks later or as necessary.

**OLIVES:** For BUD MITES. **AS A SPRAY** use 2 pounds per 100 gallons of water. Apply at the first sign of mite build up and repeat for control as needed or at 10 to 14 day intervals.

**PEACHES, NECTARINES:** For BROWN ROT, POWDERY MILDEW, SILVER MITES, & SPIDER MITES. **AS A SPRAY** use 4 to 6 pounds per 100 gallons of water for ground applications. Use 10 to 30 lbs. in 15 to 20 gallons water per acre for aerial application. Apply at bloom or early petal fall. Repeat as necessary, usually 10 to 14 day intervals or after periods of wet weather for disease control. **AS A DUST** for POWDERY MILDEW use 50 lbs. per acre at petal fall, 2 weeks later and again at beginning of pit hardening. For RUST and POWDERY MILDEW use 50 lbs. after pit hardening. For SILVER MITE use 25 to 35 lbs. per acre after thinning. For BROWN ROT use 50 to 60 lbs./acre 4 weeks pre-harvest and again 2 weeks later.

**PISTACHIOS:** For the aid of control of MITES. Use 44 1/2 lbs. per acre, making ground or air applications. This product may be applied to non-bearing and bearing pistachio trees up to the day of harvest.

**RASPBERRIES:** For POWDERY MILDEW. Apply 5 1/3 to 13 1/3 lbs. per acre. Spray each week starting at first bloom color. Continue until all fruit is set, then repeat as weather requires.

**STRAWBERRIES:** For POWDERY MILDEW and RED SPIDER MITES as a spray apply 3 to 5 lbs. per 100 gallons of water. Start in early leaf stage and repeat applications at 10 to 14 day intervals or as necessary for control with or without pesticide sprays. **AS A DUST** use 10 to 20 lbs. per acre preharvest. See When to Use and Re-entry Statement.

**WALNUTS:** For PACIFIC MITE, ALMOND MITE, TWO-SPOTTED MITE, RED SPIDER MITE. Apply 9 to 17 3/4 lbs. per acre and repeat as necessary. Apply with mite sprays during season to increase the effectiveness. **PRECAUTION:** Certain varieties of walnuts are sensitive to sulfur sprays under certain conditions. Do not apply unless the varieties are known to be tolerant of sulfur.

#### **AS A SPRAY OR DUST FOR VEGETABLES**

**BEANS, PEAS, BROCCOLI, CABBAGE, CAULIFLOWER, COLLARDS, KALE, AND PEPPERS** for POWDERY MILDEW and on **BEANS, PEAS AND PEPPERS** for SPIDER MITES. **AS A SPRAY** use 3 to 5 pounds per acre in (15 to 50 gallons water) aerial or ground application respectively. Start in the early leaf stage. Repeat application every 10 to 14 days with or without pesticides or immediately after a rain. **AS A DUST** for POWDERY MILDEW & SPIDER MITES use 15 to 20 lbs./acre. Do not apply to varieties susceptible to sulfur injury. See **WHEN TO USE & RE-ENTRY STATEMENT**.

**BEST AVAILABLE COPY**

**CARROTS, ONIONS AND GARLIC:** For control of PETROBIA MITE. **AS A DUST** use 25 to 30 lbs. per acre as needed for control. See **WHEN TO USE**.

**CUCUMBERS, SQUASH:** For POWDERY MILDEW. Apply 25 lbs. per acre when disease first appears and continue as necessary. **PRECAUTION:** Certain varieties of cucumbers and squash are sensitive to sulfur sprays under certain conditions. Do not apply unless the varieties are known to be tolerant of sulfur.

**MELONS:** **AS A DUST** for POWDERY MILDEW use 15 to 20 lbs. per acre. For SPIDER MITES use 20 to 40 lbs /acre. Do not apply to varieties susceptible to sulfur injury. Melon foliage is extremely sensitive to sulfur burn. Observe use precautions and apply only when temperatures are well below 90° F. See paragraphs on **WHEN TO USE** and **HOW MUCH SPRAY OR DUST TO USE**.

**POTATOES:** **AS A SPRAY** for control of POWDERY MILDEW use 3 to 4 lbs. in sufficient water to cover 1 acre. Applying by aircraft use a minimum of 8 gallons of spray per acre. With ground equipment use a minimum of 10 gallons of spray per acre. Apply at first sign of infection and repeat at three week intervals.

**TABLE BEETS:** For POWDERY MILDEW, RED SPIDER MITE. Apply 2 2/3 to 4 1/2 lbs. per acre at 10 to 30 day intervals with or without other pesticides as soon as the first symptoms appear.

**TOMATOES:** **AS A SPRAY** for RUSSET MITES apply 3 to 5 pounds in sufficient water (15 gals. minimum for aerial application and 50 gallons minimum for ground rig) to treat an acre. Thorough coverage of upper and lower sides of leaves and fruit is required for effective control. Apply when mites are present in damaging numbers and repeat at 10 to 14 day intervals as needed. **AS A DUST** for control of RUSSET MITE use 30 to 40 lbs./acre. If crop is being grown with the aid of stakes or poles, apply 10 to 13 lbs. per acre per foot of average plant height. (Plants 4 ft. height = 40 to 52 lbs. per acre.)

#### **AS A SPRAY OR DUST FOR FIELD CROPS**

**CORN, BARLEY, OATS, RYE, SORGHUM, SOYBEANS, WHEAT:** For suppression of RED SPIDER MITE, TWO-SPOTTED MITE, GRASS BANKS MITE, and PACIFIC MITE; for control of ATLANTIC MITE (small grain only) and for control of POWDERY MILDEW (soybeans only). Apply 5 1/3 lbs. per acre. For soybeans, apply 6 2/3 to 11 1/8 lbs. per acre. Apply when disease first appears and repeat as necessary or with the regularly scheduled insect program.

**COTTON AND SEED ALFALFA:** **AS A SPRAY** for SPIDER MITES apply 3 to 5 pounds per acre in 20 to 50 gallons of water by aircraft or ground rig respectively. Apply when mites first appear to buildup on new growth and damage to plants is imminent. Thorough spray coverage must be made to the upper and lower portions of all foliage. **AS A DUST** on COTTON for the ATLANTIC MITE use 45 to 50 lbs. per acre when first observed in flower heads and before defoliation of more than the bottom leaves has occurred. Repeat as needed throughout seed-set. May cause some flower burn. Use low rate on immature plants.

**BEST AVAILABLE COPY**

9 110411  
**HOPS: AS A SPRAY** for control of SPIDER MITES, use 2 to 6 pounds per 100 gallons of water. Start applications in the early leaf stage and repeat as needed or at 10 to 14 day intervals.

**PEANUTS: AS A SPRAY** for POWDERY MILDEW, SPIDER MITES and LEAF SPOT diseases apply 3 to 5 pounds per acre in 15 to 50 gallons of water by aircraft or ground rig respectively. Start in the early leaf stage and repeat applications as necessary or at 10 to 14 day intervals.

**SPEARMINT, PEPPERMINT:** For POWDERY MILDEW. Apply 3 1/2 to 5 1/3 lbs. per acre when mint is 5-6" tall or when disease appears. Repeat twice at 30 day intervals. Do not apply within 30 days of harvest.

**SUGAR BEETS:** For POWDERY MILDEW & SPIDER MITES<sup>c</sup> **AS A SPRAY** apply 3 to 5 pounds per acre in 15 to 50 gallons of water by aircraft or ground rig respectively. Apply at first signs of insects or disease. Repeat applications should be made at 10 to 30 day intervals or as necessary throughout the season. Under most disease conditions only one or two applications will be needed. **AS A DUST:** For SPIDER MITES use 45 to 50 lbs. per acre. For POWDERY MILDEW use 25 to 50 lbs. per acre. For PUCCINIA RUST use 7-1/2 to 12-1/2 lbs. per acre. Apply at the first signs of disease. Repeat at 7 to 10 day intervals. This product may be applied by ground or air. Use the higher rate when applying by air. The use of this product in hot weather over 90° F may cause foliage burn. See paragraphs on HOW MUCH TO USE AND RE-ENTRY STATEMENT.

### ORNAMENTALS

**AS A SPRAY** use 2 to 6 pounds per 100 gallons of water and for use **AS A DUST** use 45 to 50 lbs. per acre for POWDERY MILDEW of Alder, Azalea, Buttonbush, Catalpa, Chrysanthemum, Dahlia, Delphinium, Honeysuckle, Horse Chestnut, Hydrangea, Lilac, Linden, Matrimony Vine, Phlox, Rhododendron, Rose, Staghorn Sumac, Sweetpea, Sycamore, Trumpet Vine, Witch-Hazel, Zinnia, RUSTS spp. of Aster, Carnation, Chrysanthemum, Rose, and Snapdragon. Make full coverage applications on both sides of leaves. Begin before disease appears. Repeat at 7 to 10 day intervals. See WHEN TO USE. Apply to plant foliage, stems and flowers to leave a "barely visible" dust residue. Do not over apply during full sun nor during times when temperatures exceed 90° F.

**NOTICE:** Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms on this label.

**BEST AVAILABLE COPY**