

55467-6

2/14/2008

1/5



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

FEB 14 2008

Mr. Robert E. Ridsdale
Tenkoz, Inc
100 North Point Center East, Suite 330
Alpharetta, GA 30022-8242

Dear Mr. Ridsdale:

Subject: Volley Atz Lite Corn Herbicide (Correct Restricted Use Box on Supplementals for Use
on Sweet Corn and Revision of Rotational Crop Intervals)
EPA Registration No. 55467-6
Application Dated January 16, 2008

The labeling referred to above, submitted in connection with registration under the Federal
Insecticide, Fungicide, and Rodenticide Act as amended is acceptable. Amended labeling
supersedes supplemental labeling approved December 20, 2007. A stamped copy of labeling is
enclosed for your records.

Sincerely,

James A. Tompkins
James A. Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505P)

2/5

ACCEPTED
FEB 14 2008
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended for the pesticides
registered under
EPA Reg. No. 55467-6

Supplemental Labeling

Tenkoz Inc.

Tenkoz Inc. 100 North Point Center East Alpharetta, GA 30022 USA

RESTRICTED USE PESTICIDE
Due to Ground and Surface Water Concerns

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

This product is a restricted use herbicide due to ground and surface water concerns. Users must read and follow all precautionary statements and instructions for use in order to minimize potential for atrazine to reach ground and surface water.

Volley® Atz Lite Corn Herbicide

EPA Reg. No. 55467-6

Weed Control in Sweet Corn

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for **Volley® Atz Lite Corn Herbicide** before applying. Follow all applicable use directions and precautions on the EPA-registered label.
- Use of **Volley Atz Lite** according to this supplemental labeling is subject to all restrictions and limitations imposed by the label affixed to the product container.

Directions for Use

Volley Atz Lite Applied Alone

Used alone, **Volley® Atz Lite Corn Herbicide** may be applied to sweet corn using rates stated in the product label for **Volley Atz Lite** and application methods and timing listed below.

Carriers and Spray Volume: Either water or liquid fertilizers such as solutions, slurries or suspensions may be used as liquid carriers. Refer to product label for **Volley Atz Lite** for details.

Application Timing and Methods: For the optimum period of effective weed control during the time most critical to corn production, preplant applications of **Volley Atz Lite** should occur as close as possible to planting. Preemergence applications should occur as close as possible to planting, but prior to weed emergence. Refer to product label for **Volley Atz Lite** for details.

Note: Do not apply **Volley Atz Lite** to sweet corn as an early postemergence application.

- **Early Preplant:** On medium and fine textured soils (see Table 1 in main label), **Volley Atz Lite** may be applied up to 30 days prior to planting.
- **Preplant Incorporation:** **Volley Atz Lite** and certain tank mixes may be mechanically incorporated in the top 2 inches of the soil with field cultivators, discs, or spring tooth harrows at any time within 14 days prior to planting.
- **Preemergence Surface:** **Volley Atz Lite** and certain tank mixes may be applied to the soil surface as a broadcast or banded application.

- **Postplant-Preemergence: Volley Atz Lite** may be applied immediately after planting but prior to corn emergence.
- **Banding Preemergence: Volley Atz Lite** may be applied in a 10 to 14 inch band after corn planting but prior to corn emergence.

Sprinkler Irrigation: Do not apply Volley Atz Lite by sprinkler irrigation. Use a sprinkler system only to incorporate **Volley Atz Lite** after application. Refer to product label for **Volley Atz Lite** for details.

Cultivation: Cultivation should be delayed as long as possible. If weeds develop, a shallow cultivation or rotary hoeing will generally result in improved weed control. Refer to product label for **Volley Atz Lite** for details.

Soil Texture and Organic Matter: The use rate of **Volley Atz Lite** is determined by a combination of two factors, soil texture and organic matter, which must be determined prior to application. Refer to Table 1 in main product label for soil texture classifications.

Use Rates for Conventional Tillage Systems: Refer to table 2 in main label for application rates and detailed information for conventional systems.

Use Rates for Reduced Tillage Systems: Refer to table 3 in main label for application rates and detailed information for reduced tillage systems.

Weeds Controlled: Refer to table 4 in main label for details.

Volley Atz Lite Applied in Tank Mix Combination

Additional weeds may be controlled with tank mixes of **Volley Atz Lite** and other products labeled for use on sweet corn. Tank mix combinations may be used in conventional, reduced, or no-till systems and may be applied using application methods and timing listed above unless otherwise specified in the tank mix product label. Refer to the product label for **Volley Atz Lite** for details on carriers, mixing instructions and testing for compatibility of tank mixtures.

®Trademark of Dow AgroSciences LLC

EPA Accepted:
Initial printing.

4/5

ACCEPTED
FEB 14 2008
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended for the pesticide registered under EPA Reg. No. 55467-6

Supplemental Labeling

Tenkoz Inc.

Tenkoz Inc. 100 North Point Center East Alpharetta, GA 30022 USA

RESTRICTED USE PESTICIDE
Due to Ground and Surface Water Concerns
For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.
This product is a restricted use herbicide due to ground and surface water concerns. Users must read and follow all precautionary statements and instructions for use in order to minimize potential for atrazine to reach ground and surface water.

Volley® Atz Lite Corn Herbicide

EPA Reg. No. 55467-6

Rotational Crop Restrictions After Final Application

- ATTENTION**
- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
 - This label must be in the possession of the user at the time of pesticide application.
 - Read the label affixed to the container for Volley® Atz Lite Corn Herbicide before applying. Follow all applicable use directions and precautions on the EPA-registered label.
 - Use of Volley Atz Lite according to this supplemental labeling is subject to all restrictions and limitations imposed by the label affixed to the product container.

Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

Rotational Crop Restrictions

When tank mixing with other herbicides, follow the most restrictive crop rotation guidelines on the label of each product used. The following rotational crops may be planted as indicated:

| Crop | Interval |
|---|--------------------------------------|
| Corn (1) | Anytime – 0 Months After Application |
| Sorghum, Soybean (2) | Spring Following Application |
| Alfalfa, Barley, Dry beans (3), Lupin (4), Millet, pearl or proso, Oats, Pea (6), Potatoes, Rye, Sugar beets Sunflower, Tobacco (7), Triticale, Wheat, Wild rice | 15 Months After Application (5) |

Numbers within parentheses (-) in the table refer to Specific Rotational Crop Requirements or crop descriptions below.

- (1) If crop treated with Volley Atz Lite is lost, corn may be replanted immediately. Do not make a second application of Volley Atz Lite. Do not apply Volley Atz Lite after June 10, unless only corn will be planted the following year.
- (2) Due to the risk of atrazine carryover, injury to soybeans may occur the year following corn when planted in north central and northwest Iowa, south central and southwest Minnesota, northern Nebraska and southeast South Dakota on soils having a calcareous surface layer and relatively high pH.
- (3) Dry beans includes: adzuki, kidney, lima, navy, pinto
- (4) Lupin includes: grain, white, white sweet
- (5) Approved rotation crops list does not include any species of succulent beans and peas.
- (6) Pea includes: blackeyed, chick, cow, Crowder, field, pigeon, Southern
- (7) Because of atrazine carryover, injury may occur to tobacco.

5/5

®Trademark of Dow AgroSciences LLC

EPA Accepted:
Initial printing.