

PLEASE NOTE

This image contains more than one label approved for this product on this date.

55467-8

01/07/2008

1/27



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

JAN -7 2008

Mr. Robert E. Ridsdale
Tenkoz, Inc.
100 North Point Center East, Suite 330
Alpharetta, GA 30022

Dear Mr. Ridsdale:

Subject: Volley Corn Herbicide (Revise Master Label)
EPA Registration No. 55467-8
Application Dated October 10, 2007

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, is acceptable. Amended labeling supersedes all previously approved ones. A stamped copy of labeling is enclosed for your records.

Sincerely,

Vicki K. Williams for
James A. Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505P)

(Base label for rigid containers 5 gal or less):

Volley[®] Corn Herbicide


Herbicide

55467-8

For use only on field corn, production seed corn, silage corn, sweet corn and popcorn

Active Ingredients:

acetochlor: 2-chloro-2'-methyl-6'-ethyl-N-ethoxymethylacetanilide	70.87%
Other Ingredients:	29.13%
Total	100.00%



Contains 6.4 lb active ingredient per gallon

**Keep Out of Reach of Children
WARNING AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Skin And Eye Irritation • Harmful If Swallowed, Inhaled, Or Absorbed Through Skin

Do not get on skin, in eyes, or on clothing. Avoid breathing spray mist.

Personal Protective Equipment (PPE)

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing, or loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

Immediately start the procedures below. If further treatment is required, contact a Poison Control Center, a physician or the nearest hospital.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

Environmental Hazards

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the groundwater is shallow, may result in groundwater contamination.

Acetochlor has properties that may result in surface water contamination via dissolved runoff and runoff erosion. Practices should be followed to minimize the potential for dissolved runoff and/or runoff erosion.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with vermiculite, earth, or synthetic absorbent.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

4/27

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of label booklet. If terms are not acceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-424-9300. If you wish to obtain additional product information, visit our web site at www.tenkoz.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 55467-8

EPA Est. _____

®Trademark of Dow AgroSciences LLC

Tenkoz Inc. • 100 North Point Center East, Alpharetta, GA 30022 U.S.A.

Net Contents ____

(Base label for refillable rigid containers larger than 5 gal):

Volley[®] Corn Herbicide

Herbicide

For use only on field corn, production seed corn, silage corn, sweet corn and popcorn

Active Ingredients:

acetochlor: 2-chloro-2'-methyl-6'-ethyl-N-ethoxymethylacetanilide	70.87%
Other Ingredients:	29.13%
Total	100.00%

Contains 6.4 lb active ingredient per gallon

Keep Out of Reach of Children

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Skin And Eye Irritation • Harmful If Swallowed, Inhaled, Or Absorbed Through Skin

Do not get on skin, in eyes, or on clothing. Avoid breathing spray mist.

Personal Protective Equipment (PPE)

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing, or loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

Immediately start the procedures below. If further treatment is required, contact a Poison Control Center, a physician or the nearest hospital.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

Environmental Hazards

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the groundwater is shallow, may result in groundwater contamination.

Acetochlor has properties that may result in surface water contamination via dissolved runoff and runoff erosion. Practices should be followed to minimize the potential for dissolved runoff and/or runoff erosion.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with vermiculite, earth, or synthetic absorbent.

Container Reuse: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

7/27

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of label booklet. If terms are not acceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-424-9300. If you wish to obtain additional product information, visit our web site at www.tenkoz.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 55467-8

EPA Est. _____

®Trademark of Dow AgroSciences LLC

Tenkoz Inc. • 100 North Point Center East, Alpharetta, GA 30022 U.S.A.

Net Contents _____

(Base label for nonrefillable rigid containers larger than 5 gal):

Volley® Corn Herbicide

Herbicide

For use only on field corn, production seed corn, silage corn, sweet corn and popcorn

Active Ingredients:

acetochlor: 2-chloro-2'-methyl-6'-ethyl-N-ethoxymethylacetanilide	70.87%
Other Ingredients:	29.13%
Total	100.00%

Contains 6.4 lb active ingredient per gallon

Keep Out of Reach of Children

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Skin And Eye Irritation • Harmful If Swallowed, Inhaled, Or Absorbed Through Skin

Do not get on skin, in eyes, or on clothing. Avoid breathing spray mist.

Personal Protective Equipment (PPE)

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing, or loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

Immediately start the procedures below. If further treatment is required, contact a Poison Control Center, a physician or the nearest hospital.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

Environmental Hazards

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the groundwater is shallow, may result in groundwater contamination.

Acetochlor has properties that may result in surface water contamination via dissolved runoff and runoff erosion. Practices should be followed to minimize the potential for dissolved runoff and/or runoff erosion.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with vermiculite, earth, or synthetic absorbent.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

10/27

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of label booklet. If terms are not acceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-424-9300. If you wish to obtain additional product information, visit our web site at www.tenkoz.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 55467-8

EPA Est. _____

®Trademark of Dow AgroSciences LLC

Tenkoz Inc. • 100 North Point Center East, Alpharetta, GA 30022 U.S.A.

Net Contents _____

11/27

(cover)

Volley[®] Corn Herbicide

Herbicide

For use only on field corn, production seed corn, silage corn, sweet corn and popcorn

Active Ingredients:

acetochlor: 2-chloro-2'-methyl-6'-ethyl-N-ethoxymethylacetanilide 70.87%

Inert Ingredients: 29.13%

Total 100.00%

Contains 6.4 lb active ingredient per gallon

Keep Out of Reach of Children

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Agricultural Use Requirements
Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies at end of label booklet. If terms are not acceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-424-9300. If you wish to obtain additional product information, visit our web site at www.tenkoz.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 55467-8

EPA Est. _____

®Trademark of Dow AgroSciences LLC

Tenkoz Inc. • 100 North Point Center East, Alpharetta, GA 30022 U.S.A.

Net Contents _____

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals**WARNING****Causes Skin And Eye Irritation • Harmful If Swallowed, Inhaled, Or Absorbed Through Skin****Do not get on skin, in eyes, or on clothing. Avoid breathing spray mist.****Personal Protective Equipment (PPE)**

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing, or loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

Immediately start the procedures below. If further treatment is required, contact a Poison Control Center, a physician or the nearest hospital.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

Environmental Hazards

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the groundwater is shallow, may result in groundwater contamination.

Acetochlor has properties that may result in surface water contamination via dissolved runoff and runoff erosion. Practices should be followed to minimize the potential for dissolved runoff and/or runoff erosion.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. **Exception:** If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves made of any waterproof material
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with vermiculite, earth, or synthetic absorbent.

Nonrefillable containers 5 gallons or less:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers larger than 5 gallon:

Container Reuse: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable containers larger than 5 gallons:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

General Information

Volley® Corn Herbicide is intended for preplant, preemergence, or early postemergence use in corn. Use of this product in corn is limited to field corn, production seed corn, silage corn, sweet corn and popcorn. Do not apply this product to any crop other than corn.

15/27

Volley is a unique combination of the herbicide acetochlor and the antidote or safener dichlormid. While the acetochlor provides weed control, the dichlormid safens corn against herbicide injury. Volley may be applied to the surface or incorporated into the top 1 to 2 inch layer of soil. It is recommended for control alone, or in tank mix combinations as indicated, for the weeds listed in the Target Weeds section of these use directions. Volley controls weeds by interfering with normal germination and seedling development. Volley will not control established or germinated weeds present at application.

General Use Precautions and Restrictions

- Do not apply to the following soils if groundwater depth is 30 feet or less: sand with less than 3% organic matter; loamy sand with less than 2% organic matter; or sandy loam with less than 1% organic matter.
- Do not apply this product using aerial application equipment.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- Do not use flood irrigation to apply or incorporate this product.
- **Maximum Application Rate:** The maximum application rate for Volley on corn is 3.75 pints (3 lb of acetochlor active ingredient) per acre per year.
- This product may not be mixed or loaded within 50 feet of any wells including abandoned wells and drainage wells, sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas.
- Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or washwater, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.
- Product must be used in a manner that will prevent back siphoning in wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.
- Do not apply under conditions that favor runoff or wind erosion of soil containing this product to non-target areas. To prevent off-site movement due to runoff or wind erosion:
 - Avoid treating powdery dry or light sandy soils when conditions are favorable for wind erosion. Under these conditions, the soil surface must first be settled by rainfall or irrigation.
 - Do not apply to impervious substrates such as paved or highly compacted surfaces or frozen or snow covered soils.
 - Do not use tailwater from the first flood or furrow irrigation of treated fields to treat non-target crops unless at least 1/2 inch of rainfall has occurred between application and the first irrigation.
- Do not apply when wind conditions favor drift to non-target sites. To minimize spray drift to non-target areas:
 - Use low pressure application equipment capable of producing a large droplet spray.
 - Do not use nozzles that produce a fine droplet spray.
 - Minimize drift by using sufficient spray volume to ensure adequate coverage with large droplet size sprays.
 - Keep ground-driven spray boom as low as possible above the target surface.

16/27

- Make application when the wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid application when gusts approach 15 mph.
- Low humidity and high temperatures increase the likelihood of spray drift to sensitive areas. Avoid spraying during conditions of low humidity and/or high temperatures. Do not apply during inversion conditions.
- Do not apply more than 3 3/4 pints of Volley per acre per season.

Rotational Crop Restrictions:

When tank mixing with other herbicides, follow the most restrictive crop rotation guidelines on the label of each product used. The following rotational crops may be planted as indicated:

Rotational Crop	Timing or Interval
corn (1)	Anytime - 0 months after application
alfalfa, barley, buckwheat, clover, dry beans (2), guar, kudzu, lentil, lespedeza, lupin (4), millet, pearl or proso, oats, pea (5), potatoes, rye, sorghum, soybeans, sugar beets, sunflower, trefoil, tobacco, triticale, vetch, wild rice	Spring following application (3)
wheat	4 months after application

Numbers within parentheses (-) in the table refer to Specific Rotational Crop Requirements below.

1. If crop treated with Volley is lost, corn may be replanted immediately. Do not make a second application of Volley.
2. Dry beans includes: adzuki, kidney, lima, navy, pinto
3. Approved rotation crops list does not include any species of succulent beans and peas
4. Lupin includes: grain, white, white sweet
5. Pea includes: blackeyed, chick, cow, Crowder, field, pigeon, Southern

Application Directions - Corn

Carriers and Spray Volume

Liquids: Either water or liquid fertilizers such as solutions, slurries, or suspensions may be used as liquid carriers. If fluid fertilizers are used, a physical compatibility with these must be done **before combining** in the spray tank. See Appendix I for details of the compatibility testing procedure. Even if Volley is physically compatible with a fluid fertilizer, constant agitation is necessary to maintain a uniform mixture during application.

Apply in a minimum broadcast spray volume of 10 gallons per acre using boom equipment for ground applications. Use low pressure nozzles designed for application of herbicides. Use sufficient operating pressure to produce the desired spray pattern for the nozzle (15 to 40 psi) and follow manufacturer's recommendations for nozzle spacing and operating height to ensure uniform spray distribution at the soil surface. Use 50-mesh or coarser screens, if needed.

Dry Bulk Fertilizer: Volley may be impregnated on dry bulk fertilizer and applied as the fertilizer is spread. Use at least 200 lb of dry bulk fertilizer per acre. See Appendix II for more details including which fertilizers are compatible.

Adding to Spray Tank

The spray tank must be clean, thoroughly rinsed, and decontaminated before adding either Volley alone or in tank mix combinations. If water is used as the carrier, use clean water. All return lines to the spray tank must discharge below the liquid level.

17/27

Used Alone: If Volley is used alone, add the recommended amount to the spray tank before the tank is half filled, then add the rest of the water or fluid fertilizer. Provide sufficient agitation to ensure thorough mixing and to maintain a uniform spray mixture during application.

Tank Mixed: If a tank mixture is used, it is recommended that a small-scale test of compatibility be done before actual tank mixing. See Appendix I for details on the procedure for such a test.

Water Carrier

Allow time for complete dispersion/mixing before adding another product to the spray mixture. Add products to the tank mixture in the following order:

- To start, add one-half of the required amount of water to the spray tank. Begin agitation.
- Products in water soluble packaging. Important: Allow time for complete dispersion.
- Wettable powders or dry flowables (slurry if recommended by tank mix product label)
- Liquid flowables
- Volley or other emulsifiable concentrates
- Suspension concentrates
- Urea ammonium nitrate (UAN) or ammonium sulphate (AMS), if required
- Compatibility agent if needed
- Soluble liquids such as glyphosate, paraquat, 2,4-D amine
- Crop oil concentrate (COC) or nonionic surfactant (NIS), if required
- Finish filling spray tank to required spray volume

Liquid Fertilizer Carrier

Allow time for complete dispersion/mixing before adding another product to the spray mixture. Add products to the tank mixture in the following order:

- To start, add one-half of the required amount of liquid fertilizer to the spray tank. Begin agitation.
- Compatibility agent if needed
- Products in water soluble packaging. Important: Products in water soluble packaging must be premixed with water (slurried) prior to addition to the spray tank.
- Wettable powders or dry flowables (slurry if recommended by tank mix product label)
- Liquid flowables
- Volley or other emulsifiable concentrates
- Suspension concentrates
- Ammonium sulphate (AMS), if tank mixing with glyphosate.
- Soluble liquids such as glyphosate, paraquat, 2,4-D amine
- Crop oil concentrate (COC) or nonionic surfactant (NIS), if required
- Finish filling spray tank to required spray volume.

Note: For all tank mixtures, maintain agitation during mixing and throughout application to ensure spray mixture remains uniformly suspended.

Application Timing and Methods

For the optimum period of effective weed control during the time most critical to corn production, preplant application of Volley should occur as close as possible to planting. Preemergence applications should occur as close as possible to planting, but prior to weed emergence. Postemergence applications should occur prior to weed emergence or in tank mix combination with a product that controls emerged weeds.

Note: Do not apply Volley to sweet corn as an early postemergence application.

Early Preplant: On medium and fine textured soils (see Table 1) Volley and certain tank mixtures may be applied up to 30 days before planting.

Preplant Incorporation: Volley and certain tank mixes may be mechanically incorporated into the top 2 inches of the soil with field cultivators, discs, or spring tooth harrows any time up to 14 days prior to

planting. Improper incorporation, excessive crop residues, or poor soil tilth may result in erratic, streaked, or otherwise unsatisfactory weed control. Do not mix Volley deeper than 2 inches into the soil and avoid moving or shaping soil after incorporation.

Preemergence Surface: Volley and certain tank mixes may be applied to the soil surface as a broadcast or banded application. Precipitation or sprinkler irrigation of at least 0.25 inch is required to bring Volley into contact with germinating seeds. If rain or sprinkler irrigation does not occur within 7 days after application, weed control may be improved by using a rotary hoe or similar equipment to incorporate the herbicide. Incorporation equipment should be run at a shallow depth to avoid disturbance of germinating corn seed. Erratic weed control resulting from exposure of untreated soil may occur if surface soil is moved or reshaped after incorporation.

Postplant-Preemergence: Volley may be applied after planting but prior to corn emergence. If rain or sprinkler irrigation does not occur within 7 days after application, weed control may be improved by using a rotary hoe or similar equipment to shallowly incorporate the herbicide. Incorporation equipment should be run at a shallow depth to prevent disturbance of the germinating corn. Erratic weed control resulting from exposure of untreated soil may occur if surface soil is moved or reshaped during incorporation.

Banding-Preemergence: Volley may be applied in a 10- to 14-inch band after corn planting but prior to emergence. If rain or sprinkler irrigation does not occur within 7 days after application, weed control may be improved by shallow incorporation using a rotary hoe or similar equipment. Do not disturb the germinating corn seed. Erratic weed control resulting from exposure of untreated soil may occur if surface soil is moved or reshaped during incorporation.

Early Postemergence: Volley may be applied early postemergence to corn up to 11 inches tall. Application must be made prior to weed seedling emergence or in a tank mixture that controls the emerged weeds. Read and follow restrictions and directions on tank mix product labels.

Sprinkler Irrigation: Do not apply Volley by sprinkler irrigation. Use a sprinkler system only to incorporate Volley after application. After Volley has been applied, a sprinkler irrigation system set to deliver 0.25 to 0.75 inch of water per acre may be used to incorporate the product. Using more than 0.75 inch of water could result in reduced performance. On sandy soil low in organic matter, use no more than 0.5 inch of water. Do not use flood irrigation to apply or incorporate Volley.

Fall Application - For use in IA; IL (North of Route 136); NE (North of Route 20); MN; ND; SD; WI: Following soybean harvest, apply to soybean stubble after October 15, when the sustained soil temperature at 4-inch depth is less than 50°F, but before ground freezes. Use on medium and fine textured soils with greater than 2.5% organic matter. Only corn may be planted the following spring.

Ground may be tilled before or after application. Do not exceed 2-inch incorporation depth if tilled after application.

If a spring application is made, the total rate of the fall plus spring application must not exceed the maximum labeled rate for corn grown on that soil.

Cultivation

Cultivation should be delayed as long as possible. If weeds develop, a shallow cultivation or rotary hoeing will generally result in improved weed control. If Volley was incorporated, cultivate to a depth of less than half the depth of incorporation.

If cultivation is necessary due to soil crusting or compaction, adjust equipment to run shallow and minimize soil movement. This will decrease the possibility of diluting or moving the herbicide from the weed control zone.

Soil Texture and Organic Matter

19/27

The use rate of Volley is determined by a combination of two factors, soil texture and organic matter, which must be determined prior to application. Different soil textures are grouped into three textural classes (coarse, medium, and fine) as outlined in Table 1. Soil texture and organic matter content of the soil may be determined from soil survey information and/or by laboratory analysis and must be known in order to select the proper rate from Table 2.

Table 1: Soil Texture Groupings for Volley Use Rate Selection.

Coarse	Medium	Fine
Sand Loamy Sand Sandy Loam	Loam Silt Silt Loam Sandy Clay Loam	Silty Clay Loam Silty Clay Sandy Clay Sandy Clay Loam Clay Loam Clay

Use Rates in Conventional Tillage Systems

Table 2: Use Rates for Volley by Soil Texture and Organic Matter Content in Conventional Tillage Systems.

The following use rates are for preplant incorporated, preemergence, and early postemergence applications (see Application Timing and Methods). Consult Table 3 if no-till applications are made or application is made more than 14 days prior to planting under conventional tillage.

Soil Texture	Soil Organic Matter Content		
	Less than 3%	3% or Greater	Greater than 7%
Coarse	1.5 - 2.25 pt/acre	1.5 - 2 1/2 pt/acre	2 - 3 pt/acre
Medium	1.5 - 2.5 pt/acre	1.5 - 2 1/2 pt/acre	2.5 - 3.75 pt/acre
Fine	1.5 - 2.75 pt/acre	2 - 3 pt/acre	3 - 3.75 pt/acre

Rate Ranges: If the weed infestation is light and/or organic matter is in the lower end of the range, use a rate at the lower end of the rate range for the soil texture and organic matter content. If the weed infestation is heavier and/or organic matter is in the upper end of the range, use the higher rates in the rate range for the soil texture and organic matter.

Use Rates for Reduced Tillage Systems

Volley may be used in reduced and no-till systems. Application may occur up to 30 days prior to planting or after planting but before corn emergence. Optimal weed control will be obtained when applications are made as close as possible to planting but before the weeds emerge. In reduced or no-till systems, it is recommended that a burndown herbicide such as paraquat (Gramoxone) or glyphosate (Glyphomax, Roundup or Touchdown) or 2,4-D be tank mixed with Volley if emerged weeds are present at application.

Table 3: Use Rates for Volley by Soil Texture and Organic Matter Content in Reduced and No-Till Systems.¹

Soil Texture	Soil Organic Matter Content		
	Less Than 3%	3% or Greater	Greater than 7%
Coarse	2 pt/acre	2 pt/acre	2 - 3 pt/acre
Medium	2 - 2.5 pt/acre	2.5 pt/acre	2.5 - 3.75 pt/acre
Fine	3 pt/acre	3 pt/acre	3 - 3.75 pt/acre

¹ Rates are for single applications. Split applications of Volley may be used; apply at least 60% of the recommended rate up to 30 days before planting and the remaining balance, up to 40%, at planting.

20/27

Band Applications

This product may be applied as a band treatment. Use the following formulas below to determine the appropriate rate and volume per treated acre.

Band width in inches

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast rate per acre} = \text{Band rate per treated acre}$$

Band width in inches

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast volume per acre} = \text{Band volume per treated acre}$$

Weeds Controlled

Volley, applied as directed in this label, will provide control or partial control of the weeds listed in Table 4.

Partially controlled weeds will be severely stunted, or experience reduced height, vigor, or population compared to untreated areas. Depending on the infestation level or density, a follow-up treatment with another herbicide registered for use on corn may be needed to provide complete control.

Additional weeds may be controlled with tank mixes. See the Tank Mix Combinations section for recommended tank mix combinations and the additional weeds controlled.

Table 4: Weeds Controlled or Partially Controlled by Volley at Recommended Use Rates.

Grasses and Sedges	C = Control PC = Partial Control	Broadleaves	C = Control PC = Partial Control
barnyardgrass	C	beggarweed, Florida	C
crabgrass spp.	C	carpetweed	C
crowfootgrass	C	galinsoga	C
cupgrass, prairie	C	kochia	PC
cupgrass, southwestern	C	lambsquarters, common (2)	C
cupgrass, woolly	PC	nightshade, black	C
foxtail, bristly	C	nightshade, hairy	C
foxtail, giant	C	pigweed	C
foxtail, green	C	purslane, common	C
foxtail, robust (purple, white)	C	pusley, Florida	C
foxtail, yellow	C	ragweed, common	C
goosegrass	C	sida, prickly	C
johnsongrass, seedling	PC	smartweed spp.	C
millet, foxtail	C	waterhemp, tall	C
millet, wild proso	PC	waterhemp, common	C
nutsedge, yellow (1)	C		
panicum, browntop	C		
panicum, fall	C		
panicum, Texas (3)	C		
rice, red	C		
sandbur, field	PC		
shattercane	PC		
signalgrass, broadleaf (3)	C		
sprangletop, red	C		
witchgrass	C		

1. Yellow nutsedge requires a minimum of 2.5 pints. Incorporation will improve control.

2. Light to moderate infestations will be controlled. Heavy infestations may require a tank mixture or sequential herbicide.
3. Best control is achieved when Volley is applied within 5 days of planting and rainfall occurs shortly after application or mechanical incorporation is used to activate the herbicide. If it does not rain within 7 days, shallow cultivation will enhance activity. Excessive rainfall after application may reduce control. Under adverse weather conditions and/or heavy infestations, a cultivation or follow-up herbicide may be needed

Tank Mix Combinations

Additional weeds may be controlled with tank mixes of Volley and other products labeled for use on field corn, production seed corn, silage corn, sweet corn and popcorn. Tank mix combinations may be used in conventional, reduced, or no-till systems and may be applied by the same methods and at the same application timing as Volley unless otherwise specified in the tank mix product label.

Volley may be tank mixed with any other herbicide labeled for use on corn provided the compatibility of the tank mix is verified by a jar test and tank mixing with Volley is not prohibited by the label of the tank mix product. The compatibility of a tank mixture can be determined by mixing the ingredients of the herbicide mixture in their relative proportions in a glass jar as described for fluid fertilizer mixtures in Appendix I by substituting water for fluid fertilizer. Refer to the label of the tank mix product for applicable use directions, precautions and limitations, including additional weeds controlled. Do not exceed application rates on the respective product labels. Do not tank mix with another pesticide product that contains the same active ingredient as this product unless the label of either tank mix partner specifies the maximum dosages that may be used.

When tank mixing Volley with atrazine, do not exceed the maximum allowable rate of atrazine in your county or state. In some atrazine management areas, atrazine is more restricted. Consult your county extension office or state university for further information.

Use of Spray Adjuvants

Volley is a preemergence herbicide for which spray adjuvants have little or no influence on performance. However, several herbicides used in tank mixtures with Volley require use of adjuvants to aid in the burndown of emerged weeds. Use only those adjuvants recommended on the label of the tank mix product and approved for use in growing crops. Surfactants and/or low rate liquid fertilizers (28%, 30% or 32% UAN) or ammonium sulfate (AMS) adjuvants may be used with tank mixes applied preplant or preemergence to the crop.

Preemergence Tank Mix Combinations

Conventional Tillage (Volley Plus):

Tank Mix Herbicide †	Comments
Atrazine 4L	<ul style="list-style-type: none"> • May be applied preplant surface, preplant incorporated, preemergence. If emerged weeds are greater than 1.5 inches tall at the time of application, add an appropriate postemergence herbicide • Provides control or partial control of cocklebur, giant ragweed, ground cherry (spp), jimsonweed, kochia, morningglory (spp), mustards, sicklepod and velvetleaf • Use when there is heavy broadleaf weed pressure
Balance Pro	<ul style="list-style-type: none"> • Not labeled in all states; please refer to the label for Balance Pro for applicable directions for use, geographic and other restrictions • For use in field corn only • Refer to the use rates section for minimum use rates for Volley
Hornet WDG	<ul style="list-style-type: none"> • Tank mix with 4 – 5 oz/acre of Hornet® WDG herbicide to provide

	consistent control of velvetleaf, lambsquarters, pigweed species, waterhemp and triazine resistant varieties of these species. Also provides improved control of cocklebur, common ragweed, giant ragweed, common sunflower and jimsonweed.
Princep 4L	<ul style="list-style-type: none"> • Provides improve crabgrass or fall panicum control.
Python WDG	<ul style="list-style-type: none"> • Tank mix with 0.8 – 1.33 oz/acre of Python[®] WDG herbicide to provide consistent control of velvetleaf, lambsquarters, pigweed species, waterhemp and triazine resistant varieties of these species.

† Formulations that are not listed may be used. Perform a compatibility test and check the label of the tank mix product label for application rates, applicable use directions, precautions and limitations.

Reduced or No-Tillage Corn (Volley plus):

Tank Mix Herbicide †	Comments
Atrazine 4L	<ul style="list-style-type: none"> • Provides control or partial control of cocklebur, giant ragweed, ground cherry (spp), jimsonweed, kochia, morningglory (spp), mustards, sicklepod and velvetleaf • If emerged weeds are greater than 1.5 inches tall at the time of application, add an appropriate postemergence herbicide • Use in areas with heavy broadleaf weed pressure
Balance Pro	<ul style="list-style-type: none"> • Not labeled in all states; refer to the label for Balance Pro label for precautionary statements, directions for use, geographic and other use restrictions • For use in field corn only • Refer to use rate section for minimum use rates for Volley
Banvel/Clarity Marksman	<ul style="list-style-type: none"> • Apply preplant or preemergence in reduced/ no-till systems for burndown of existing weeds
Glyphomax Plus, Durango, Roundup WeatherMax, Touchdown	<ul style="list-style-type: none"> • Apply preplant for burndown of existing weeds • Weeds less than 6 inches tall are easiest to control with burndown herbicides applied in combination with Volley. • Always add ammonium sulphate (AMS) to tank mixes prior to addition of glyphosate (8.5 to 17 lb per 100 gal of spray).
Gramoxone Max	<ul style="list-style-type: none"> • Control annuals, suppress perennials
Pendimax [®] herbicide/ Prowl	<ul style="list-style-type: none"> • Preemergence to early postemergence to corn up to 3 inches tall, but before weeds are more than 1 inch tall.
Princep 4L	<ul style="list-style-type: none"> • Provides improved control of crabgrass and fall panicum
2,4-D	<ul style="list-style-type: none"> • Burndown existing weeds

† Formulations that are not listed may be used. Perform a compatibility test and check the product label for application rates, applicable use directions, precautions and limitations.

Postemergence Tank Mix Combinations

Volley may be applied before, with, or following the use of one or more of the following herbicides: Accent, Accent Gold, Aim, atrazine, Banvel, Basis, Basis Gold, Beacon, Bladex, Buctril, Buctril/atrazine, Clarity, Distinct, Exceed, Hornet WDG, Liberty, Lightning, Marksman, Peak, Permit, Princep, Prowl, Pendimax, Pursuit, Shotgun, Spirit and Steadfast. Refer to the label of the tank mix product for applicable directions for use, precautions and restrictions, and a list of weeds controlled. Volley may be tank mixed with any product approved for use on corn unless it is prohibited by the tank mix product label. **Note:** Do not use liquid fertilizer as the carrier when Volley is applied postemergence to corn as severe injury may result. The addition of liquid fertilizers used as adjuvants with Volley tank mixes applied postemergence to corn under environmental stress conditions may result in significant crop injury and should be avoided if the risk of crop injury is unacceptable.

23/27

When tank mixing, refer to the label of the tank mix product and follow additional use directions in the following table. **Volley can be applied to corn up to 11 inches tall.**

Postemergence Tank Mixes (Volley plus):

Tank Mix Herbicide	Rate	Comments
Hornet WDG	2 - 5 oz/acre	<ul style="list-style-type: none"> Always add NIS at 0.25% v/v or COC at 1% v/v.
Aim EW	0.5 oz/acre	<ul style="list-style-type: none"> Always add a NIS at 0.25% v/v.
Banvel Clarity Marksman	0.5 - 1 pt/acre 0.5 - 1 pt/acre 2 - 3.5 pt/acre	<ul style="list-style-type: none"> Early postemergence up to 8 inches tall corn on all soils. If grasses are more than 2-leaf stage, combine with another herbicide to control these weeds.
Buctril Buctril/atrazine Shotgun herbicide	1.5 pt/acre 2 pt/acre 2 - 3 pt/acre	<ul style="list-style-type: none"> Refer to tank mix product labels for applicable use directions, precautions and restrictions.
Atrazine	0.5 - 2 lb ai/acre	<ul style="list-style-type: none"> May be applied preplant surface, preplant incorporated, preemergence or early postemergence to corn up to 8 inches tall. If emerged weeds are greater than 1.5 inches tall at the time of application, add an appropriate postemergence herbicide. Note: The maximum atrazine application rate per year for corn is 2 lb active if applied only postemergence or 2.5 lb active if pre- and postemergence applications are made.
Distinct	4 - 6 oz/acre	<ul style="list-style-type: none"> Always add a NIS at 0.25% v/v and 1.25% UAN. Can be applied up to corn up to 10 inches tall.
Exceed	1 oz/acre	<ul style="list-style-type: none"> Always add crop oil concentrate at 1% v/v. See label for Exceed for geographic restrictions.
Liberty	16 - 28 oz/acre	<ul style="list-style-type: none"> For use on Liberty tolerant corn only. Apply to grass and broadleaf weeds up to 6 inches tall. Do not add additional surfactant.
Lightning	1.28 oz/acre	<ul style="list-style-type: none"> For use on Clearfield corn only. Use an NIS at 25%v/v and a liquid nitrogen fertilizer at 1 to 2 qt per acre or ammonium sulfate at 2.5 lb per acre.
Pendimax / Prowl	1.8 - 3.6 pt/acre	<ul style="list-style-type: none"> Apply preemergence or apply early postemergence to corn up to 3 inches tall, but before weeds are more than 1 inch tall.
Princep	1 - 3 lb ai/acre	<ul style="list-style-type: none"> May be applied preplant surface, preplant incorporated, preemergence to corn.
Pursuit 2.5L Pursuit 70DG	4 fl oz/acre 1.4 fl oz/acre	<ul style="list-style-type: none"> Use only on Clearfield varieties. Apply preplant incorporated, preplant surface, preemergence or early postemergence to weeds up to 3 inches tall.
Resource	4 - 6 oz/acre	<ul style="list-style-type: none"> Apply to weeds less than 5 inches tall. Add a crop oil concentrate at 1 to 2 pt/acre and either 28% nitrogen at 2% v/v or ammonium sulfate at 2.5 lb/acre. May cause some burn or spotting to corn leaves.
Spirit	1 oz/acre	<ul style="list-style-type: none"> Always add crop oil concentrate at 1% v/v. See label for Spirit for geographic restrictions.
2,4-D Ester	See label	<ul style="list-style-type: none"> Apply preplant surface or preemergence to

Accent 75WDG Beacon 75WDG Basis Steadfast	1/4 - 2/3 oz/acre 0.76 oz/acre 1/4 - 2/3 oz/acre 0.75 oz/acre	control emerged broadleaf weeds in corn. <ul style="list-style-type: none"> Minimum Volley use rates (pt/acre): <table border="1"> <tr> <td>Soil</td> <td><3%OM</td> <td>3-7%OM</td> <td>>7%OM</td> </tr> <tr> <td>Coarse</td> <td>1.5</td> <td>1.5</td> <td>2</td> </tr> <tr> <td>Medium</td> <td>1.5</td> <td>1.5 - 2</td> <td>2</td> </tr> <tr> <td>Fine</td> <td>1.5</td> <td>1.5 - 2</td> <td>2</td> </tr> </table> Always add NIS at 0.25% (v/v) and, in addition if applied under dry conditions, add 4% (v/v) clear liquid fertilizer. Banvel, Clarity, Marksman, Buctril, Buctril/atrazine may be added to this mixture to provide burndown and residual control of broadleaf weeds. 	Soil	<3%OM	3-7%OM	>7%OM	Coarse	1.5	1.5	2	Medium	1.5	1.5 - 2	2	Fine	1.5	1.5 - 2	2
Soil	<3%OM	3-7%OM	>7%OM															
Coarse	1.5	1.5	2															
Medium	1.5	1.5 - 2	2															
Fine	1.5	1.5 - 2	2															
Basis Gold	14 oz/acre	<ul style="list-style-type: none"> Minimum Volley use rates (pt/acre): <table border="1"> <tr> <td>Soil</td> <td><3%OM</td> <td>3-7%OM</td> <td>>7%OM</td> </tr> <tr> <td>Coarse</td> <td>1.5</td> <td>1.5</td> <td>2</td> </tr> <tr> <td>Medium</td> <td>1.5</td> <td>1.5 - 2</td> <td>2</td> </tr> <tr> <td>Fine</td> <td>1.5</td> <td>1.5 - 2</td> <td>2</td> </tr> </table> Always add COC at 1.0% v/v or, under dry arid conditions, add COC at 2.0% v/v plus 2 qt/acre of 28% liquid nitrogen or 2 lb/acre of ammonium sulfate. Banvel, Clarity, Marksman, Buctril, or Tough may be added to this mixture to provide burndown and residual control of broadleaf weeds. 	Soil	<3%OM	3-7%OM	>7%OM	Coarse	1.5	1.5	2	Medium	1.5	1.5 - 2	2	Fine	1.5	1.5 - 2	2
Soil	<3%OM	3-7%OM	>7%OM															
Coarse	1.5	1.5	2															
Medium	1.5	1.5 - 2	2															
Fine	1.5	1.5 - 2	2															

Appendix I

Procedure for Testing the Compatibility of Volley and Tank Mixes with Fluid Fertilizers

Since fluid fertilizers vary, the following procedure is suggested for determining whether Volley may be combined with a specific fluid fertilizer for spray tank application.

Materials Needed:

- Volley and any tank mix products
- Fluid fertilizer to be used
- Adjuvant for fertilizer tank mix: Use any adjuvant cleared for use on growing crops under 40 CFR 180.1001 to improve the compatibility of Volley with fluid fertilizers. The adjuvant that provides the best emulsification depends upon the specific fertilizer under consideration.
- Two 1 quart, wide mouth glass jars with lid or stopper
- Measuring spoons (a 25 ml pipette or graduated cylinder provides more accurate measurement)
- Measuring cup, 8 oz (257 ml)

Procedure:

1. Pour a pint (about 473 ml) of the fluid fertilizer into each of the quart jars.
2. Add Volley and any tank mix combination to the jars. The order of addition is wettable powders first with mixing, followed by flowables with mixing and the ECs last. The rate of wettable powders and dry flowables is 1 1/2 teaspoon per pound of product per acre to be applied. ECs should be added at the rate of 1/2 teaspoon for each pint per acre to be applied. Premixing the wettable powders in 1 oz of water before adding to the pint of fluid fertilizer will improve the compatibility of the final mixture.
3. Add 1/2 teaspoon (2 ml) adjuvant to one of the jars, label it as "with," and mix. The rate of 1/2 teaspoon per pint is equal to 3 pints of adjuvant per 100 gallons of fluid fertilizer.
4. Close both jars with lids or stoppers and mix the contents by turning the jars upside down 10 times.
5. Inspect the surface and body of the mixtures-

- (a) Immediately after completing the jar inversions
- (b) After allowing the jars to stand undisturbed for 30 minutes
- (c) And then again after turning the jars upside down 10 times after the 30 minute inspection

Evaluation:

If either mixture remains uniform for 30 minutes, the combination may be used. Should either mixture separate after 30 minutes, but readily remix uniformly with 10 jar inversions, the mixture can be used if adequate agitation is maintained in the tank. If the mixture with adjuvant is satisfactory but the one without adjuvant is not, be sure to use the adjuvant in the spray tank. Add the adjuvant first at a rate of 3 pints per 100 gallons of fluid fertilizer. Foaming may be minimized by using moderate agitation. **If non-dispersible oil, sludge, or clumps of solids form in the mixtures, the combination should not be used.**

Appendix II

Dry Bulk Fertilizer Impregnation

All individual state regulations relating to dry bulk fertilizer blending, registration, labeling, and application are the responsibility of the individual and/or company selling Volley, Volley fertilizer mixtures.

When applying Volley alone or in tank mixes with dry bulk fertilizers, follow all directions for use and precautions on the respective tank mix product labels regarding use rates, soil texture, application methods, and rotational restrictions. Use a minimum of 200 lb of dry bulk fertilizer per acre.

Approved Dry Fertilizer Ingredients for Use with Volley¹

Fertilizer	N	P	K
Ammonium Phosphate-Sulfate	16	20	0
Ammonium Sulfate	21	0	0
Diammonium Phosphate	18	46	0
Monoammonium Phosphate	11	56	0
Potassium Chloride	0	0	60
Potassium Sulfate	0	0	52
Urea ²	45	0	0

¹ Do not impregnate on fertilizers containing ammonium nitrate, potassium nitrate, or sodium nitrate.

² Some ureas may be phytotoxic when high rates are applied to corn. Use only urea rates known to be safe for corn application.

For impregnating pesticides on dry fertilizers, use suitable mixers equipped with suitable spraying equipment. The spray nozzles should be positioned inside the mixer to provide uniform spray coverage of the tumbling fertilizer. Volley should be sprayed uniformly onto the fertilizer using a fine spray pattern. Tank mix components may be applied as separate ingredients with powders and dry flowables added first or they may be mixed in a slurry in the proper ratio and added jointly. Volley may also be impregnated on dry bulk fertilizer in the field while the fertilizer is being spread using a pneumatic applicator equipped to impregnate herbicides.

The following table provides a reference to determine the amount of Volley to be mixed per ton of dry bulk fertilizer for a range of herbicide rates.

Fertilizer Rate (lb/acre)	Acres Covered (per ton)	Volley (pints/acre)		
		2 pt/acre	2.5 pt/acre	3 pt/acre
Pints Herbicide/Ton Fertilizer				
200	10	20	25	30
300	6.7	13.4	16.8	20.1

400	5	10	12.5	15
500	4	8	10	12
600	3.3	6.6	8.3	9.9
700	2.9	5.8	7.3	8.7

To determine the amount of Volley needed for other rates of fertilizer, use this formula:

$$\frac{\text{Volley (pints/acre)}}{\text{Pounds of fertilizer/acre}} \times 2000 = \text{Pints of Volley per ton of fertilizer}$$

If the herbicide/fertilizer mixture is too wet, use of a drying agent is required to provide a dry, free-flowing mixture. For mixtures to be used in spinning-disc applicators, Micro-Cel E calcium silicate powder (Manville, Filtration & Minerals) is recommended for use as a drying agent. Mixtures to be used in pneumatic applicators should use Micro-Cel E or Agsorb 16/30 RVM-MS granular clay (Oil-Dri Corporation). The drying agents should be added separately and uniformly to the prepared pesticide/fertilizer mixture, in a quantity that is sufficient to provide a suitable free-flowing mixture. Generally, less than 2% Micro-Cel E or 5% Agsorb 16/30 RVM-MS by weight is required.

Precaution: To avoid potential for explosion, do not impregnate Volley on ammonium sorbate nitrate, potassium nitrate, or sodium nitrate fertilizer or fertilizer blends. Do not impregnate on a single (0-20-0) or triple (0-46-0) super phosphate. Do not attempt to impregnate Volley on agricultural limestone as the herbicide will not be adequately absorbed.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Tenkoz warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, Tenkoz MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Tenkoz or the seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Tenkoz' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, Tenkoz shall not be liable for losses or damages resulting from handling or use of this product unless Tenkoz is promptly notified of such loss or damage in writing. To the extent

permitted by law, in no case shall Tenkoz be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Tenkoz or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of Dow AgroSciences LLC
EPA accepted

NEXT

LABEL

55467-8

01/07/2008

1/3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

JAN -7 2008

Mr. Robert E. Ridsdale
Tenkoz, Inc
100 North Point Center East, Suite 330
Alpharetta, GA 30022

Dear Mr. Ridsdale:

Subject: Volley Corn Herbicide (Supplemental for Use on Sweet Corn)
EPA Registration No. 55467-8
Application Dated October 10, 2007

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended is acceptable, provided you make the following change before you release the product for shipment.

--In the first paragraph under "Volley Corn Herbicide Applied Alone" revise "...to sweet corn using rates recommended in the product label..." to read "...to sweet corn using rates **stated** in the product label." The Agency no longer allows the word "recommended" on product labels when pertaining to application rates.

Submit one copy of your final printed labeling incorporating the above changes before you release the product for shipment. A stamped copy of labeling is enclosed for your records.

Sincerely,

Victoria K. Walters for
James A. Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505P)

Supplemental Labeling

2/3
Tenkoz Inc.

Tenkoz Inc. 100 North Point Center East Alpharetta, GA 30022 USA

Volley[®] Corn Herbicide

EPA Reg. No. 55467-8

Weed Control in Sweet Corn

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for **Volley[®] Corn Herbicide** before applying. Follow all applicable use directions and precautions on the EPA-registered label.
- Use of **Volley Corn Herbicide** according to this supplemental labeling is subject to all restrictions and limitations imposed by the label affixed to the product container.

Directions for Use

Volley Corn Herbicide Applied Alone

Used alone, **Volley[®] Corn Herbicide** may be applied to sweet corn using rates recommended in the product label for **Volley** and application methods and timing listed below.

Carriers and Spray Volume: Either water or liquid fertilizers such as solutions, slurries or suspensions may be used as liquid carriers. Refer to product label for **Volley** for details.

Application Timing and Methods: For the optimum period of effective weed control during the time most critical to corn production, preplant applications of **Volley** should occur as close as possible to planting. Preemergence applications should occur as close as possible to planting, but prior to weed emergence. Refer to product label for **Volley** for details.

Note: Do not apply **Volley** to sweet corn as an early postemergence application.

- **Early Preplant:** On medium and fine textured soils (see Table 1 in main label), **Volley** may be applied up to 30 days prior to planting.
- **Preplant Incorporation:** **Volley** and certain tank mixes may be mechanically incorporated in the top 2 inches of the soil with field cultivators, discs, or spring tooth harrows at any time within 14 days prior to planting.
- **Preemergence Surface:** **Volley** and certain tank mixes may be applied to the soil surface as a broadcast or banded application.
- **Postplant-Preemergence:** **Volley** may be applied immediately after planting but prior to corn emergence.
- **Banding Preemergence:** **Volley** may be applied in a 10 to 14 inch band after corn planting but prior to corn emergence.

Sprinkler Irrigation: Do not apply **Volley** by sprinkler irrigation. Use a sprinkler system only to incorporate **Volley** after application. Refer to product label for **Volley** for details.

Cultivation: Cultivation should be delayed as long as possible. If weeds develop, a shallow cultivation or rotary hoeing will generally result in improved weed control. Refer to product label for **Volley** for details.

Soil Texture and Organic Matter: The use rate of **Volley** is determined by a combination of two factors, soil texture and organic matter, which must be determined prior to application. Refer to Table 1 in main product label for soil texture classifications.

Use Rates for Conventional Tillage Systems: Refer to table 2 in main label for application rates and detailed information for conventional systems.

Use Rates for Reduced Tillage Systems: Refer to table 3 in main label for application rates and detailed information for reduced tillage systems.

Weeds Controlled: Refer to table 4 in main label for details.

Volley Applied in Tank Mix Combination

Additional weeds may be controlled with tank mixes of **Volley** and other products labeled for use on sweet corn. Tank mix combinations may be used in conventional, reduced, or no-till systems and may be applied using application methods and timing listed above unless otherwise specified in the tank mix product label. Refer to the product label for **Volley** for details on carriers, mixing instructions and testing for compatibility of tank mixtures.

ACCEPTED
with COMMENTS
In EPA Letter Dated:
JAN -7 2008

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
55467-8

®Trademark of Dow AgroSciences LLC

EPA Accepted:
Initial printing.

NEXT

LABEL

55467-8

01/07/2008

1/2



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

JAN -7 2008

Mr. Robert E. Ridsdale
Tenkoz, Inc
100 North Point Center East, Suite 330
Alpharetta, GA 30022

Dear Mr. Ridsdale:

Subject: Volley Corn Herbicide (Supplemental for Rotational Crops after Final Application)
EPA Registration No. 55467-8
Application Dated October 10, 2007

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended is acceptable. A stamped copy of labeling is enclosed for your records.

Sincerely,

James A. Tompkins for
James A. Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505P)

Supplemental Labeling

Tenkoz Inc.

Tenkoz Inc.

100 North Point Center East

Alpharetta, GA 30022 USA

Volley® Corn Herbicide

EPA Reg. No. 55467-8

Rotational Crop Restrictions After Final Application

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for **Volley® Corn Herbicide** before applying. Follow all applicable use directions and precautions on the EPA-registered label.
- Use of **Volley Corn Herbicide** according to this supplemental labeling is subject to all restrictions and limitations imposed by the label affixed to the product container.

Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

Rotational Crop Restrictions

When tank mixing with other herbicides, follow the most restrictive crop rotation guidelines on the label of each product used. The following rotational crops may be planted as indicated:

Crop	Interval
Corn (1)	Anytime – 0 Months After Application
Wheat	4 Months After Application
Alfalfa, Barley, Buckwheat, Clover, Dry beans (2), Guar, Kudzu, Lentil, Lespedeza, Lupin (4), Millet, pearl or proso, Oats, Pea (5), Potatoes, Rye, Sorghum, Soybeans, Sugar beets, Sunflower, Trefoil, Tobacco, Triticale, Vetch, Wild rice	Spring Following Application (3)

Numbers within parentheses (-) in the table refer to Specific Rotational Crop Requirements below.

- (1) If crop treated with **Volley** is lost, corn may be replanted immediately. Do not make a second application of **Volley**.
- (2) Dry beans includes: adzuki, kidney, lima, navy, pinto
- (3) Approved rotation crops list does not include any species of succulent beans and peas
- (4) Lupin includes: grain, white, white sweet
- (5) Pea includes: blackeyed, chick, cow, Crowder, field, pigeon, Southern

Note: These rotational crop restrictions supercede prior restrictions of 4 months for wheat and the spring following application for sorghum, soybeans, and tobacco.

®Trademark of Dow AgroSciences LLC

EPA Accepted:
Initial printing.

