

55467-8

2-14-2008

1/4



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

FEB 14 2008

Mr. Robert E. Ridsdale  
Tenkoz, Inc  
100 North Point Center East, Suite 330  
Alpharetta, GA 30022-8242

Dear Mr. Ridsdale:

Subject: Volley Corn Herbicide (Revise Supplementals for Use on Sweet Corn and Revision of  
Rotational Crop Intervals)  
EPA Registration No. 55467-8  
Application Dated January 16, 2008

The labeling referred to above, submitted in connection with registration under the Federal  
Insecticide, Fungicide, and Rodenticide Act as amended is acceptable. Amended labeling  
supersedes supplemental labeling approved January 7, 2008. A stamped copy of labeling is  
enclosed for your records.

Sincerely,

*James A. Tompkins*  
James A. Tompkins  
Product Manager 25  
Herbicide Branch  
Registration Division (7505P)

# Supplemental Labeling

ACCEPTED

FEB 14 2008

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended for the pesticide  
registered under  
EPA Reg. No. 55467-8

Tenkoz Inc.

Tenkoz Inc.

100 North Point Center East

Alpharetta, GA 30022 USA

## Volley® Corn Herbicide

EPA Reg. No. 55467-8

### Weed Control in Sweet Corn

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for **Volley® Corn Herbicide** before applying. Follow all applicable use directions and precautions on the EPA-registered label.
- Use of **Volley Corn Herbicide** according to this supplemental labeling is subject to all restrictions and limitations imposed by the label affixed to the product container.

#### Directions for Use

##### Volley Corn Herbicide Applied Alone

Used alone, **Volley® Corn Herbicide** may be applied to sweet corn using rates stated in the product label for **Volley** and application methods and timing listed below.

**Carriers and Spray Volume:** Either water or liquid fertilizers such as solutions, slurries or suspensions may be used as liquid carriers. Refer to product label for **Volley** for details.

**Application Timing and Methods:** For the optimum period of effective weed control during the time most critical to corn production, preplant applications of Volley should occur as close as possible to planting. Preemergence applications should occur as close as possible to planting, but prior to weed emergence. Refer to product label for **Volley** for details.

**Note:** Do not apply **Volley** to sweet corn as an early postemergence application.

- **Early Preplant:** On medium and fine textured soils (see Table 1 in main label), **Volley** may be applied up to 30 days prior to planting.
- **Preplant Incorporation:** **Volley** and certain tank mixes may be mechanically incorporated in the top 2 inches of the soil with field cultivators, discs, or spring tooth harrows at any time within 14 days prior to planting.
- **Preemergence Surface:** **Volley** and certain tank mixes may be applied to the soil surface as a broadcast or banded application.
- **Postplant-Preemergence:** **Volley** may be applied immediately after planting but prior to corn emergence.
- **Banding Preemergence:** **Volley** may be applied in a 10 to 14 inch band after corn planting but prior to corn emergence.

**Sprinkler Irrigation:** Do not apply **Volley** by sprinkler irrigation. Use a sprinkler system only to incorporate Volley after application. Refer to product label for **Volley** for details.

**Cultivation:** Cultivation should be delayed as long as possible. If weeds develop, a shallow cultivation or rotary hoeing will generally result in improved weed control. Refer to product label for **Volley** for details.

2/4

**Soil Texture and Organic Matter:** The use rate of **Volley** is determined by a combination of two factors, soil texture and organic matter, which must be determined prior to application. Refer to Table 1 in main product label for soil texture classifications.

3/4

**Use Rates for Conventional Tillage Systems:** Refer to table 2 in main label for application rates and detailed information for conventional systems.

**Use Rates for Reduced Tillage Systems:** Refer to table 3 in main label for application rates and detailed information for reduced tillage systems.

**Weeds Controlled:** Refer to table 4 in main label for details.

### **Volley Applied in Tank Mix Combination**

Additional weeds may be controlled with tank mixes of **Volley** and other products labeled for use on sweet corn. Tank mix combinations may be used in conventional, reduced, or no-till systems and may be applied using application methods and timing listed above unless otherwise specified in the tank mix product label. Refer to the product label for Volley for details on carriers, mixing instructions and testing for compatibility of tank mixtures.

®Trademark of Dow AgroSciences LLC

EPA Accepted:  
Initial printing.

# Supplemental Labeling

4/4  
Tenkoz Inc.

Tenkoz Inc.

100 North Point Center East

Alpharetta, GA 30022 USA

## Volley® Corn Herbicide

EPA Reg. No. 55467-8

### Rotational Crop Restrictions After Final Application

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for **Volley® Corn Herbicide** before applying. Follow all applicable use directions and precautions on the EPA-registered label.
- Use of **Volley Corn Herbicide** according to this supplemental labeling is subject to all restrictions and limitations imposed by the label affixed to the product container.

#### Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

#### Rotational Crop Restrictions

When tank mixing with other herbicides, follow the most restrictive crop rotation guidelines on the label of each product used. The following rotational crops may be planted as indicated:

Crop	Interval
Corn (1)	Anytime – 0 Months After Application
Wheat	4 Months After Application
Alfalfa, Barley, Buckwheat, Clover, Dry beans (2), Guar, Kudzu, Lentil, Lespedeza, Lupin (4), Millet, pearl or proso, Oats, Pea (5), Potatoes, Rye, Sorghum, Soybeans, Sugar beets, Sunflower, Trefoil, Tobacco, Triticale, Vetch, Wild rice	Spring Following Application (3)

Numbers within parentheses (-) in the table refer to Specific Rotational Crop Requirements below.

- (1) If crop treated with **Volley** is lost, corn may be replanted immediately. Do not make a second application of **Volley**.
- (2) Dry beans includes: adzuki, kidney, lima, navy, pinto
- (3) Approved rotation crops list does not include any species of succulent beans and peas
- (4) Lupin includes: grain, white, white sweet
- (5) Pea includes: blackeyed, chick, cow, Crowder, field, pigeon, Southern

**Note:** These rotational crop restrictions supercede prior restrictions of 4 months for wheat and the spring following application for sorghum, soybeans, and tobacco.

®Trademark of Dow AgroSciences LLC

EPA Accepted:  
Initial printing.

