

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

9-28-07

Robert Ridsdale 100 North Point Center E. Suite 330 Alpharetta, GA 30022-8242

Subject: EPA Reg. No. 55467-9 / Buccaneer Plus Glyphosate Herbicide Supplemental Labeling

Dear Mr. Ridsdale:

The labeling referred to above is acceptable provided you make the following changes:

1. Change "General Information" to "Product Information".

A stamped copy of the label is enclosed for your records. You must submit one copy of the final printed label before you release the product for shipment. This supplemental label must be incorporated within the master label within 18 months from the date of this letter. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA. Your release for shipment of the product constitutes acceptance of these conditions. If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov

Sincerely,

James A. Tompkins

Product Manager 25

Herbicide Branch

Registration Division (7505P)

Erik Kruft An

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR BUCCANEER PLUS GLYPHOSATE HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using BUCCANEER PLUS GLYPHOSATE HERBICIDE as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application. CEPTED with COMMENTS

BUCCANEER PLUS GLYPHOSATE HERBICIDE EPA Letter Dated:

EPA Reg. No. 55467-9

Buccaneer is a trademark used under license.

9-28-09 Under the Federal Insecticide, Fungicide, and Rodenticide Act

FOR APPLICATIONS TO Dry Beans, Dry Peas, Lentils, and amended, for the pesticide Chickpeas only

55467.9

Keep out of reach of children. WARNING! AVISO!

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.) In case of emergency involving this product, call day or night, 1-800-424-9300.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This label must be in the possession of the user at the time of application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY® CROPS) DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for BUCCANEER PLUS GLYPHOSATE HERBICIDE for essential product performance information.

See "Vegetable Crops" section of the label booklet for General Directions that apply to all vegetable crops

USE DIRECTIONS

Spot Treatment, or Preharvest (Dry Beans, Dry Peas, Lentils and Chickpeas only)

USE INSTRUCTIONS: This product may be applied as an over-the-top broadcast spray or as a spot treatment to control labeled weeds in dry beans, dry peas, lentils or chickpeas. For spot treatment, to control troublesome weeds such as Canada thistle, quackgrass, mayweed (dog fennel), and milkweed, apply in 10 to 20 gallons of water per acre through ground broadcast spray equipment or use a 2 percent solution in a hand-held sprayer. For preharvest treatments, apply in 3 to 20 gallons of water per acre at the hard dough stage of the legume seed (30 percent grain moisture or less).

PRECAUTIONS, RESTRICTIONS: Follow the limitations listed in the table below. Only one application per year may be made; do not combine a preharvest spray with a spot treatment on the same crop area. Employ at least a 30-day plant-back interval between treatment and replanting for any crop not specified for treatment in this label. Do not feed treated vines and hay from these crops to livestock. Do not treat cowpeas or field (feed) peas, since these crop are considered to be grown as livestock feed. Preharvest application is not recommended for dry legumes grown for seed, as a reduction in germination or vigor may occur.

Crop	Maximum Rate (fluid ounces per Acre)	Minimum Pre- Harvest Interval (days)	
Dry Beans	32	7	
Dry Peas, Lentils, and Chickpeas	26	14	

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for BUCCANEER PLUS GLYPHOSATE HERBICIDE before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Tenkoz Inc. 100 North Point Center East Alpharetta, GA 30022

08/09

