

# CATTLE MINERAL FOL

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See label for directions. Do not use if contents are past expiration date.

**Directions:** Feed 1/2 pound per 400 pounds of live weight per month. For animals weighing less than 400 pounds, feed 1/4 pound per 400 pounds of live weight per month. For animals weighing more than 400 pounds, feed 3/4 pound per 400 pounds of live weight per month. Feed continuously throughout the year.

**Warnings:** Do not use if contents are past expiration date. Do not use if contents are wet or clumpy. Do not use if contents are contaminated.

**Net Weight: 50 LBS.**

# 803

# CATTLE MINERAL FOL

**Directions:** Feed 1/2 pound per 400 pounds of live weight per month. For animals weighing less than 400 pounds, feed 1/4 pound per 400 pounds of live weight per month. For animals weighing more than 400 pounds, feed 3/4 pound per 400 pounds of live weight per month. Feed continuously throughout the year.

**Warnings:** Do not use if contents are past expiration date. Do not use if contents are wet or clumpy. Do not use if contents are contaminated.

**Net Weight: 50 LBS.**

**ENVIRONMENTAL PRECAUTIONS:** This product is toxic to fish. Do not use in areas where it may come in contact with surface water by cleaning of equipment.

**STORAGE AND DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Store in a dry place in original container. Contents should be kept away from fire, loading equipment. Then dispose of this product by burning, if allowed by State and local health officials, or by other means approved by local health officials.

**BEST AVAILABLE COPY**

12-22-86

01158

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460  <b>NOTICE OF PESTICIDE:</b> <input type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION <i>(Under the Federal Insecticide, Fungicide,          and Rodenticide Act, as amended)</i>	EPA REGISTRATION NO. 55591-1	DATE OF ISSUANCE DEC 22 1986
	TERM OF ISSUANCE Until Reregistration	
	NAME OF PESTICIDE PRODUCT Gallon of Shell	

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

[ ]

100 West 100th Street  
 Danville, NY 12832

100/101/562  
 11/1

**NOTE:** Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered under the Act, provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.

2. Make the labeling changes listed below before you release the product for shipment:

- Change the name "Gallon of Shell" to "55591-1."
- Change the name "Gallon of Shell" to "Gallon of Shell."
- Change the sentence "Do not apply directly to water." to its place and "Do not apply directly to water."
- Change the sentence "Do not apply directly to water." to its place and "Do not apply directly to water."

3. Submit five (5) copies of your final product labeling before you release the product for shipment.

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

*[Handwritten Signature]*

12/22/86

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

George T. LaRocca  
Product Manager (15)  
Insecticide-Rodenticide Branch  
Registration Division (25-767C)

Enclosure



# CATTLE MINERAL ROL

For Self-Feeding

(With RABON® Oral Larvicide)

To prevent the development of Horn Flies, Face Flies, House Flies and Stable Flies in the manure of treated cattle.

ACTIVE INGREDIENT	BY WEIGHT
2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate	0.95%*
INERT INGREDIENTS	99.05%**
TOTAL	100.00%

\*Each pound contains 4.3 grams RABON.  
\*\*Refers only to ingredients which are not larvicidal.

### GUARANTEED FEED ANALYSIS

Calcium	Minimum	13.00 %
Calcium	Maximum	15.00 %
Phosphorus	Minimum	8.00 %
Salt	Minimum	20.00 %
Salt	Maximum	24.00 %
Magnesium	Minimum	1.50 %
Zinc	Minimum	.15 %
Iron	Minimum	.10 %
Manganese	Minimum	.05 %
Copper	Minimum	.02 %
Iodine	Minimum	.005%
Cobalt	Minimum	.001%
Vitamin A	Minimum USP Units/lb.	400,000
Vitamin D <sub>3</sub>	Minimum USP Units/lb.	75,000
Vitamin E	Minimum INT Units/lb.	100

### FEED INGREDIENTS

Dicalcium phosphate, monocalcium phosphate, calcium carbonate, salt, potassium sulfate, magnesium sulfate, magnesium oxide, dried cane molasses, bentonite, zinc oxide, copper oxide, iron sulfate, manganous oxide, potassium iodide, cobalt carbonate, cane molasses, mineral oil, vitamin A supplement, vitamin D<sub>3</sub> supplement, vitamin E supplement, sodium selenite, natural and artificial feed flavors.

IMPORTANT: See reverse side of tag for feeding directions.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See reverse of tag for additional precautionary statements.

EPA Reg. No. \_\_\_\_\_

EPA Est. No. \_\_\_\_\_

RABON is the registered trademark of Shell Oil Company.

Manufactured By **BURKMANN MILLS**  
Danville, Kentucky 40422  
NET WT. 50 LBS.

DEC 2 2 1986

803

Under the Federal Insecticide, Fungicide, and Rodenticide Act  
EPA Reg. No. \_\_\_\_\_

55591-1

Product Number 803

### FEEDING DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Allow free choice to cattle. Cattle should consume an average of 1/2 pound of the mineral per 100 pounds of body weight per month. This will supply the recommended average daily intake of 70 mg. RABON Oral Larvicide per hundred pounds body weight. Animals should not be starved for minerals before initiation of the RABON Cattle Mineral Mix program but OMIT all salt from other feed after the program is started. Do not feed other free choice sources of salt or mineral containing salt in any form. Allow one mineral feeder for each 15-20 head, and place near watering and loafing areas. Put out no more than one month's supply at a time, and protect from rain.

If consumption of Cattle Mineral ROL is above 1/2 pound per 100 lbs. of body weight per month, reduce the number of feeding locations or relocate. If consumption is below 1/2 pound per 100 lbs. of body weight per month, increase the number of feeding locations. Start feeding Cattle Mineral ROL early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall.

Cattle Mineral ROL may be fed up to slaughter and to lactating dairy cows.

Cattle Mineral ROL prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, Cattle Mineral ROL should be used in conjunction with other good management and sanitation practices.

### CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Storage: Store in a dry place in original container. Container Disposal: Completely empty bag into feeding equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke. Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.