EPA Form 8570-6 (Rev. 5-76) PREVIOUS EDITION MAY BE USED UNTIL SUPPLY IS EXHAUS

WILBRO FERTILIZER with .75% RONSTAR® FOR ORNAMENTAL and RECREATIONAL TURF

50 lb. Bag Treats up to 8,150 Sq.Ft. FOR USE ON PARKS, GOLF COURSES, ATHLETIC FIELDS, AND OTHER RECREATIONAL / ORNAMENTAL TURFGRASS

FOR SALE TO, USE AND STORAGE BY COMMERCIAL TURF AND LANDSCAPE PERSONNEL ONLY Not for use on turf being grown for sale or other commerical use as sod, or for commercial seed production, or for research purposes. Not for use on Home Lawns.

Controls: Crabgrass, Annual Bluegrass, Goosegrass, Florida Pusley, Oxalis, Stinging Nettle, Carpetgrass, Pigweed & Other Weeds Listed

ACTIVE INGREDIENT: Oxadiazon; [2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl) /\(\lambda^2\)-1,3,4-oxadiazolin-5-one] -INERT INGREDIENTS:-0.75% Total

Contains RONSTAR® Herbicide a Reg.Trade Mark of Rhone Poulenc Inc.

KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (if you do not understand the label, find someone to explain it to you in detail.) STATEMENTS OF PRACTICAL TREATMENT

in sWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists. IF ON SKIN: Wash

with plenty of soap and water. Set medical attention if irritation persists. If ON SKIN! Wash with plenty of soap and water. Set medical attention if irritation persists.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes eye and skin irritation. Avoid contact with skin, eyes or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Do not allow domestic animals to graze treated areas. Avoid contamination of feed and foodstuffs

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters. Apply only as specified on this

STORAGE AND DISPOSAL

not contaminate water, food, or feed by storage or disposal

AGE: Store product in its original bag, in a cool, dry, locked place out of reach of ren. in case of spill, sweep up, and use according to directions or dispose of as follows.

PESTICIDE DISPOSAL: Pesticide wastes are tooc. Improper disposal of excess pesticide, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance. CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary tandfill or by incineration, or, if allowed by State and local

authorities, by burning, If burned stay out of smoke.

GUARANTEED FERTILIZER ANALYSIS Total Nitrogen (N).
Nitrate Nitrogen
Ammoniacal Nitrogen
Water Soluble Nitrogen
and/or Urea Nitrogen 15% Water insoluble Nitrogen Water insoluble Nitrogen
Available Prosphono Acid (P_iO₂). 0%
Soluble Potash (K_iO). 15%
Derived from Ammonium Sulfate, Ureatorm, Muriate of
Potash; Manganese as Mn.5%; Iron as Fe 1.0%; Chlonne <15%

FOR LAWNS AND ORNAMENTAL TURF

When applied according to directions this product can be used safely on the following established turfgrass varieties: Bermudagrass, Bluegrass, Buffalograss, Perennial Ryegrass, St. Augustinegrass, Tall Fescue, and Zoysiagrass, it may be used in the control of weeds in lawns and turfgrass areas associated with apartment and condominium complexes, parks, golf courses, athletic fields, and cemeteries.

EPA Reg. No. 55615-U

EPA EST. NO. 55615-SC-1

NET WEIGHT 50 POUNDS Notice 1996.

CONTROLS WEED GRASSES & BROADLEAF WEEDS

Weed Grasses

**Annual Bluegrass *Barnyardgrass Bristly Foxtail Carpetgrass

Crabgrass Fall Panicum Field Sandbur "Goosegrass Green Foxtail "Jungle Ricegrass
"Panama Paspalum

Groundsel, common Common Purslane *Evening Primrose Florida Pusiey *Galinsoga Golden Ragwort Lambsquarter

Bittercress

Carpetweed

"Livewort

Broadleaf Weeds Pennsylvania Smartweed Petty Spurge
Prostrate Spurge Redroot Pigweed **Spotted Catsear Stinging Nettle
*Swinecress Yellow Woodsorrel

and the second distribution of the second distri

For Weeds without "Asterisks, Use 50 lbs. to treat 8,150 sq.ft.
For weeds with 1" or 2" asterisks see rates in "Tolerant Weeds" paragraphs below.

HOW TO APPLY

For best weed control rake the area free of leaves and thatch and mow if necessary so that granules will reach the soil surface. Measure the area to be treated and uniformly apply the required amount using a fertilizer spreader (see Spreader Settings chart). Do not apply to well turf. Do not over-lap spreader applications, on St. Augustinegrass since discoloration "streaking" may temporarily occur from "double rate". If rain is not expected within 24 hours sprinkle treated turf with 1/4 to 1/2 inch to increase effectiveness.

HOW MUCH TO USE

FOR CRABGRASS and OTHER WEEDS LISTED ABOVE

FOR CRABGRASS and OTHER WEEDS LISTED ABOVE
For susceptible weeds listed above (without "asterisk") apply 50 lbs. per 8,150 sq.Ft. when
turn has been in a wearly maintenance program (equal to 2 lbs. A.I. herbicide per acre.) Apply
50 lbs. to treat 5,500 sq.Ft. of lawn if it has been heavily infested with Crabgrass and other
susceptible weeds (equal to 3 lb. A.I. herbicide /acre). Apply in early spring 1 to 2 weeks prior
to weed seed germination. This product is a pre-emergence herbicide. It will not control
growing weeds; only germinating weed seeds. Late applications will control only those weed
seeds that germinate after the application. See "When To Apply" Chart.

* MORE TOLERANT WEEDS
BARNYARDGRASS, EVENING PRIMROSE, GALINSOGA,
GROUNDSEL and SWINECRESS

GROUNDSEL, and SWINECRESS
Where these weeds are a problem apply 50 lbs. to treat 5,500 Sq.Ft. of Lawn; (equal to 3 lbs. active herbicide per acre), in early spring or 1 to 2 weeks before expected weed seed germination. Will not control emerged and growing weeds; only effective against germinating weed seeds. Observe 50#/5,500 sq.ft. maximum rate for St. Augustine grass.

** MOST TOLERANT WEEDS

GOOSEGRASS, JUNGLERICE, LIVERWORT, PANAMA PASPALUM, PROSTRATE SPURGE

For these most tolerant weeds apply 50 lbs. to treat 4,100 Sq.Ft. (4 lbs. A.I./acre) and apply 1 to 2 weeks prior to seed germination. See Chart for When To Apply. Observe 50#/5,500 sq.ft. maximum rate for St. Augustine grass.

FOR ANNUAL BLUEGRASS (Poa Annua)

Apply 50 lbs. per 4,100 to 5,500 Sq.Ft. of Lawn; equal to 3 to 4 lbs. active herbicide per acre. Apply in late summer to fall or in early spring 1 to 2 weeks before expected weed seed germination. Use 4 lb. rate for expected heavy seed germination. Will not control annual bluegrass present at time of application; therefore only germinating seeds. Refer to chart for proper timing in your locale. Observe 50#/5,500 sq.ft. maximum rate for St. Augustine grass.

SPREADER SETTINGS FOR SQUARE FOOT COVERAGE

AMT. OF for 8,1500 Sq.Ft. 5,500 Sq.Ft. 4,100 Sq.Ft.

EADER PRODUCT 2#A,L/Acre 3#A,L/Acre 4#A,L/Acre SPREADER 50 LBS. CYCLONE B-1 6 1/2 4 1/2 SCOTTS #35

NOTE: Spreader rate varies with walking speed

WHEN TO APPLY

APPROPRIATE TIMES TO APPLY THIS PRE-EMERGENT HERBICIDE JAN, FEB. MAR. APR. MAY JUNE JULY AUG. SEPT. OCT. NOV. DEC 1. 3.

3a •

For CRABGRASS and other weeds germinating in the Spring in Northern turf.
For GOOSEGRASS and other Late Germinating weeds such as PROSTRATE SPURGE.
For GOOSEGRASS in Southern Turf.
For POA ANNUA in Northern Turf.
For POA ANNUA in Southern Turf.
For Fall-Winter-Spring control of ANNUAL WEEDS in Southern Turf.

PRECAUTIONS IN USING

Do not use on desirable Red Fescue, Benigrass varieties, Dichongra, or Centipede grass since damage will occur. Do not use on Putting greens or 7 (ext. due to varying cultural practices used. Use only the rate of 50 lbs. (o tract 5,500 Sq.Ft. on St. Augustinegrass since temporary discoloration will occur from heavier rates. Avoid contact with flowers and shrubs except as recommended on the Ornamental Use Supplemental Labeling available for this product. For newly seeded lawns do not spx/rtkis product until after the second or third cutting. For best lawn care, place new grass speed in the fall of the year. Spring pre-emergence weed control can then be applied. weed control can then be applied.

RESEEDING TREATED AREAS: Delay new seeding upitil 4 triopths after treatment.

NOTICE: Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label. (55-RSTT5.PM5/ EPA revi. of 02-28-95 versions this dated 04-04-95)

P.O. BOX 318 - NORWAY, S.C. 29113 U.S.A.

ACCEPTED
with COMMENTS
in EPA Letter Dated

APR 17 1995

Under the Federal Insecticide, Fundamental and Redenticide Act as amended for the pesticide registered under EPA Reg. No. 55615-4