

pm 18

55638-12

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460	EPA REGISTRATION NO. 55638-12	DATE OF ISSUANCE MAY 08 1992
	TERM OF ISSUANCE	
NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION <i>(Under the Federal Insecticide, Fungicide,          and Rodenticide Act, as amended)</i>	NAME OF PESTICIDE PRODUCT Condor WP	

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Ecogen, Inc.  
 2005 Cabot Boulevard West  
 Langhorne, PA 19047-1810

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5). Once a pesticide is unconditionally registered, however, it is not regarded as permanently acceptable. Unconditional registration does not eliminate the need for continual reassessment of a pesticide. If EPA determines, at any time, that additional data are required to maintain in effect an existing registration, the Agency will require submission of such data under Section 3(c)(2)(B) of the Federal Insecticide Fungicide and Rodenticide Act (FIFRA).

A stamped copy is enclosed for your records.

CC: Owen Beeder, RSB/RD

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

*[Handwritten Signature]*

DATE

5/7/92

**CONDOR<sup>R</sup>**  
**Wettable Powder Bioinsecticide**

**Active Ingredient:**

Bacillus thuringiensis subspecies  
kurstaki strain EG2348  
Lepidopteran active toxin. . . . . 10.0%  
Inert Ingredients. . . . . 90.0%  
Total. . . . . 100.0%  
1.6 oz active ingredient per pound.

CONDOR<sup>R</sup> Bioinsecticide is a biological insecticide for the control of lepidopteran pests.

**KEEP OUT OF REACH OF CHILDREN**  
**DANGER**

**STATEMENT OF PRACTICAL TREATMENT**

- If in Eyes: Flush with plenty of water. Call a physician.
- If Swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
- If Inhaled: Remove to fresh air. Get medical attention.
- If on Skin: Wash with soap and water.

**PELIGRO**

Precaution Al Usuario:

Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

EPA Reg. No. 55638-12  
EPA Est. No. 769-GA-1

Ecogen Inc.  
2005 Cabot Blvd. West  
Langhorne, PA 19047-1810

Net Contents 4 U.S. Pounds

55638-12

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND Domestic Animals

Causes irreversible eye damage. Harmful if inhaled. Avoid breathing dust or spray mist. Harmful if swallowed. Do not get in eyes, on skin or on clothing. Applicators, mixers and loaders must wear goggles. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

### ENVIRONMENTAL HAZARDS

Do not contaminate water when disposing of equipment washwaters.

Not for Use or Storage In or Around the Home

### RE-ENTRY STATEMENT

Do not apply this product in such a manner as to directly, or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in areas about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is a reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: DANGER: Area treated with CONDOR<sup>R</sup> on (date of application). Do not enter without protective clothing.

### Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

**Storage:** Store in a cool, dry place inaccessible to children.

#### Pesticide

**Disposal:** Do not contaminate water when disposing of equipment washwaters. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

#### Container

##### Disposal:

**Bag:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke!

### Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Preharvest Interval:** There are no restrictions for applying CONDOR<sup>R</sup> bioinsecticide up to the time of harvest.

CONDOR<sup>R</sup> is a selective insecticide for use against the lepidopteran larvae listed in the Application Rate table. Larvae must consume deposits of CONDOR<sup>R</sup> to be affected. Always follow these directions:

- o Treat when larvae are young (early instars) and are actively feeding on exposed plant parts.
- o Apply before foliar damage has occurred.
- o Thorough spray coverage is essential for good insect control. Directed drop nozzles may be required for some crops.

- o When insect infestations are heavy, use the higher label rates, shorten the spray interval, and/or use larger total spray volume to improve spray coverage.
- o Repeat applications at an interval sufficient to maintain control, usually 3-14 days depending on plant growth, insect activity, and weather conditions after spraying.
- o For improved durability of spray deposits, an approved spreader/sticker is strongly recommended for hard-to-wet crops, such as cole crops.
- o Spray water/spray tank solution should not exceed pH 8.0. If necessary, buffer water to near neutral pH (7.0).
- o After consuming a lethal dose of CONDOR<sup>R</sup>, larvae will cease feeding, but may remain on the foliage until they die within several days. Dying larvae move slowly, discolor, shrivel and blacken prior to death.
- o Local conditions may affect the use of CONDOR<sup>R</sup>. Consult State Agricultural Extension Specialists for specific recommendations related to local crop protection problems.

#### Application Instructions

CONDOR<sup>R</sup> bioinsecticide may be applied with conventional ground or aerial equipment with quantities of water sufficient to provide thorough coverage of infested plants. Do not apply this product through any type of irrigation system. To obtain a suitable mixture with water, add enough water to allow maximum agitation. With agitator running, slowly add in the CONDOR<sup>R</sup>. Continue agitation. Then add remainder of water and other spray materials and agitate until mixed. Maintain suspension while loading and spraying. Do not mix more CONDOR<sup>R</sup> than can be used in a 12-hour period. Rinse and flush spray equipment thoroughly following each use. Do not contaminate water when disposing of equipment washwaters.

## Warranty and Conditions of Sale

Ecogen warrants that this product conforms to the description on this label and is reasonably fit for the purposes stated on this label when used in accordance with the directions on this label under normal conditions of use.

ECOGEN MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

If this product is defective, Buyer's exclusive remedy shall be the replacement of the product, or if replacement is impracticable, refund of the purchase price. In no case will Ecogen be liable for incidental, consequential or special damages resulting from the handling, storage or use of this product.

CONDOR<sup>R</sup> is a registered trademark of Ecogen, Inc.  
Patent Pending  
Vers. 3/91 c 1991 Ecogen, Inc.

CONDOR<sup>R</sup> for Vegetable and Field Crops

Crop	Pest
Arugala Asparagus Beans Beets Bok Choy Broccoli Brussel sprouts Cabbage Canola Cauliflower Celery Citrus Chicory Collards Corn, field & sweet Cucumber Cucurbits Eggplant Escarole Endive Grapes Horseradish Kale Lettuce (Head, Leaf, Romaine) Melons Mustard Greens Parsnips Peanuts Peas Peppers Potatoes Radishes Soybeans Spinach Squash Strawberries Sugar Beets Sweet Potatoes Swiss chard Tomatoes Turnips Watercress	Armyworms Beet armyworm Cabbage looper Cabbage webworm Celery leaf-tier Corn earworm Cross striped cabbageworm Diamondback moth European corn borer Fall armyworm Grape Leaf folder Grapeleaf Skeletonizer Green cloverworm Imported cabbageworm Melonworm Omniverous leafroller Orange dog Pickleworm Rind worm complex Saltmarsh caterpillar Soybean looper Tobacco budworm Tomato fruitworm Tomato hornworm Tomato pinworm Velvetbean caterpillar

CONDOR<sup>R</sup> for Vegetable and Field Crops (Continued)

Crop	pest
Cotton <sup>1</sup>	Beet armyworm, Cotton bollworm, Tobacco budworm, Cabbage looper, Soybean looper
Pasture	ARMYWORM
Alfalfa, Hay & Seed	Alfalfa Caterpillar, Armyworms, Loopers
Tobacco	Tobacco budworm, Tobacco hornworm
Rate to Apply <sup>2</sup>	1.0 to 3.0 lbs/acre

1 Use CONDOR<sup>R</sup> to control light to moderate populations of newly hatched worms in integrated pest management programs. Repeat treatments at 4 to 5 day intervals or as long as necessary and results are acceptable. For control of cotton bollworm and tobacco budworm ovicides such as LARVIN<sup>R</sup> should be combined with CONDOR<sup>R</sup> at each application. LARVIN<sup>R</sup> is a trademark of Rhone Poulenc.

2 For ground applications, use a minimum of 5 gallons of water per acre for field crops and 20 gallons per acre for vegetables. For aerial applications, use a minimum of 2 gallons of water per acre.