

055638-00037-073197

Printed: 12:44:27 Friday, 21 May, 1999 # 4 / 2870

Systems Integration Group, Inc.

Lepinox[®] wdg bioinsecticide

LEPINOX[®] WDG water dispersible granule bioinsecticide is a biological insecticide for the control of lepidopterous pests.

Active Ingredient:

<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i> EG7826	
Lepidopteran active toxin	15.00%
Inert Ingredients	85.00%
TOTAL	100.00%

2.4 oz. active ingredient per pound
Potency: 52,500 Fall Armyworm Units per milligram of product. Potency units should not be used to adjust use rates beyond those specified in the Directions for Use section.

KEEP OUT OF REACH OF CHILDREN WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

STATEMENT OF PRACTICAL TREATMENT

If In Eyes: Hold eyelid open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS


Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear goggles or faceshield. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

Personal Protective Equipment (PPE):

Applicators and other handlers must wear:

- Long sleeved shirt and long pants
- Shoes plus socks
- Protective eyewear

EPA REG. No. 55638-37

EPA EST. Nos. 67250-IL-1 

(Subscript refers to last 2 digits of lot number on container.)

Net Contents: 5 U.S. Pound Bag

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark.

Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.


Pesticide Storage: Store in a cool, dry place inaccessible to children.

Pesticide Disposal: Do not contaminate water when disposing of equipment washwaters. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.



97 APR -9 P2:31

REC'D EPA/OPP/DPD1



Ecogen Inc.
2005 Cabot Blvd. West, P.O. Box 3023
Langhorne, PA 19047-3023
215/757-1590 or 800/220-2135

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this section only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: coveralls, waterproof gloves, shoes plus socks, and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to the uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

Preharvest Interval: LEPINOX WDG may be applied to the crops listed in the APPLICATION RATE TABLE at any time, up to and on the day of harvest.

Mode of Action: After consuming a lethal dose of LEPINOX WDG, larvae will cease to feed, but may remain alive on foliage for several days before disappearing. Immediately after ingestion of LEPINOX WDG, larvae begin to move slowly, become discolored, shrivel and blacken prior to death.

MIXING INSTRUCTIONS

LEPINOX WDG may be applied with conventional ground, aerial or hand held application equipment with quantities of water sufficient to provide thorough coverage of infested plants. Do not apply this product through any type of irrigation system. To obtain a suitable mixture with water, add enough water to allow maximum agitation. With agitator running, slowly add in the LEPINOX WDG. Continue agitation. Add remainder of water and other spray materials and agitate until mixed. Maintain suspension while loading and spraying. Do not mix more LEPINOX WDG than can be used in a 12-hour period. Rinse and flush spray equipment thoroughly following each use. Do not contaminate water when disposing of equipment washwaters.

In order to make proper decisions on application rates to be used, follow the recommendations in the APPLICATION RATE TABLE.

APPLICATION INSTRUCTIONS

LEPINOX WDG is a bioinsecticide for use against the lepidopteran larvae listed in the APPLICATION RATE TABLE. Larvae must consume deposits of LEPINOX WDG to be affected. Always follow these directions:

- Make applications when larvae are still small (early instars) and actively feeding on foliage or other plant parts.
- Make applications before noticeable foliar damage occurs.
- Thorough spray coverage is essential for good insect control. For ground applications, directed drop nozzles should be used for certain vegetable crops.
- When insect infestations are heavy, use the higher label rates, shorten the spray interval, and/or use larger total spray volume to improve spray coverage.
- Applications should be repeated at an interval sufficient to maintain control, depending upon plant growth, insect pressure and weather conditions after spraying.
- For crops such as Fruits, Nuts and Vines, applications are often timed to stage of development and recommendations from local Extension personnel should always be followed.
- Local conditions may affect the use of LEPINOX WDG. Consult your State Agricultural Extension Specialist for specific recommendations related to local crop protection problems.
- Spray water/spray tank solutions should not exceed pH 8.0. If necessary, buffer water to near neutral pH.

HAND HELD EQUIPMENT

When using hand held equipment, mix 4 teaspoons per gallon of water or 3 pounds per 100 gallons of spray solution. Spray to wet, but not to runoff.

TANK MIX

Combinations of LEPINOX WDG with commonly used insecticides, fungicides or other spray tank adjuvants are generally not deleterious to performance. It is advisable to test physical compatibility by mixing all components in a small container in proportionate quantities prior to mixing in spray tank. This product cannot be mixed with any product containing a label prohibition against such mixing. No label dosage rate should be exceeded. Application must be made in accordance with the more restrictive of label limitation and precautions.

- For improved durability of spray deposits, a spreader/sticker approved for use on growing crops may be used for hard-to-wet crops such as cole crops.

APPLICATION RATE TABLE

I. VEGETABLES & COLE CROPS (Fresh and Processed)

Crop such as:		Insect Pest
Artichokes	Melons	Alfalfa looper
Arugala	(Cantaloupe,	Amyworm
Asparagus	Crenshaw,	Artichoke plume
Beans	Honeydew,	moth
Beets	Muskmelon,	Beet armyworm
Bok Choy	Watermelon,	Cabbage
Broccoli	(etc.)	budworm
Brussels sprouts	Napa	Cabbage looper
Cabbage	Okra	Cabbage
Cardoni	Onions	webworm
Carrots	Parsley	Celery leaflier
Cauliflower	Parsnips	Corn earworm
Celeriac	Peas	Cross-striped
Celery	Peppers	cabbageworm
Chick peas	Potatoes	Diamondback
Chicory	Pumpkins	moth
Chinese cabbage	Radishes	European corn
Collards	Rutabaga	borer
Cucumber	Salsify	Fall armyworm
Cucurbits	Shallots	Green
Dry bulb onions	Soybean foliage	cloverworm
Eggplants	Spinach	Imported
Escarole	Squash	cabbageworm
Endive	Sugar beets	Melonworm
Garlic	Sweet potatoes	Omnivorous
Green onions	Swiss chard	leafroller
Greens (Beets,	Tomatoes	Pickeworm
China, Dande-	Turnips	Rindworm
lion, Mustard,		complex
Turnip)		Saltmarsh
Horseradish		caterpillar
Kale		Soybean looper
Kohlrabi		Tobacco
Leeks		budworm
Lentils		Tomato fruitworm
Lettuce (Head,		Tomato
Leaf, Romaine)		hornworm
Malanga		Tomato pinworm
		Velvetbean
		caterpillar
		Yellowstriped
		armyworm

Rate/Acre: 1.0 - 2.0 pounds

II. HERBS & SPICES

Crop such as:	Insect Pest
Basil	Alfalfa looper
Chives	Amyworm
Cilantro	Diamondback moth
Dill	European corn borer
Oregano	Green cloverworm
Peppermint	Imported cabbageworm
Thyme	Loopers
	Saltmarsh caterpillar

Rate/Acre: 1.0 - 2.0 pounds

III. PASTURE & HAY CROPS

Crop such as:	Insect Pest
Alfalfa (hay & seed)	Alfalfa caterpillar
Pasture (grasses & hay)	Amyworm
Silage	Beet armyworm*
	European skipper
	Fall armyworm
	Loopers*
	Webworm
	Yellowstriped armyworm*

Rate/Acre: 1.0 - 2.0 pounds

*Product should be applied when early instar larvae first appear. If infestation persists, make a second application 7-10 days later. Combination of LEPINOX WDG with a contact insecticide is recommended for control of 4th and 5th instar larvae.

IV. FRUITS, NUT & VINE CROPS

Crop such as:	Insect Pest
Pome and Stone Fruit Trees: Apples Apricots Cherries Nectarines Peaches Pears Plums Prunes Quince	Cankerworm (Spring & Fall) Cherry fruitworm Eastern tent caterpillar Fall webworm Fruittree leafroller Greentree fruitworm Gypsy moth Navel orangeworm Obliquebanded leafroller Omnivorous leafroller Oriental fruit moth Pandemis leafroller Peach twig borer Redbanded leafroller Redhumped caterpillar Tortrix moth (Orange and Garden) Tufted apple budmoth Variegated leafroller Walnut caterpillar Western tent caterpillar
Nut Trees: Almonds Chestnuts Filberts Pecans Pistachios Walnuts	Citrus cutworm Filbert leafroller Filbert webworm Fruittree leafroller Hickory shuckworm Navel orangeworm Obliquebanded leafroller Omnivorous leafroller Pecan nut casebearer Peach twig borer Redhumped caterpillar Roughskinned cutworm Western tent caterpillar
Citrus	Amorbia Citrus cutworm Fruittree leafroller Omnivorous leafroller Orangedog
Small Fruit and Berries: Blackberries Blueberries Boysenberries Cranberries Currants Loganberries Raspberries Strawberries	Achema sphinx moth Armyworm Blackheaded fireworm Blueberry leafroller Cranberry girdler Fruittree leafroller Grape berry moth Gypsy moth Loopers Obliquebanded leafroller Omnivorous looper Tobacco budworm
Grapes	Grape berry moth Cherry fruitworm Grape leafroller Grapeleaf skeletonizer Green fruitworm Omnivorous leafroller Orange tortrix Saltmarsh caterpillar Yellowstriped armyworm
Tropical and Other Fruit: Avocados	Amorbia Loopers Orange tortrix Omnivorous leafroller Omnivorous looper Spanworm
Bananas	Banana skipper
Kwi	Omnivorous leafroller
Persimmons Pomegranate	Citrus cutworm Fall webworm Filbert webworm Omnivorous leafroller Redhumped caterpillar Tent caterpillar
Pineapple	Gummosos-Batrachedra commosae Thecla-Thecla basilides
Tropical fruits	Hornworms Leafrollers Loopers Omnivorous leafroller

Rate/Acre: 1.0 - 2.0 pounds

V. FIELD CROPS

Crop such as:	Insect Pest
Canola/Rape Seed Evening Primrose Meadow foam	Armyworm Diamondback moth Imported cabbageworm Loopers
Corn (Field, Sweet, Popcorn, Seed)	Armyworm European corn borer Fall armyworm Southwestern corn borer
Cotton	Beet armyworm Bollworm Cabbage looper Cotton leaf perforator Fall armyworm Saltmarsh caterpillar Tobacco budworm
Hops	Armyworm Loopers Obliquebanded leafroller Omnivorous leaflier Spotted cutworm
Jojoba	Looper (<i>Anacamptodes spp.</i>)
Peanuts	Fall armyworm Green cloverworm Loopers Podworm Velvetbean caterpillar
Rice	Armyworm Green cloverworm Loopers Saltmarsh caterpillar Velvetbean caterpillar
Safflower	Armyworm Loopers Saltmarsh caterpillar
Small Grains (Barley, Oats, Rye, Wheat, etc.)	Armyworm Loopers
Sorghum	European corn borer Fall armyworm Headworm Saltmarsh caterpillar Velvetbean caterpillar
Soybeans	Fall armyworm Green cloverworm Podworm Soybean looper Velvetbean caterpillar
Sunflowers	Banded sunflower moth Beet armyworm Headmoth Loopers Sunflower moth
Tobacco	Tobacco budworm Tobacco hornworm Loopers

Rate/Acre: 1.0 - 2.0 pounds

VI. COMMERCIAL FLOWERS & ORNAMENTAL PLANTS

Crop such as:	Insect Pest
Bedding plants Flowers (Greenhouse and field) Greenhouse Ornamentals Vegetables Container stock	Armyworm Azalea moth Beet armyworm Diamondback moth Elo moth (hornworm) Florida fern caterpillar lo moth Loopers Oleander moth Omnivorous leafroller Omnivorous looper Tobacco budworm

Rate/Acre: 1.0 - 2.0 pounds

VII. FOREST, SHADE TREE & NURSERY STOCK

Crop such as:	Insect Pest
Forest Shade trees Nursery trees	Bagworm Blackheaded budworm Browntail moth California oakworm Douglas fir tussock moth Elm spanworm Fall webworm Fruitree leafroller Greenstriped mapleworm Gypsy moth Jack pine budworm Mimosa webworm Pine butterfly Redhumped caterpillar Saddleback caterpillar Saddle prominent caterpillar Spring and Fall cankerworm Spruce budworm Tent caterpillar Tortrix Western tussock moth

Rate/Acre: 1.0 - 2.0 pounds

VIII. TURF

Crop such as:	Insect Pest
Turf	Armyworm Fall armyworm Sod webworm Tropical sod webworm

Rate/Acre: 1.0 - 2.0 pounds

WARRANTY AND CONDITIONS OF SALE

Ecogen warrants that this product conforms to the description on this label and is reasonably fit for the purposes stated on this label when used in accordance with the directions on this label under normal conditions of use.

ECOGEN MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

If this product is defective, Buyer's exclusive remedy shall be the replacement of the product, or if replacement is impracticable, refund of the purchase price. In no case will Ecogen be liable for incidental, consequential or special damages resulting from the handling, storage or use of this product.