

PM 25

55947-1

1077

AUG 31 1993

Dr. Jonathan Bryant
Sandoz Agro, Inc.
1300 East Touhy Ave.
Des Plaines, IL 60018

Dear Dr. Bryant:

Subject: Increase in Use Rate- Wheat
BANVEL HERBICIDE
EPA Reg. No. 55947-1
Your application dated Aug. 23, 1993

The amendment referred to above, submitted in connection with registration under FIFRA sec. 3(c)(7)(A), is acceptable provided that you:

1. Submit and/or cited all data or other material required for registration/reregistration of your product under FIFRA sec. 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.

2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

This acceptance of your label does not relieve you of any obligation to comply with the Worker Protection Standard (WPS). Under the WPS labeling regulations at 40 CFR part 158, subpart K, § 156.200(c)(3), you are prohibited from distributing or selling any product within the scope of the WPS requirements after April 21, 1992, without amended labeling accepted by the Agency.

2 of 3

-2-

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

Robert J. Taylor
Product Manager 25
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

5074

BANVEL^(R) Herbicide

SUPPLEMENTAL LABELING

EPA Reg. No. 55947-1

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE PRECAUTIONS IMPORTANT

Before using this product, read and carefully observe all applicable directions, restrictions, and precautionary statements on the container label and the Directions for Use booklet accompanying the BANVEL[®] Herbicide container.

SENSITIVE CROP PRECAUTIONS

BANVEL Herbicide may cause injury to desirable trees and plants, particularly beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes and other broadleaf plants when contacting their roots, stems or foliage. These plants are most sensitive to BANVEL Herbicide during their development or growing stage. **FOLLOW THE PRECAUTIONS LISTED ON THE BANVEL HERBICIDE LABEL WHEN USING BANVEL HERBICIDE.**

FALL SEEDED WHEAT

RATES AND TIMINGS

Application of BANVEL Herbicide may be made before, during or after planting to emerged and actively growing weeds. Use BANVEL Herbicide at 4 fluid ounces (0.125 lb.ai) per treated acre in wheat. BANVEL Herbicide is most effective when used in a tank mix with other herbicides. For best performance, make application when weeds are in the 2-3 leaf stage and rosettes are less than 2 inches across. Use the higher level of listed rate ranges when treating more mature weeds or dense vegetative growth.

BANVEL HERBICIDE MUST BE APPLIED TO FALL SEEDED WHEAT PRIOR TO THE JOINTING STAGE.

ACCEPTED
with COMMENTS
in EPA Letter Dated:

AUG 31 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

55947-1

11-27-66

FOR SUPPRESSION OF PERENNIAL BROADLEAF WEEDS IN FALL SEED WHEAT

PERIODS OF EXTENDED COLD AND WET WEATHER MAY ENHANCE THE POSSIBILITY OF CROP INJURY. DO NOT USE IF THE POTENTIAL FOR CROP INJURY IS NOT ACCEPTABLE.

Apply 8 fl. oz. of BANVEL Herbicide to winter wheat after it exceeds the 3 leaf stage. Do not apply with surfactants or oils as injury may occur. BANVEL Herbicide may be tank mixed with 2,4-D amine at 0.25 lbs. a.i./A after wheat begins to tiller.

For best control apply BANVEL Herbicide to actively growing bindweed. Applications may be made in the Fall following frost but before the first killing frost.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION.