

JAN 11 1994

Ms. Nancy Huebl
Sandoz Agric. U.S.A.
1300 East Touhy Ave.
Des Plaines, IL 60018

Dear Ms. Huebl:

Subject: Minor Changes
MonDak® Herbicide
EPA Reg. No. 55947-7
Your submission dated Dec. 14, 1993

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable and a stamped copy is enclosed for your records.

We note that this labeling does not bear the required worker protection statements as outlined in PR Notices 93-7 and 93-11. We are aware that you have submitted labels in accordance with the instructions in those Notices. The new protective language in those submissions must be melded into this label for any shipments released after the deadlines that have been established. You may opt to do so immediately if you are electing the self-verification option, or after acceptance by the special Agency Worker Protection Standard review group (WSP). Finished printed labels should be submitted after acceptance by the WSP and in accordance with their instructions.

Sincerely yours,

Robert J. Taylor
Product Manager 25
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

BEST AVAILABLE COPY

29 18

MonDak® Herbicide
For Commercial or Agricultural Use Only

ACTIVE INGREDIENTS:

Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)	16.0%
Dimethylamine salt of related acids	4.0%
Dimethylamine salt of 2-methyl-4- chlorophenoxy-acetic acid	32.6%
INERT INGREDIENTS:	47.4%

TOTAL 100.0%

Contains: 1.25 Pounds Per Gallon Dicamba
2.50 Pounds Per Gallon MCPA

ACCEPTED
with COMMENTS
in EPA Letter Dated:

SHAKE WELL BEFORE USING
READ THE LABEL
FOLLOW DIRECTIONS

Under the Federal Insecticide,
Fungicide, and Herbicide Act
as amended, for the
registered under...

Keep Out of the Reach of Children

55947-7

DANGER

STATEMENT OF PRACTICAL TREATMENT

If in eyes, flush with water for 15 minutes.
See Back Panel for additional precautionary statements.

Net Contents: 2 1/2 Gallons
EPA Reg. No. 55947-7
EPA Est. No. 37507-MT-01

SANDOZ AGRO, INC.
1300 EAST TOUHY AVENUE
DES PLAINES, ILLINOIS 60018

CONTAINER/HMONDAKA.93C

DECEMBER 1993

BEST AVAILABLE COPY

62

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER

Corrosive, causes eye damage. Do not get in eyes, or on clothing. Wear goggles or face shield when handling. Harmful if swallowed. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

If in eyes, flush with water for 15 minutes. Get medical attention.

If swallowed, drink 1 or 2 glasses of water and induce vomiting by sticking finger down back of throat. Never induce vomiting or give anything by mouth to an unconscious person.

If on skin, wash with plenty of soap and water. Get medical attention.

Environmental Hazards

Keep out of lakes, streams or ponds. Drift or run-off may adversely affect nontarget plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling MCPA pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

47 18

Apply this product only as directed on the label.

IMPORTANT

Do not apply MONDAK® herbicide directly to or allow spray drifts to come in contact with vegetables, flowers, grapes, tomatoes, potatoes, sugar beets, beans, or other desirable plants. (Coarse sprays are less likely to drift.). Do not contaminate domestic or irrigation water. Do not store in direct contact with fertilizer, seed, insecticides or fungicides. Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes. When container is empty, immediately wash thoroughly and destroy. Do not graze treated areas or harvest for dairy feed prior to crop maturity. Do not apply (except as recommended) or clean equipment on or near desirable trees or other plants, or on areas where roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

5718

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage or disposal.

STORAGE

Always use original container to store pesticides in a secured warehouse or storage building. Do not store near seeds, fertilizers, insecticides or fungicides. Store at temperature above 32° F. If allowed to freeze, warm to at least 40° and remix before using. Freezing does not alter this product. Containers should be opened in well-ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

6918

LIMITED WARRANTY AND LIABILITY

NOTICE: Read this Limited Warranty and Liability before buying or using this product. If the terms are not acceptable, return it at once unopened.

It is critical that this product be used and mixed only as specified on the label. The laws of a State may make some or all of this paragraph inapplicable or may give you rights in addition to your rights hereunder. Except, to the extent prohibited by applicable law, the exclusive remedy of the user or buyer and the limit of liability of this Company or any other Seller for any and all losses, personal injuries or damages resulting from the use of this product, shall be the purchase price paid by the user or buyer for the quantity of product involved. Except to the extent prohibited by State law, there is no warranty, and this Company and other Sellers disclaim all liability for losses, personal injury or damages: (i) arising from any use of this product in a manner or for a purpose not recommended in its label directions, or from mixing this product before use with any substance except as recommended by the product's label, (ii) arising from handling or storage in violation of label instructions, (iii) for all indirect, special or consequential damages, (iv) when not reported to this Company within one year of discovery, and (v) arising from product not used within the label-designated shelf life of four years from date of purchase, whichever first occurs. **THERE ARE NO IMPLIED WARRANTIES AND NO WARRANTIES OF MERCHANTABILITY OR FITNESS.**

79 18
7

MonDak® Herbicide
For Commercial or Agricultural Use Only

ACTIVE INGREDIENTS:

Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)	16.0%
Dimethylamine salt of related acids	4.0%
Dimethylamine salt of 2-methyl-4- chlorophenoxy-acetic acid	32.6%
INERT INGREDIENTS:	47.4%
 TOTAL	 100.0%

Contains: 1.25 Pounds Per Gallon Dicamba
2.50 Pounds Per Gallon MCPA

**SHAKE WELL BEFORE USING
READ THE LABEL
FOLLOW DIRECTIONS**

Keep Out of the Reach of Children

DANGER

STATEMENT OF PRACTICAL TREATMENT
If in eyes, flush with water for 15 minutes.
See Back Panel for additional precautionary statements.

Net Contents: 2 1/2 Gallons
EPA Reg. No. 55947-7
EPA Est. No. 37507-MT-01

**SANDOZ AGRO, INC.
1300 EAST TOUHY AVE.
DES PLAINES, IL 60018**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER

Corrosive, causes eye damage. Do not get in eyes, or on clothing. Wear goggles or face shield when handling. Harmful if swallowed. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

If in eyes, flush with water for 15 minutes. Get medical attention.

If swallowed, drink 1 or 2 glasses of water and induce vomiting by sticking finger down back of throat. Never induce vomiting or give anything by mouth to an unconscious person.

If on skin, wash with plenty of soap and water. Get medical attention.

Environmental Hazards

Keep out of lakes, streams or ponds. Drift or run-off may adversely affect nontarget plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling MCPA pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Apply this product only as directed on the label.

IMPORTANT

Do not apply MONDAK® herbicide directly to or allow spray drifts to come in contact with vegetables, flowers, grapes, tomatoes, potatoes, sugar beets, beans, or other desirable plants. (Coarse sprays are less likely to drift.). Do not contaminate domestic or irrigation water. Do not store in direct contact with fertilizer, seed, insecticides or fungicides. Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes. When container is empty, immediately wash thoroughly and destroy. Do not graze treated areas or harvest for dairy feed prior to crop maturity. Do not apply (except as recommended) or clean equipment on or near desirable trees or other plants, or on areas where roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

BRUSH AND BROADLEAF WEED CONTROL

SPECIES CONTROLLED

When used as directed, MONDAK® Herbicide will control or suppress many herbaceous broadleaf weeds (annual, biennial, and perennial) as well as many unwanted woody plant and vine species. Species controlled include:

ANNUALS

Buckwheat,	Wild Lambsquarters
Carpetweed	Morningglory
Chickweed	Mustard
Clover	Pigweed
Cocklebur	Purslane
Daisy, English	Ragweed
Henbit	Smartweed
Knawel	Velvetleaf

BIENNIALS

Ragwort, Tansy	Thistle, Musk
----------------	---------------

PERENNIALS

Bindweed, Field	Milkweed
Carrot, Wild	Ragweed, Perennial
(Queen Anne's Lace)	Sorrel, Sheep
Dock, Curly	Spurge, Leafy
Dogfennel	Thistle, Canada
Knapweed, Russian	Toadflax, Dalmatian

WHEAT (not underseeded to legumes)

For the control of "hard to kill" weeds (phenoxy resistant) including:

Bedstraw	Henbit
Corn cockle	Knawel (German moss)
Cow cockle	Ladysthumb
Corn buttercup (crowfoot)	Miners lettuce
Dogfennel	Prickly lettuce
Fiddleneck (tarweed)	Smartweed
Gromwell	Wild buckwheat

And for control of other broadleaf weeds including:

Canada thistle (top growth)	Perennial Sowthistle
Cocklebur	(top growth)
Dandelion	Pigweed (redroot)
Dragonhead (mint)	Puncturevine
Field pennycress (fanweed)	Ragweed (common)
Field pepperweed	Russian thistle
Hempnettle	Shepherdspurse
Knotweed	Sunflower
Kochia	Vetch
Lambsquarters	Wild radish
Mustard (annual)	and others

For ground application use 5 to 20 gallons of water per acre.

For aerial application use 3 to 5 gallons of water per acre.

Spring Seed Wheat: For the control of the listed weeds, apply 0.8 pint of MONDAK® herbicide (1/8 pound of dicamba plus 1/4 pound of MCPA) per acre. One gallon of MONDAK® herbicide treats 10 acres. Apply as a foliar spray when the wheat is in the 2 to 5 leaf stage.

Fall Seeded Wheat: For control or suppression of the listed weeds, apply 0.8 to 1.2 pints of MONDAK® herbicide (1/8 to 3/16 pound of dicamba plus 1/4 to 3/8 pound of MCPA) per acre. One gallon of MONDAK® herbicide treats 7 to 10 acres. Use the higher rates on hard to kill or more mature weeds. Apply as a foliar spray immediately after winter dormancy and before the wheat begins to joint.

11 3 18
3

NONCROPLAND USE: (Roadsides, fence rows and similar areas.)

For top growth control of field bindweed, Canada thistle, perennial sowthistle, Russian knapweed and leafy spurge, apply 3.2 to 6.4 pints of MONDAK® herbicide (1/2 to 1 pound of dicamba and 1 to 2 pounds of MCPA) per acre. For spot treatment apply 1 teaspoonful of MONDAK® herbicide per 1 to 2 square rods. Apply as a foliar spray to actively growing plants.

GRASS SEED PRODUCTION

(Lawn-type fescues, perennial bentgrasses, perennial ryegrasses, and Kentucky bluegrass.)

FOR USE ONLY IN THE STATES OF IDAHO, OREGON AND WASHINGTON.

Apply to grass stands established at least one year prior to treatment.

For control of:

Corn cockle	Henbit
Cornflower	Knawel
(Bachelor's button)	Lambsquarters
Corn spurry	Pepperweed
Chickweed	Pineappleweed
Curly dock	Prickly lettuce
Dandelion	Red sorrel
Dogfennel	Shepherdspurse
False dandelion	Vetch (common)
Field pennycress (tarweed)	Wild radish
Filaree	

Apply 1 1/2 pints of MONDAK® herbicide (3 3/4 ounces dicamba acid equivalent plus 7 1/2 ounces MCPA acid equivalent) per acre. Apply to early seedling weeds (4" or less in height after winter dormancy up to the early boot stage of the grass. For ground application, use a minimum of 20 gallons of total spray per acre.

WOODY BRUSH AND VINES

Alder
Ash
Aspen
Basswood
Beech
Birch
Blackberry
Cherry
Creeper, Virginia
*Creosotebush
Cucumbertree
Dogwood
Elderberry
Elm
Gum
Hawthorn
Hemlock
Honeysuckle
Ivy, Poison
Kudzu
Locust
Maple
Oak
Olive, Russian

Persimmon
Pine
Plum, Wild
Poplar
Puncturevine
Raspberry
Redcedar, Eastern
Redvine
Rose, Multiflora
Sagebrush
Sassafras
Schinus
(Florida Holly,
Brazil Peppertree,
Christmas-berry)
Serviceberry
Snowberry
Spruce
Sumac
Sycamore
Trumpetcreeper
Waxmyrtle
Willow
Witchhazel
Yaupon

*Suppression

APPLICATION DIRECTIONS

Regardless of the species to be controlled, spray volumes should be high enough to allow for good spray coverage. Make applications when weeds and brush are actively growing. The addition of surfactants can increase control. Biennials are best controlled when treated in the rosette stage. Regrowth may occur on resistant species. Retreatments may be made as needed; however, do not exceed a total of 4.8 gallons (6 lbs. a.i. dicamba + 12 lbs. a.i. MCPA) of MONDAK® Herbicide per treated acre during a growing season. For control of hard-to-kill woody species, a tankmix may be necessary. Tankmix recommendations are for use only in states where the tank mix product and application site are registered. Read and follow all label directions.

HERBACEOUS BROADLEAF WEED CONTROL

Apply 3/4 to 3 1/4 quarts MONDAK Herbicide in 20 to 100 gallons of water per treated acre. 3/4 to 2 quarts of MONDAK® Herbicide is recommended for annuals, 2 to 3 1/4 quarts for biennials and easy-to-kill perennials, and 3 1/4 quarts for established perennials. Do not apply more than 2 1/3 gallons of product per treated acre.

WOODY BRUSH AND VINE CONTROL

High-Volume Foliar Spot Applications

Mix 1 to 2 1/3 gallons of MONDAK® Herbicide in enough water to make 100 gallons of spray mix. Spray volume applied will depend on the size and density of the brush to be treated, but do not apply more than 2 1/3 gallons of product per treated acre. Direct the spray to treat all foliage, stems, and root collars to wet.

WOODY BRUSH AND VINE CONTROL

Broadcast Applications with Ground Equipment

Apply 1 to 2 1/3 gallons MONDAK® Herbicide in 20 to 100 gallons of water per treated acre. Spray volume applied will depend on the size and density of the brush to be treated, but do not apply more than 2 1/3 gallons of product per treated acre. Spray all foliage, stems, and root collars to wet.

AERIAL APPLICATIONS

Aerial applications may be made to control either herbaceous or woody plants. Apply 1 to 3 1/4 quarts MONDAK® Herbicide (for herbaceous weeds) or 1 to 2 1/3 gallons MONDAK Herbicide (for woody brush and vines) in 5 to 40 gallons of water per treated acre. Coverage is important, so increase spray volume when treating dense stands of brush or weeds. Do not apply more than 2 1/3 gallons of product per treated acre.

TANK MIX TREATMENTS

TANK MIX RECOMMENDATIONS ARE FOR USE ONLY IN STATES WHERE THE TANK MIX PRODUCT AND APPLICATION SITE ARE REGISTERED. READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, AND OTHER RESTRICTIONS. For broader spectrum control, MONDAK® Herbicide may be tank mixed with one or more of the following Herbicides for noncropland uses (e.g. railroad, highway, pipeline, etc.) including forest management applications, if permitted by product labeling (e.g. MCPA). Add water to the spray tank prior to the addition of the tank mix products. Do not premix concentrates.

Herbicide	Rate (lbs. a.i./treated acre)
Asulam (Asulox®)	2.92 to 6.68
Bromacil (Hyvar®)	1.6 to 24.0
Chlorsulfuron (Telar®)	0.1875 to 2.25 oz.**
Dicamba (Banvel®)	0.25 to 8.0***
Diuron (Karmex®)	4.0 to 48.0
Fosamine ammonium (Krenite®)	6.0 to 12.0
Glyphosate (Roundup®)	0.75 to 3.75
Hexazinone (Velpar®)	0.675 to 10.8
+ Imazapyr (Arsenal®)	0.5 to 1.5
MSMA	1.0 to 2.475
Picloram* (Tordon®)	2.0 to 3.0
Simazine* (Princep®)	4.8 to 40.0
Sulfometuron methyl (Oust®)	0.75 to 9.0 oz.**
Tebuthiuron (Spike®)	1.0 to 16.0
Triclopyr (Garlon®)	0.75 to 12.0
2,4-D	0.475 to 16.92****
2,4-DP	0.5 to 11.1

Due to variations that may occur in formulated products and specific use ingredients (e.g. water supplies) a COMPATIBILITY TEST as described on this label is recommended prior to actual tank mixing.

*Restricted use pesticides limited to certified applicators

**Telar = 0.1875 to 2.25 oz. a.i./A (0.25 to 3.0 oz. product/A)

Oust = 0.75 to 9.0 oz. a.i./A (1.0 to 12.0 oz. product/A)

***Do not exceed a total of 8 lbs. a.i. dicamba per treated acre per growing season. (MONDAK® Herbicide contains 1.25 lb. a.i. dicamba per gallon).

****Use of extremely hard water (500 ppm) may cause this product to form a precipitate when used in a tank mix with MONDAK® Herbicide. Do not exceed 16 lbs. total MCPA acid equivalent per treated acre per growing season. (MONDAK® Herbicide contains the equivalent of 2.5 lbs./gal. MCPA acid).

+ Not registered for use in California.

COMPATIBILITY TEST

Before mixing in the spray tank, it is advisable to test compatibility by mixing all components in a small container in proportionate quantities (see following table).

**Amount of Herbicide to Add to
One Pint of Spray carrier
(Assuming Volume is 25 Gallons per Acre)**

HERBICIDE FORMULATIONS	RATE PER ACRE	LEVEL TEASPOONS
Dry	1 lb.	1 ½
Liquid	1 Pt.	2

If herbicides do not ball-up or form flakes, sludge, gels, oily films or layers, or other precipitates, then the tested components are compatible. Usually incompatibility in any of the above described forms will occur within 5 minutes after mixing.

If components are incompatible, the use of a compatibility agent is recommended. Rerun the above COMPATIBILITY TEST with a suitable compatibility agent (¼ teaspoon is equivalent to 2 pints per 100 gallons of spray solution).

PROCEDURE FOR CLEANING SPRAY EQUIPMENT

The steps listed below are suggested for thorough cleaning of spray equipment following applications of MONDAK® Herbicide or tank mixes of MONDAK® Herbicide plus MCPA amine.

- 1) Hose down thoroughly the inside as well as outside surfaces of equipment while filling the spray tank half full of water. Flush by operating sprayer until the system is purged of the rinse water.
- 2) Fill tank with water while adding 1 quart of household ammonia for every 25 gallons of water. Operate the pump to circulate the ammonia solution through the sprayer system for 15 to 20 minutes and discharge a small amount of the ammonia solution through the boom and nozzles. Let the solution stand for several hours, preferable overnight.
- 3) Flush the solution out of the spray tank through the boom.
- 4) Remove the nozzles and screens and flush the system with two full tanks of water.

The steps listed below are suggested for thorough cleaning of spray equipment used to apply MONDAK® Herbicide as a tank mix with wettable powders (WP), emulsifiable concentrates (EC), or other types of water-dispersible formulations. MONDAK® Herbicide tank mixes with water-dispersible formulations require the use of a water/detergent rinse.

5) **Complete step 1.**

6) **Fill tank with water while adding 2 lbs. of detergent for every 40 gallons of water. Operate the pump to circulate the detergent solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.**

7) **Flush the detergent solution out of the spray tank through the boom.**

8) **Repeat step 1, and follow with steps 2, 3 and 4.**

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage or disposal.

STORAGE

Always use original container to store pesticides in a secured warehouse or storage building. Do not store near seeds, fertilizers, insecticides or fungicides. Store at temperature above 32° F. If allowed to freeze., warm to at least 40° and remix before using. Freezing does not alter this product. Containers should be opened in well-ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more that two pallets high.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

17 9 15

Arsenal® is a registered trademark of American Cyanamid Company.
Princep® is a registered trademark of Ciba-Geigy Corporation.
Garlon® and Tordon® are registered trademarks of DowElanco Company.
Hyvar®, Karmex®, Krenite®, Oust®, Telar®, and Velpar® are registered trademarks of E.I. duPont de Nemours & Co., Inc.
Spike® is a registered trademark of DowElanco.
Roundup® is a registered trademark of Monsanto Agricultural Products Co.
Asulox® is a registered trademark of Rhone-Poulenc Chemical Company.
Banvel® is a registered trademark of Sandoz Ltd.
Mondak® is a registered trademark of Sandoz Agro Inc.

LIMITED WARRANTY AND LIABILITY

NOTICE: Read this Limited Warranty and Liability before buying or using this product. If the terms are not acceptable, return it at once unopened.

It is critical that this product be used and mixed only as specified on the label. The laws of a State may make some or all of this paragraph inapplicable or may give you rights in addition to your rights hereunder. Except, to the extent prohibited by applicable law, the exclusive remedy of the user or buyer and the limit of liability of this Company or any other Seller for any and all losses, personal injuries or damages resulting from the use of this product, shall be the purchase price paid by the user or buyer for the quantity of product involved. Except to the extent prohibited by State law, there is no warranty, and this Company and other Sellers disclaim all liability for losses, personal injury or damages: (i) arising from any use of this product in a manner or for a purpose not recommended in its label directions, or from mixing this product before use with any substance except as recommended by the product's label, (ii) arising from handling or storage in violation of label instructions, (iii) for all indirect, special or consequential damages, (iv) when not reported to this Company within one year of discovery, and (v) arising from product not used within the label-designated shelf life of four years from date of purchase, whichever first occurs. THERE ARE NO IMPLIED WARRANTIES AND NO WARRANTIES OF MERCHANTABILITY OR FITNESS.