hancy number sandoz Agro, Inc.
1300 rast rouby Avenue per Flaines, In 60015

near ms. nuebl:

Subject: panVel® 720 Herbicide

EPA Registration No. 55947-20

Application Dated August 9, 1995, Requested

Amendments To Reduce Rates of Dicamba (Not To

Exceed 2 lbs./A, Acid Equivalent), To

Add Limits of Restrictions To Reduce Exposure
To Dicamba and Residues of Dicamba; to Add

To List of Weeds Control

The proposed amendments to the subject posticide product registration as listed above have been reviewed and found acceptable under the Federal Insecticide, Fundicide and Rodenticide act as amended, provided nations:

o sepuit rive (5) printed copies or your rinal printed labeling before you release the product for shipment.

It these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFPA, section b(a). Your release for shipment of the product constitutes acceptance of this condition.

A stumped copy of the labeling is enclosed for your records.

submit an explanation for the increased restrictions to exposures from the use-patterns claimed on the label. If analytical data support the more restrictive ase of this product product, you should submit the data. The record for this product should sear an explanation for the requested amenoments.

Sincerely yours,

Joanne I. Miller Product Manager (23) Fungicide-herbicide Brunch Redistration Division (750%)

	EUG LOSULE							
SYMBOL								
SURNAM	F. N							·
DATE	Witteom: w	i skerrear	13:05-11	9.5				
EPA For	m 1320-1 (12-70)	<u>. </u>	BEST	COPY	AVAIL	ABLE	.LOFF1CI	IAL FILE COPY

Ve har & Godien

REGULATORY LABEL DATA SHEET

(I:\labels\tools\covrshet.wpt)

File Name:

PBAN720A.95A

Data Sheet Date:

July 1995

Product:

BANVEL 720 HERBICIDE

EPA Reg #:

55947-20

VETERAN 720

Approved:

(RIVERDALE)

Supercedes:

8/8/94

OVERVIEW: Reduce rates - 2 lb/A dicamba maximum use rate (stewardship on all dicamba labels)

Page #	Description of Changes
4	Grazing and feeding restrictions for dairy animals revised to match other dicamba labels.
6	Some woody brush and vines species changed from control to suppression: Blackberry, Dogwood, Hawthorn, Wild Plum, Eastern Redcedar, Multiflora Rose and Yaupon
7-8	Rates revised to 2 lb dicamba (ai)/A/yr
8	"Imazapyr is the proposed common name for Arsenal" removed from footnote **.

REGULATORY:

Nancy Huebl/J. Bryant

CREATIVE SERVICES:

QUALITY ASSURANCE: Fred Mitchell

PRODUCT DEVELOPMENT:

Luke Bozeman

MARKETING: Leo Schoenhofen

EPA PRODUCT MANAGER: J. Miller/PM23

ACCEPTED with COMMENTS In EPA Letter Dated

SEP 12

Under the Federal Insecticide, Fundicide, and Rodenticide Act as amended, for the posticide giotograd under EPA Reg. No.

BANVEL® 720 HERBICIDE

WATER SOLUBLE INDUSTRIAL HERBICIDE FOR CONTROL OF BRUSH AND BROADLEAF WEEDS ON RIGHTS-OF-WAY, FOREST BRUSH, FENCE ROWS, NON-IRRIGATION DITCHBANKS, WASTELAND, AND SIMILAR NONCROPLAND AREAS

Active Ingredients:

Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)	12.82%
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid**	24.58%
Inert Ingredients:	<u>62.60%</u>
TOTAL	100.00%

^{*} This product contains 10.6% 3,6-dichloro-<u>o</u>-anisic acid (dicamba) or 1 lb per gallon (120 g/L).

SHAKE WELL BEFORE USING KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional precautionary statements

EPA Reg. No. 55947-20

Net Contents:

EPA Est. No. Sandoz Agro, Inc. Des Plaines, IL 60018

[&]quot;This product contains 20.4% 2,4-dichlorophenoxyacetic acid (2,4-D) or 1.9 lb per gallon (228 g/L).

^{) &}quot; Isomer specific by AOAC method 978.05, 15th Edition.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. If swallowed, drink 1 or 2 glasses of water and induce vomiting by sticking finger down back of throat. Never induce vomiting or give anything by mouth to an unconscious person. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. In case of contact, wash exposed skin with plenty of soap and water; for eyes flush with water for 15 minutes and get medical attention.

When mixing, loading, or applying this product or repairing or cleaning equipment used with this product, wear eye protection (face shield or safety glasses), chemical-resistant gloves, long sleeved shirt, long pants, socks and shoes. For aerial applicators in enclosed cockpits and applicators applying this product from a tractor that has a completely enclosed cab, eye protection is not required.

Wash hands, face and arms with soap and water as soon as possible after mixing, loading, or applying this product. Wash hands, face and arms with soap and water before eating, smoking, or drinking. Wash hands and arms before using toilet. After work, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing and loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry. Remove saturated clothing as soon as possible and shower.

SANVEL 720/55947-20 sban720a.95a July 1995

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply this product only as directed on label.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use on plants in non-crop and non-timber areas only. Not for use on trees, crops, or other plants being grown for sale or other commercial use, or for commercial seed production, or for the production of timber or wood products, or for research purposes.

John profes

The following directions apply to all uses of BANVEL® 720 Herbicide.

Do not contaminate irrigation ditches or water used for domestic purposes.

Do not apply this product through any type of irrigation system.

SENSITIVE CROP PRECAUTIONS

BANVEL 720 Herbicide may cause injury to desirable trees and plants, particularly beans, cotton, flowers, fruit trees, grapes, prnamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes and other broadleaf plants when contacting their roots, stems or foliage. These plants are most sensitive to BANVEL 720 Herbicide during their development or growing stage. FOLLOW THE PRECAUTIONS LISTED BELOW WHEN USING BANVEL 720 HERBICIDE.

Do not treat areas where either possible downward movement into the soil or surface washing may cause contact of BANVEL 720 Herbicide with the roots of desirable plants such as trees and shrubs.

Avoid making application when spray particles may be carried by air currents to areas where sensitive crops and other desirable plants are growing. Always make application where there is some air movement in order to determine the direction and distance of possible spray drift. Leave an adequate buffer zone between the area to be treated and desirable sensitive plants. Coarse sprays are less likely to drift out of target area.

Do not use aerial equipment to apply BANVEL 720 Herbicide when sensitive crops and plants are growing in the vicinity of area to be treated.

Select nozzles designed to produce minimal amounts of fine spray particles. Drift control agents and equipment designed to minimize fine spray particles may be used.

Do not graze dairy animals on treated areas within 7-days after application.

Do not graze dairy animals on treated areas until 21 days after application for 1 gallon (1 lb ai) or 40 days for up to 2 gallons (2 lb ai).

Remove meat animals from treated areas-3 30 days prior to slaughter.

Do not out grass for hay within 30 days of application.

Do not cut grass for hay within 51 days of application of 1 gallon (1 lb ai) or within 70 days for 2 gallons (2 lb ai).

To avoid injury to desirable plants, equipment used to apply BANVEL 720 Herbicide should be thoroughly cleaned (see PROCEDURE FOR CLEANING SPRAY EQUIPMENT on pages ---) before reusing to apply any other chemicals.

Consult your local or state authorities for possible application restrictions and acvice concerning these and other special local use situations.

John Chart

BANVEL 720 Water Soluble Industrial Herbicide for Brush and Broadleaf Weed Control is recommended for use on rights-of-way (utility, railroad, highway, pipeline), nonselective forest brush control, fence rows, non-irrigation ditchbanks, wasteland and similar non-cropland areas.

MIXING AND APPLICATION

If this container has a volume less than 5 gallons, all persons engaged in open pouring of this product must also wear coveralls or a chemical-resistant apron.

If this container has a volume of 5 gallons or more, a mechanical system (probe and pump) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

BANVEL 720 Herbicide is a water-soluble formulation which can be applied using water or oil/water emulsions as the carrier. Herbicide adjuvants or other spray additives (emulsifiers, surfactants, wetting agents, drift control agents, or penetrants) may be used for wetting, penetration, or drift control. Read and follow all use recommendations and precautions on the label(s) of product(s) used. A COMPATIBILITY TEST (page -- of this booklet) should be made prior to tank mixing.

BANVEL 720 Herbicide may be applied with ground or aerial equipment for broadcast treatments, or with handgun, backpack, or similar types of equipment for treatment of individual clumps or small areas (spot treatments).

NOTE: To prepare oil/water emulsions, half fill spray tank with water. Add a suitable emulsifier. Then, with constant agitation, slowly add the appropriate amount of herbicide followed by the necessary oil (or slowly add a premix of emulsifier and oil). Finally, fill the rest of the tank with water. Maintain vigorous agitation during spray operation to prevent oil and water from forming separate layers.

SPECIES CONTROLLED

When used as directed, BANVEL 720 Herbicide will control or suppress many herbaceous broadleaf weeds (annual, biennial, and perennial) as well as many unwanted woody plant and vine species. Species controlled include:

ANNUALS

Buckwheat, Wild Carpetweed Chickweed Clover Cocklebur Daisy, English

Lambsquarters Morningglory Mustard **Pigweed** Purslane Ragweed **Smartweed** Velvetleaf

Henbit Knawel

BIENNIALS

Thistle, Musk

Ragwort, Tansy

PERENNIALS

Bindweed, Field Carrot, Wild

Milkweed

(Queen Anne's Lace)

Ragweed, Perennial Sorrel, Sheep

Dock, Curly Dogfennel

Spurge, Leafy Thistle, Canada

Knapweed, Russian

Toadflax, Dalmation

WOODY BRUSH AND VINES

Alder Ash Aspen Basswood

Pine

Beech Birch

*Plum, Wild

Persimmon

Poplar

Puncturevine Raspberry

*Blackberry

Redced r. Eastern

Cherry

Redvine

Creeper, Virginia *Creosotebush

*Rose, Multiflora

Cucumbertree

Sagebrush Sassafras

*Dogwood > Elderberry

Schinus

Elm

(Florida Holly, Brazil Peppertree.

Gum *Hawthorn > Hemlock

Christmas-berry) Serviceberry Snowberry

Honeysuckle Ivy, Poison Kudzu

Spruce Sumac Sycamore

Locust Maple

Trumpetcreeper

Qak Olive, Russian Waxmyrtle Willow Witchhazel *Yaupon

^{*}Suppression

DIRECTIONS FOR USE

Regardless of the species to be controlled, spray volumes should be high enough to allow for good spray coverage. Make applications when weeds and brush are actively growing. The addition of surfactants can increase control. Biennials are best controlled when treated in the rosette stage. Regrowth may occur on resistant species. Retreatments may be made as needed; however, do not exceed a total of 6 2 gallons (6 2 lb ai dicamba + 12 3.8 lb ai 2,4-D) of BANVEL 720 Herbicide per treated acre during a growing season. To control additional weed species, BANVEL 720 Herbicide may be tank mixed with any of the products listed on pages ----.

HERBACEOUS BROADLEAF WEED CONTROL: Apply 1 to 4 quarts BANVEL 720 Herbicide in 20 to 100 gallons of water per treated acre. 1 to 2 quarts of BANVEL 720 Herbicide is recommended for annuals, 2 to 4 quarts for biennials and easy-to-kill perennials, and 4 quarts for established perennials. Do not apply more than 2 gallons of product per treated acre.

BRUSH AND VINE CONTROL - High Volume Foliar Spot Applications: Mix 1 to 2 gallons of BANVEL 720 Herbicide in enough water to make 100 gallons of spray mix. Spray volume applied will depend on the size and density of the brush to be treated, but do not apply more than 2 gallons of product per treated acre. Direct the spray to treat all foliage, stems, and root collars to wet.

BRUSH AND VINE CONTROL - Broadcast Applications with Ground Equipment: Apply 1 to 2 gallons BANVEL 720 Herbicide in 20 to 100 gallons of water per treated acre. Spray volume applied will depend on the lize and density of the brush to be treated, but do not apply more than 2 gallons of product per treated acre. Spray all foliage, stems and root collars to wet.

AERIAL APPLICATIONS: Aerial applications may be made to control either herbaceous or woody plants. Apply 1 to 4 quarts BANVEL 720 Herbicide (for herbaceous weeds) or 1 to 2 gallons BANVEL 720 Herbicide (for woody brush and vines) in 5 to 40 gallons of water per treated acre. Coverage is important, so increase spray volume when treating dense stands of brush or weeds. Do not apply more than 2 gallons of product per treated acre.

TANK MIX TREATMENTS

READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, AND OTHER RESTRICTIONS. For broader spectrum control, BANVEL 720 Herbicide may be tank mixed with one or more of the following herbicides for noncropland uses (e.g. railroad, highway, pipeline, etc.) including forest management applications, if parmitted by product labeling (e.g. 2,4-D). Add water to the spray tank prior to the addition of the tank mix products. Do not premix concentrates.

Rate Herbicide (lb ai/treated acre) 1.8 to 8.0 amitrol* 2.92 to 6.68 asulam (Asulox®) 4.77 to 39.96 atratol 1.6 to 24.0 bromacil (Hyvar®) 2.0 to 3.0 chlorflurecol (Maintain®) chlorsulfuron (Telar®) 0.1875 to 2.25 oz *** 4.25 to 12.75 dalapon 0.25 to 8.0 1.75 **** dicamba (Banvel) 0.5 to 1.0 diquat diuron (Karmex®) 4.0 to 48.0 fenac (Fenatrol®) 4.5 to 18.0 fosamine ammonium 6.0 to 12.0 (Krenite®)) glyphosate (Roundup®) 0.75 to 3.75 hexazinone (Velpar®) 0.675 to 10.8 imazapyr** (Arsenal®) 0.5 to 1.5 Limit®** 0.625 maleic hydrazide 2.25 to 4.5 (Royal Slo-Gro®) mefluidide (Embark®) 0.25 to 1.0 **MSMA** 1.0 to 2.475 paraquat* 0.5 picloram* (Tordon®) 2.0 to 3.0 simazine* (Princep®) 4.8 to 40.0 0.75 to 9.0 oz*** sulfometuron methyl (Oust®) 1.0 to 16.0 tebuthiuron (Spike®) triclopyr (Garlon®) 0.75 to 12.0 2,4-D 0.475 to 3.5**** 2,4-DP 0.5 to 11.1

Due to variations that may occur in formulated products and specific use ingredients (e.g. water supplies) a COMPATIBILITY TEST as described on page -- of this booklet is recommended prior to actual tank mixing.

Limit does not have a common name.

*** Telar = 0.1875 to 2.25 oz ai/A (0.25 to 3.0 oz product/A)

Oust = 0.75 to 9.0 oz ai/A (1.0 to 12.0 oz product/A)

**** Do not exceed a total of-8 2 lb ai dicamba per treated acre per growing season.
(BANVEL 720 Herbicide contains 1 lb ai dicamba per gallon.)

***** Use of extremely hard water (500 ppm) may cause this product to form a precipitate when used in a tank mix with BANVEL 720 Herbicide. Do not exceed 4 lb total 2,4-D acid equivalent per acre per application per site.

Do not exceed 16 lb total 2,4-D acid equivalent per treated acre per growing season. (BANVEL 720 Herbicide contains the equivalent to 1.9 lb/gal 2,4-D acid.)

^{*} Restricted use pesticides limited to certified applicators.

^{• *} Imazapyr is the proposed common name for Arsenal.

COMPATIBILITY TEST

Before mixing in the spray tank, it is advisable to test compatibility by mixing all components in a small container in proportionate quantities (see following table).

Amount of Herbicide to Add to One Pint of Spray Carrier (Assuming Volume is 25 Gallons per Acre)

HERBICIDE	RATE	LEVEL	
FORMULATIONS	PER ACRE	TEASPOONS	
Dry	1 lb	1 ½	
Liquid	1 pt	1/2	

If herbicide(s) do not ball-up or form flakes, sludge, gels, oily films or layers, or other precipitates, then the tested components are compatible. Usually incompatibility in any of the above described forms will occur within 5 minutes after mixing.

If components are incompatible, the use of a compatibility agent is recommended. Rerun the above COMPATIBILITY TEST with a suitable compatibility agent (1/4 teaspoon is equivalent to 2 pints per 100 gallons of spray solution).

PROCEDURE FOR CLEANING SPRAY EQUIPMENT

The steps listed below are suggested for thorough cleaning of spray equipment following applications of BANVEL 720 Herbicide or tank mixes of BANVEL 720 Herbicide plus 2,4-D amine.

- 1) Hose down thoroughly the inside as well as outside surfaces of equipment while filling the spray tank half full of water. Flush by operating sprayer until the system is purged of the rinse water.
- 2) Fill tank with water while adding 1 quart of household ammonia for every 25 gallons of water. Operate the pump to circulate the ammonia solution through the sprayer system for 15 to 20 minutes and discharge a small amount of the ammonia solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
- Flush the solution out of the spray tank through the boom.
- 4) Remove the nozzles and screens and flush the system with two full tanks of water.

15 1 14

SANVEL 720/55947-20 cban720a 95a July 1995

The steps listed below are suggested for thorough cleaning of spray equipment used to apply BANVEL 720 Herbicide as a tank mix with wettable powders (WP), emulsifiable concentrates (EC), or other types of water-dispersible formulations. BANVEL 720 Herbicide tank mixes with water-dispersible formulations require the use of a water/detergent rinse.

- 5) Complete step 1.
- 6) Fill tank with water while adding 2 lb of detergent for every 40 gallons of water.

 Operate the pump to circulate the detergent solution through the boom and nozzles.

 Let the solution stand for several hours, preferably overnight.
- 7) Flush the detergent solution out of the spray tank through the boom.
- 8) Repeat step 1, and follow with steps 2, 3 and 4.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage or disposal.

STORAGE

Store in original container in a well-ventilated area separately from fertilizer, feed and foodstuffs. Avoid cross-contamination with other pesticides. Spillage or leakage should be contained and absorbed with clay granules, sawdust, or equivalent material for disposal.

PESTICIDE DISPOSAL

Pesticide wastes are toxic. Triple rinse pesticide from containers and use rinsates in the pesticide application. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Plastic or Metal: After triple rinsing (or equivalent), offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities, such as burning of plastic containers. If burned, stay out of smoke.

REGISTERED TRADEMARKS

Arsenal® is a registered trademark of American Cyanamid Company.

Asulox® is a registered trademark of Rhone-Poulenc Chemical Company.

Embark® is a registered trademark of 3M.

Fenatrol® is a registered trademark of Union Carbide Agricultural Products Co., Inc.

Garlon®, and Tordon® are registered trademarks of Dow Chemical Company.

Hyvar®, Karmex®, Krenite®, Oust®, Telar®, and Velpar® are registered trademarks of E. I. duPont de Nemours & Co., Incorporated.

Limit® and Roundup® are registered trademarks of Monsanto Agricultural Products Company.

Maintain® is a registered trademark of U.S. Borax & Chemical Corporation.

Princep® is a registered trademark of Ciba-Geigy Corporation.

Royal Slo-Gro® is a registered trademark of Uniroyal, Incorporated.

Spike® is a registered trademark of Elanco Products Company.

BANVEL® is a registered trademark of Sandoz Ltd.

BANVEL 120/55947 20 pban720a 95# July 1995

LIMITATION OF WARRANTY AND LIMITATION OF LIABILITY

Notice: Read this Limitation of Warranty and Limitation of Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Sandoz or seller. All such risks shall be assumed by buyer or user.

Sandoz warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, under normal use conditions, subject to the risks described above. Sandoz makes no other express or implied warranty of fitness or of merchantability or any other express or implied warranty.

In no event shall Sandoz or seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. The exclusive remedy of the user or buyer, and the exclusive liability of Sandoz or seller for any and all claims, losses, injuries or damages (including claims based on breach of warranty, contract, negligence, tort, strict liability or otherwise) resulting from the use or handling of this product, shall be the return of the purchase price of the product or, at the election of Sandoz or seller, the replacement of the product.

Sandoz and seller offer this product, and buyer and user accept it, subject to the foregoing limitations of warranty and limitation of liability, which may not be modified by any oral or written agreement.