Reg # 55947-24 PM-25 1 of 1

CAUTION

Caution Harmful if swallowed Avoid breathing spray mist Avoid contact with skin, eyes or clothing Wash thoroughly after handling In case of contact, wash skin with soap and water, for eyes, flush with water for 15 minutes and get medical attention

IMPORTANT

Do not apply (except as recommended) on or near desirable trees or plants or in locations where chemicals may be washed or moved in contact with their roots, as injury may occur

When applying, prevent drift of product to desirable plants particularly beans, cotton, flowers, fruit trees, grapes, legume, forage crops, ornamentals, peas, potatoes, soybeans, tomatoes and other broadleaf crops (coarse sprays are less likely to drift), as injury may occur Sunflowers and tobacco are very sensitive to Weedmaster Herbicide It requires only a small amount of chemical per acre for symptoms to appear Affected tobacco leaves will be permanently malformed Do not make aerial applications if sensitive crops are nearby

Drift Precautions

To minimize Weedmaster Herbicide spray drift into sensitive broadleaf crops, (see above) observe the following precautions

1 Make applications when there is some air movement, but wind is less than 5 mph However, if any wind is blowing in the direction of sensitive crops, do not spray

2 Do not apply Weedmaster Herbicide when the daily temperature is expected to be over 85°F

3 Use coarse sprays Use 20 gallons of water per acre Keep the spray pressure below 20 psi

4 Spray with the nozzles as close to the weeds as is practical for good coverage

Do not make mist blower applications

Do not make aerial applications in the States of Mississippi, Alabama, Arkansas and Louisiana

Do not allow the product to come in direct contact with seeds, insecticides or fungicides

Do not contaminate irrigation ditches or water used for domestic purposes To avoid injury to desirable plants, equipment used for Weedmaster Herbicide should be thoroughly cleaned before reuse to handle or apply any other agricultural chemicals

All uses of Weedmaster Herbicide are intended for a normal crop growing interval If this interval is shortened, such as in grass seed crops which may be plowed under or pasture renovation, do not follow up with the planting of sensitive crops, such as those listed above

Grass crops growing under stress can exhibit various injury symptoms which may be more pronounced if herbicides are applied Consult your local or state authorities prior to treating stressed crops or new varieties as to their potential for injury

Do not use on bent, susceptible grass pastures, Lespedeza, wild winter peas, vetch clover and alfalfa pastures as injury may occur

If grasses are grown for seed or for seed down purposes do not apply when grass is in the boot to milk stage Do not apply to newly seeded areas or after heading begins

Observe all precautions on labeling of all products used in tank mixtures

Do not graze meat animals in treated areas within 30 days of slaughter

Do not graze lactating dairy animals within 7 days of treatment

Treated grasses may be harvested for dry hay, however, do not harvest within 37 days of treatment

Do not use seed from treated grasses for feed or food purposes

A nonionic surfactant such as X-77® may be added, follow the surfactant label

DO NOT USE IN ANY MANNER NOT SPECIFIED ON THE LABEL

CONTAINER DISPOSAL

Do not reuse empty container Wash thoroughly with water and detergent Discard in a safe place Do not burn

®Trademark of KALO LABORATORIES

DIRECTIONS FOR USE

For use on Pasture and Rangeland Grasses and non cropland areas such as fencerows, roadways and around farm buildings

For best control of listed weeds, apply Weedmaster Herbicide when weeds are actively growing

For ground equipment use 10 to 40 gallons of water per acre

For waiting period between treatment and grazing or harvest of treated grasses, see IMPORTANT section

ACCEPTED
MAY 3 1979
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 876-203

Transferred to 55947-24

DIRECTIONS FOR USE

WEED	DOSAGE PER ACRE WEEDMASTER HERBICIDE	SPOT TREATMENT
Bitterweed Broomweed Buffalobur Buttercup Wild Cocklebur Kochia Lambsquarters Pigweed Redroot (Carelessweed) Ragweed Common Smartweed Pennsylvania Snapweed Sunflower	1 quart per acre (½ lb Dicamba acid plus ¾ lb 2 4 D acid equivalent)	Mix ¾ teaspoon of WEEDMASTER HERBICIDE with 1 gallon of water to treat 1 sq rod (272 sq feet)
Buckeye Burdock Common Cocklebur Common Chicory Croton Woolly Dandelion Dewberry Dock Curly Dogfennel (Cypressweed) Eupatorium Ipse (Thoroughwort) Henbane Annual Goldenrod Missouri Hemlock Knotweed Mustards Annual Pennycress Field Peppergrass Persimmon Poorjoe Sorrel Red Sheep sorrel Thistle Russian	1 quart per acre (½ lb Dicamba acid plus ¾ lb 2 4 D acid equivalent)	Mix 1½ teaspoons of WEEDMASTER HERBICIDE with 1 gallon of water to treat 1 sq rod (272 sq feet)

WEED	DOSAGE PER ACRE WEEDMASTER HERBICIDE	SPOT TREATMENT
Thistle Canada (seedling) Thistle Musk Thistle Plumless (Cart) Vetch Hairy Ironweed Yellowtop	1 quart per acre (½ lb Dicamba acid plus ¾ lb 2 4 D acid equivalent)	Mix 1½ teaspoons of WEEDMASTER HERBICIDE with 1 gallon of water to treat 1 sq rod (272 sq feet)
Clover Bur Elderberry Hedge Prickly Honeysuckle Ivy Poison Knapweed Spotted Morningglory Tall Nettle Horse Nightshade Silverleaf Rubber Brush Sage Fringed Tansy Ragwort Redvine Smartweed Perennial Waterleaf TOP GROWTH SUPPRESSION Bindweed Field Knotweeds (Buckwheat & Russian) Sowthistle (Perennial) Sprange Leafy Thistle Canada	2 quarts per acre (1½ lb Dicamba acid plus 1 5 lbs 2 4 D acid equivalent)	Mix 3 teaspoons of WEEDMASTER HERBICIDE with 1 gallon of water to treat 1 sq rod (272 sq feet)

PROSPER WITH PESTICIDES BY USING THEM SAFELY. READ THE LABEL AND FOLLOW LABEL DIRECTIONS.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF THE PESTICIDE APPLICATION

NOTICE Read "LIMIT OF WARRANTY AND LIABILITY" on the container before buying or using If terms are not acceptable, return at once unopened

RECENTLY REGISTERED ON APRIL 2, 1979 FOR DISTRIBUTION AND USE IN ALL CONTINENTAL UNITED STATES

VELSICOL CHEMICAL CORPORATION
341 East Ohio Street, Chicago, Illinois 60611

07269-408AD

USE DIRECTIONS

VELSICOL®

Weedmaster®

HERBICIDE

Contains 1 lb Dicamba and 3 lbs 2,4-D Per Gallon

ACTIVE INGREDIENTS	
Dimethylamine salt of Dicamba (3 6-dichloro-o-anisic acid)*	12 37%
Dimethylamine salts of related acids	1 53%
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid**	37 17%
INERT INGREDIENTS:	48 93%
TOTAL	100 00%

*Equivalent to 10 04% 3 6-dichloro-o-anisic acid
**Equivalent to 30 11% 2 4-dichlorophenoxyacetic acid

EPA REG NO 876-203
EPA EST NO 876-TX-2

FOR USE ON PASTURE AND RANGELAND GRASSES AND NONCROPLAND AREAS SUCH AS FENCEROWS ROADWAYS AND AROUND FARM BUILDINGS

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

See container label for Warranty and Limited Liability statements and additional precautionary statements

BEST AVAILABLE COPY