


Reg # 55947-24

PM-25

1 7 2

VELSICOL



Weedr

HERBICIDE

**KEEP OUT OF THE
REACH OF CHILDREN**

CAUTION

See attached label booklet for application directions and additional precautionary statements.

E.P.A. REG. NO. 876-203AA
E.P.A. EST. NO. 876-TX-2

ROLL OR SHAKE DRUM BEFORE USING

NET CONTENTS:

ACTIVE INGREDIENT
Dimethylamine salt of
(3,6-dichloro-p-ent)
Dimethylamine salts
Dimethylamine salt of
phenoxyacetic acid
INERT INGREDIENT
TOTAL

Contains 1 lb. Dicamba
* Equivalent to 10.3%
** Equivalent to 29.6%
by AOAC isomer sp

This ingredient statement is on the supplemental labeling dicamba have not been for change in the chemical met

SICOL®

Weedmaster®

Herbicide

ACTIVE INGREDIENTS:

Dimethylamine salt of Dicamba (3,6-dichloro-g-anisic acid)*	12.0%
Dimethylamine salts of related acids	2.1%
Dimethylamine salt of 2,4-dichloro- phenoxyacetic acid**	35.0%
INERT INGREDIENTS	50.9%
TOTAL	100.0%

Contains 1 lb. Dicamba and 2.8** lbs 2,4-D per gallon

* Equivalent to 10.3% 3,6-dichloro-g-anisic acid

** Equivalent to 29.6% 2,4-dichlorophenoxyacetic acid
by AOAC isomer specific method 6D01-5.

This ingredient statement supersedes the ingredient statement appearing on the supplemental labeling booklet(s). Weedmaster® content of 2,4-D and dicamba have not been lowered or altered. This change only reflects a change in the chemical method of analysis.

CHEMICAL CORPORATION 341 East Ohio St., Chicago, Ill. 60611

CAUTION

Caution: Harmful if swallowed. breathing spray mist. Avoid contact with eyes or clothing. Wash thoroughly after handling. In case of contact, wash skin with soap and water; for eyes, flush with water for 15 minutes and get medical attention.

IMPORTANT

Do not apply (except as recommended) on desirable trees or plants or in locations where chemicals may be washed or moved in contact with their roots; as injury may occur.

When applying, prevent drift of product to desirable plants particularly beans, cotton, flowers, fruit, grapes, legume forage crops, ornamentals, potatoes, soybeans, tomatoes and other broadleaf crops (coarse sprays are less likely to drift); as injury may occur. Sunflowers and tobacco are very sensitive to Weedmaster Herbicide. It requires only a small amount of chemical per acre for symptoms to appear. Affected tobacco leaves will be permanently malformed. Do not make aerial applications if sensitive crops are nearby.

Drift Precautions:

To minimize Weedmaster Herbicide spray drift on sensitive broadleaf crops (see above) observe the following precautions:

1. Make applications when there is no wind movement, but wind is less than 5 mph. If any wind is blowing in the direction of sensitive crops, do not spray.
2. Do not apply Weedmaster Herbicide when the daily temperature is expected to be over 90°F.
3. Use coarse sprays. Use 20 gallons of water per acre. Keep the spray pressure below 20 psi.
4. Spray with the nozzles as close to the weeds as practical for good coverage.

Do not make mist blower applications.

Do not make aerial applications in the States of Mississippi, Alabama, Arkansas and Louisiana. Do not allow this product to come in direct contact with seeds, insecticides or fungicides.

Do not contaminate irrigation ditches or water used for domestic purposes. To avoid injury to desirable plants