

Reg # 55947-24

PM-25

172

DIRECTIONS FOR USE

WEED	DOSAGE PER ACRE WEEDMASTER HERBICIDE	SPOT TREATMENT
Blackwood Bramble Butcherbush Butternut, Wild Cockspur F. 1-3 Lambquarters Pigweed, Redroot (Carnegewood) Ragweed, Common Smartweed, Pennsylvanian Smartweed Sorrel	1 pint per acre	Mix 1/4 teaspoon of WEEDMASTER HERBICIDE with 1 gallon of water to treat 1 sq. rod (272 sq. feet)
Bedstraw Bindweed, Common Chickweed, Common Chicory Clover, Woolly Dandelion Dewberry Dock, Curly Dogfennel (Dyspnoea) Eupatorium, late (T. verticillatum) Fleabane, Annual Goldenrod, Missouri Hound Knotweed Mustard, Annual Pennywort, Field Pepperweed Persimmon Puncture Sorrel, Red (Sheep sorrel) Thistle, Bull	1 quart per acre	Mix 1 1/2 teaspoons of WEEDMASTER HERBICIDE with 1 gallon of water to treat 1 sq. rod (272 sq. feet)

WEED	DOSAGE PER ACRE WEEDMASTER HERBICIDE	SPOT TREATMENT
Thistle, Canada (seedling) Thistle, Musk Thistle, Plumose (curl) Vetch, Hairy Yarrowweed Yellowtop	1 quart per acre	Mix 1 1/2 teaspoons of WEEDMASTER HERBICIDE with 1 gallon of water to treat 1 sq. rod (272 sq. feet)
Clever, Bar Elderberry Hedge, Privet Honeysuckle Ivy, Poison Knotweed, Spotted Morningglory, Tall Nettle, Horse Nightshade, Silverleaf Rabbit Brush Sage, Fringed Tansy Ragwort Rudbeckia Smartweed, Perennial Valerian	2 quarts per acre	Mix 3 teaspoons of WEEDMASTER HERBICIDE with 1 gallon of water to treat 1 sq. rod (272 sq. feet)
TOP GROWTH SUPPRESSION Bindweed, Field Knotweed (Black & Russian) Sawtooth (Pernovis) Spurge, Lady Thistle, Canada		

THIS LABELING MUST BE IN THE
 SION OF THE USER AT THE TIM
 PESTICIDE APPLICATION.

NOTICE: Read "LIMIT OF WARRANTY
 LIABILITY" on the container before
 using. If terms are not acceptable, ret
 unopened.

VELSICOL CHEMICAL CORP

341 East Ohio Street, Chicago, Ill.

07269-408AE

CAUTION

Caution: Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling. In case of contact, wash skin with soap and water; for eyes, flush with water for 15 minutes and get medical attention.

IMPORTANT

Do not apply (except as recommended) on or near desirable trees or plants or in locations where chemicals may be washed or moved in contact with their roots; as injury may occur.

When applying, prevent drift of product to desirable plants particularly beans, cotton, flowers, fruit trees, grapes, legume, forage crops, ornamentals, peas, potatoes, soybeans, tomatoes and other broadleaf crops (coarse sprays are less likely to drift); as injury may occur. Sunflowers and tobacco are very sensitive to Weedmaster Herbicide. It requires only a small amount of chemical per acre for symptoms to appear. Affected tobacco leaves will be permanently malformed. Do not make aerial applications if sensitive crops are nearby.

Drift Precautions

To minimize Weedmaster Herbicide spray drift into sensitive broadleaf crops; (see above) observe the following precautions:

1. Make applications when there is some air movement, but wind is less than 5 mph. However, if any wind is blowing in the direction of sensitive crops, do not spray.

2. Do not apply Weedmaster Herbicide when the daily temperature is expected to be over 85°F.

3. Use coarse sprays. Use 20 gallons of water per acre. Keep the spray pressure below 20 psi.

4. Spray with the nozzles as close to the weeds as is practical for good coverage.

Do not make mist blower applications.

Do not make aerial applications in the States of Mississippi, Alabama, Arkansas and Louisiana.

Do not allow this product to come in direct contact with seeds, insecticides or fungicides.

Do not contaminate irrigation ditches or water used for domestic purposes. To avoid injury to desirable plants, equipment used for Weedmaster Herbicide should be thoroughly cleaned before reuse to handle or apply any other agricultural chemicals.

All uses of Weedmaster Herbicide are intended for a normal crop growing interval. If this interval is shortened, such as in grass seed crops which may be plowed under or pasture renovation, do not follow up with the planting of sensitive crops, such as those listed above.

Grass crops growing under stress can exhibit various injury symptoms which may be more pronounced if herbicides are applied. Consult your local or state authorities prior to treating stressed crops or new varieties as to their potential for injury.

Do not use on bent, susceptible grass pastures, Leespedeza, wild winter peas, vetch, clover and alfalfa pastures as injury may occur.

If grasses are grown for seed or for seed-do not apply when grass is in the boot to mil apply to newly seeded areas or after heading. Observe all precautions on labeling of all p tank mixtures.

Do not graze meat animals in treated areas v slaughter.

Do not graze lactating dairy animals within ment.

Treated grasses may be harvested for dry h not har ist within 37 days of treatment.

Do not use seed from treated grasses k purposes.

A nonionic surfactant such as X-77® may b the surfactant label.

DO NOT USE IN ANY MANNER NOT SPEC LABEL

*Trademark of KALO LABORATORIES