

Reg # 55947-24

PM-25

(g-22) (a) (b) (c) (d) (e) (f) (g) (h) (i) (j) (k) (l) (m) (n) (o) (p) (q) (r) (s) (t) (u) (v) (w) (x) (y) (z) (aa) (ab) (ac) (ad) (ae) (af) (ag) (ah) (ai) (aj) (ak) (al) (am) (an) (ao) (ap) (aq) (ar) (as) (at) (au) (av) (aw) (ax) (ay) (az) (ba) (bb) (bc) (bd) (be) (bf) (bg) (bh) (bi) (bj) (bk) (bl) (bm) (bn) (bo) (bp) (bq) (br) (bs) (bt) (bu) (bv) (bw) (bx) (by) (bz) (ca) (cb) (cc) (cd) (ce) (cf) (cg) (ch) (ci) (cj) (ck) (cl) (cm) (cn) (co) (cp) (cq) (cr) (cs) (ct) (cu) (cv) (cw) (cx) (cy) (cz) (da) (db) (dc) (dd) (de) (df) (dg) (dh) (di) (dj) (dk) (dl) (dm) (dn) (do) (dp) (dq) (dr) (ds) (dt) (du) (dv) (dw) (dx) (dy) (dz) (ea) (eb) (ec) (ed) (ee) (ef) (eg) (eh) (ei) (ej) (ek) (el) (em) (en) (eo) (ep) (eq) (er) (es) (et) (eu) (ev) (ew) (ex) (ey) (ez) (fa) (fb) (fc) (fd) (fe) (ff) (fg) (fh) (fi) (fj) (fk) (fl) (fm) (fn) (fo) (fp) (fq) (fr) (fs) (ft) (fu) (fv) (fw) (fx) (fy) (fz) (ga) (gb) (gc) (gd) (ge) (gf) (gg) (gh) (gi) (gj) (gk) (gl) (gm) (gn) (go) (gp) (gq) (gr) (gs) (gt) (gu) (gv) (gw) (gx) (gy) (gz) (ha) (hb) (hc) (hd) (he) (hf) (hg) (hh) (hi) (hj) (hk) (hl) (hm) (hn) (ho) (hp) (hq) (hr) (hs) (ht) (hu) (hv) (hw) (hx) (hy) (hz) (ia) (ib) (ic) (id) (ie) (if) (ig) (ih) (ii) (ij) (ik) (il) (im) (in) (io) (ip) (iq) (ir) (is) (it) (iu) (iv) (iw) (ix) (iy) (iz) (ja) (jb) (jc) (jd) (je) (jf) (jg) (jh) (ji) (jj) (jk) (jl) (jm) (jn) (jo) (jp) (jq) (jr) (js) (jt) (ju) (jv) (jw) (jx) (jy) (jz) (ka) (kb) (kc) (kd) (ke) (kf) (kg) (kh) (ki) (kj) (kk) (kl) (km) (kn) (ko) (kp) (kq) (kr) (ks) (kt) (ku) (kv) (kw) (kx) (ky) (kz) (la) (lb) (lc) (ld) (le) (lf) (lg) (lh) (li) (lj) (lk) (ll) (lm) (ln) (lo) (lp) (lq) (lr) (ls) (lt) (lu) (lv) (lw) (lx) (ly) (lz) (ma) (mb) (mc) (md) (me) (mf) (mg) (mh) (mi) (mj) (mk) (ml) (mm) (mn) (mo) (mp) (mq) (mr) (ms) (mt) (mu) (mv) (mw) (mx) (my) (mz) (na) (nb) (nc) (nd) (ne) (nf) (ng) (nh) (ni) (nj) (nk) (nl) (nm) (nn) (no) (np) (nq) (nr) (ns) (nt) (nu) (nv) (nw) (nx) (ny) (nz) (oa) (ob) (oc) (od) (oe) (of) (og) (oh) (oi) (oj) (ok) (ol) (om) (on) (oo) (op) (oq) (or) (os) (ot) (ou) (ov) (ow) (ox) (oy) (oz) (pa) (pb) (pc) (pd) (pe) (pf) (pg) (ph) (pi) (pj) (pk) (pl) (pm) (pn) (po) (pp) (pq) (pr) (ps) (pt) (pu) (pv) (pw) (px) (py) (pz) (qa) (qb) (qc) (qd) (qe) (qf) (qg) (qh) (qi) (qj) (qk) (ql) (qm) (qn) (qo) (qp) (qq) (qr) (qs) (qt) (qu) (qv) (qw) (qx) (qy) (qz) (ra) (rb) (rc) (rd) (re) (rf) (rg) (rh) (ri) (rj) (rk) (rl) (rm) (rn) (ro) (rp) (rq) (rr) (rs) (rt) (ru) (rv) (rw) (rx) (ry) (rz) (sa) (sb) (sc) (sd) (se) (sf) (sg) (sh) (si) (sj) (sk) (sl) (sm) (sn) (so) (sp) (sq) (sr) (ss) (st) (su) (sv) (sw) (sx) (sy) (sz) (ta) (tb) (tc) (td) (te) (tf) (tg) (th) (ti) (tj) (tk) (tl) (tm) (tn) (to) (tp) (tq) (tr) (ts) (tu) (tv) (tw) (tx) (ty) (tz) (ua) (ub) (uc) (ud) (ue) (uf) (ug) (uh) (ui) (uj) (uk) (ul) (um) (un) (uo) (up) (uq) (ur) (us) (ut) (uu) (uv) (uw) (ux) (uy) (uz) (va) (vb) (vc) (vd) (ve) (vf) (vg) (vh) (vi) (vj) (vk) (vl) (vm) (vn) (vo) (vp) (vq) (vr) (vs) (vt) (vu) (vv) (vw) (vx) (vy) (vz) (wa) (wb) (wc) (wd) (we) (wf) (wg) (wh) (wi) (wj) (wk) (wl) (wm) (wn) (wo) (wp) (wq) (wr) (ws) (wt) (wu) (wv) (ww) (wx) (wy) (wz) (xa) (xb) (xc) (xd) (xe) (xf) (xg) (xh) (xi) (xj) (xk) (xl) (xm) (xn) (xo) (xp) (xq) (xr) (xs) (xt) (xu) (xv) (xw) (xx) (xy) (xz) (ya) (yb) (yc) (yd) (ye) (yf) (yg) (yh) (yi) (yj) (yk) (yl) (ym) (yn) (yo) (yp) (yq) (yr) (ys) (yt) (yu) (yv) (yw) (yx) (yy) (yz) (za) (zb) (zc) (zd) (ze) (zf) (zg) (zh) (zi) (zj) (zk) (zl) (zm) (zn) (zo) (zp) (zq) (zr) (zs) (zt) (zu) (zv) (zw) (zx) (zy) (zz)

(Container Label)

**WEEDMASTER® Herbicide**

**Active Ingredients:**

Dimethylamine salt of dicamba

(3,6-dichloro-p-anisic acid)\* ..... 12.4%

Dimethylamine salt of 2,4-dichloro-

phenoxyacetic acid\*\* ..... 35.7%

**Inert Ingredients:** ..... 51.9%

TOTAL.... 100.0%

\*This product contains 10.3% 3,6-dichloro-p-anisic acid (dicamba) or 1 pound per gallon (120 g/L).

\*\*This product contains 29.6% 2,4-dichlorophenoxyacetic acid (2,4-D) or 2.87 lbs. per gallon (344 g/L).

\*\*Isomer specific by AOAC method 6.D01-5

**SHAKE WELL BEFORE USING**

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See side panel for additional precautionary statements.....


EPA Reg. No. 55947-24

Net Contents: \_\_\_\_\_

EPA Est. No. 55947-TX-1..

Sandoz Crop Protection Corporation

Des Plaines, IL 60018



## **PRECAUTIONARY STATEMENTS**

### **HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

#### **CAUTION**

Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. In case of contact, wash skin with soap and water; for eyes, flush with water for 15 minutes and get medical attention.

### **ENVIRONMENTAL HAZARDS**

Keep out of lakes, streams or ponds. This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters. Apply this product only as directed on label.

## **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Refer to the DIRECTIONS FOR USE booklet attached to this container for proper use directions and additional precautionary statements.

## **STORAGE AND DISPOSAL**

### **PROHIBITIONS**

Do not contaminate water, food or feed by storage or disposal.

### **STORAGE**

Store in original container in a well-ventilated area separately from fertilizer, feed and foodstuffs. Avoid cross-contamination with other pesticides. Spillage or leakage should be contained and absorbed with clay granules, sawdust, or equivalent material for disposal.

### **PESTICIDE DISPOSAL**

Triple rinse pesticide from containers and use rinsates in the pesticide application. Wastes which cannot be used according to label instructions may be disposed of on site or at an approved waste disposal facility.

### **CONTAINER DISPOSAL**

Plastic or Metal: After triple rinsing (or equivalent), offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities, such as burning of plastic containers. If burned, stay out of smoke.

4915

## **LIMITED WARRANTY AND LIABILITY**

**NOTICE:** Read this Limited Warranty and Liability before buying or using this product. If the terms are not acceptable, return it at once, unopened.

It is critical that this product be used and mixed only as specified on the label. The laws of a State may make some or all of this paragraph inapplicable or may give you rights in addition to your rights hereunder. Except to the extent prohibited by applicable law, the exclusive remedy of the user or buyer and the limit of liability of this Company or any other Seller for any and all losses, personal injuries or damages resulting from the use of this product shall be the purchase price paid by the user or buyer for the quantity of product involved. Except to the extent prohibited by State Law, there is no warranty, and this Company and other Sellers disclaim all liability for losses, personal injury or damages: (i) arising from any use of this product in a manner or for a purpose not recommended in its label directions or from mixing this product before use with any substance except as recommended by the product's label; (ii) arising from handling or storage in violation of label instructions; (iii) for all indirect, special or consequential damages; and (iv) when not reported to this Company within one year of discovery. **THERE ARE NO IMPLIED WARRANTIES AND NO WARRANTIES OF MERCHANTABILITY OR FITNESS.**

Proposed Label

1-14-95

0724800 40 AC  
SANDOX  
CROP PROTECTION  
CORPORATION  
001 000000 0000



PROPERLY WITH  
PESTICIDES BY  
USING THEM IN  
PROPERLY READ  
DIRECTIONS  
AND FOLLOW LABEL



NOTICE: READ LIMITED  
WARRANTY AND LIABILITY  
ON THE CONTAINER BEFORE  
BUYING OR USING. RETURN  
ARE NOT ACCEPTABLE. RETURN  
AT ONCE UNOPENED.

THIS LABELING MUST BE IN THE  
POSSESSION OF THE USER AT  
THE TIME OF THE PESTICIDE  
APPLICATION.

HERBICIDE

**Weedmaster**

**Weedmaster**

HERBICIDE

DIRECTIONS FOR USE

IF YOU ARE NOT A FARMER,  
LAW TO USE THIS PRODUCT IN A  
GARDEN OR HOME PLANT  
WITH THE LABELING.


THIS PRODUCT MUST BE IN  
THE POSSESSION OF THE  
USER AT THE TIME OF  
THE PESTICIDE APPLICATION

E.P.A. Reg. 03-56317-2A

Weedmaster® is a Registered  
Trademark of Sandoz Ltd

©SANDOZ CROP PROTECTION CORP 1988

**SANDOZCROP PROTECTION**



ACC  
Under the F  
Fungicide, c  
as awarded  
registered u  
EPA Reg. N

TAB

GENERAL PRECAUT  
MIXING AND APPL  
PROCEDURE FOR C  
PASTURE RANGELAN  
BETWEEN CROPPING  
FOR BROADLEAF W  
CROPLAND ROTATE  
SUGARCANE  
REGISTERED TRADE

**BEFORE USING WEEDMASTER® HERBICIDE  
READ AND FOLLOW THE PRECAUTIONS  
APPEARING ON THE CONTAINER**

**IMPORTANT**

The following directions apply to all uses of WEEDMASTER Herbicide.

Do not contaminate irrigation ditches or water used for domestic purposes.

**SENSITIVE CROP PRECAUTIONS:** WEEDMASTER Herbicide may cause injury to desirable trees and plants, particularly beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes and other broadleaf plants when contacting their roots, stems or foliage. These plants are most sensitive to WEEDMASTER Herbicide during their development or growing stage. **FOLLOW THE PRECAUTIONS LISTED BELOW WHEN USING WEEDMASTER HERBICIDE.**

- Do not treat areas where either possible downward movement into the soil or surface washing may cause contact of WEEDMASTER Herbicide with the roots of desirable plants such as trees and shrubs.
- Avoid making applications when spray particles may be carried by air currents to areas where sensitive crops and plants are growing. Do not spray near sensitive plants if wind is gusty or in excess of 5 mph and moving in the direction of nearby sensitive crops. However, always make applications when there is some air movement to determine the direction and distance of possible spray

2

drift. Leave an adequate buffer zone between area to be treated and sensitive plants. Coarse sprays are less likely to drift out of the target area than fine sprays. Agriculturally-approved drift-reducing additives may be used.

- Do not use aerial equipment to apply WEEDMASTER Herbicide when sensitive crops and plants are growing in the vicinity of area to be treated.
- To avoid injury to desirable plants, equipment used to apply WEEDMASTER Herbicide should be thoroughly cleaned (see PROCEDURE FOR CLEANING SPRAY EQUIPMENT on page 4) before reusing to apply any other chemicals.

Crops growing under stress conditions such as drought, poor fertility, or foliar damage due to hail, wind or insects, can exhibit various injury symptoms that may be more pronounced if herbicides are applied.

Consult your local or state authorities for possible application restrictions and advice concerning these and other special local use situations.

**Do not apply this product through any type of irrigation system.**

**MIXING AND APPLICATION**

WEEDMASTER Herbicide is a water-soluble formulation that may be applied using water.

WEEDMASTER Herbicide may be applied broadcast using either ground or aerial equipment. When using ground equipment, apply 5 to 40 gallons of di-

**MIXING AND APPLICATION (Cont'd)**

luted spray per treated acre. Volume of spray applied will height, density, and type of weeds or brush being treated or mow being used. When using aerial equipment, apply 3 to 10 ed spray per treated acre.

WEEDMASTER Herbicide may be applied to individual clump (SPOT TREATMENT) of undesirable vegetation using hand-ap of application equipment. Apply diluted sprays to allow com to runoff or foliage and stems. For knapsack or other small c utilize the table below to calculate material needed. (The table on the assumption that the spot treatment rate equates to 60 on the broadcast basis).

**KNAPSACK SPRAYER DILUTION INSTRU**

Sprayer Capacity (Gallons of Water)	Wt. of herbicide
1 gallon	
2 1/2 gallons	
3 gallons	
5 gallons	

The addition of a surfactant can help improve control. Add volume. For example, 5 gallons (40 pts./640 oz.) of herbicide require .2 pt. (3.2 oz.) of surfactant.

\*1 pint = 16 oz. = 2 cups and 1 cup (liquid) = 8 table

Select nozzles designed to produce minimal amounts of fine Spray with nozzles as close to the weeds as is practical coverage.

WEEDMASTER Herbicide should not be applied during peno or when wind is in excess of 15 mph as uneven spray cover

Avoid disturbing (e.g. cultivating or mowing) treated areas to following application.

5

## PROCEDURE FOR CLEANING SPRAY EQUIPMENT

The steps listed below are suggested for thorough cleaning of spray equipment following applications of WEEDMASTER He bicide.

- 1) Hose down thoroughly the inside as well as outside surfaces of equipment while filling the spray tank half full of water. Flush by operating sprayer until the system is purged of the rinse water.
- 2) Fill tank with water while adding 1 quart of household ammonia for every 25 gallons of water. Operate the pump to circulate the ammonia solution through the sprayer system for 15 to 20 minutes and discharge a small amount of the ammonia solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
- 3) Flush the solution out of the spray tank through the boom.
- 4) Remove the nozzles and screens and flush the system with two full tanks of water.

REFER TO THE CONTAINER LABEL FOR INSTRUCTIONS CONCERNING DISPOSAL OF WASTE AND CLEANING RINSES.

8

## PASTURE RANGELAND AND NONCROPLAND IMPORTANT

Do not use on bent, susceptible grass pastures, lespedeza, wild winter peas, vetch, clover and alfalfa pastures as, injury will occur. For pasture renovations, wait 3 weeks per quart per acre of WEEDMASTER Herbicide used before re-seeding or injury may occur.

If grasses are grown for seed or for seed-down purposes, do not apply when grass is in the boot to milk stage. Do not apply to newly seeded areas or after heading begins.

**Do not graze meat animals within 30 days of slaughter.**

**Do not graze lactating dairy animals within 7 days of treatment.**

Treated grasses may be harvested for dry hay, however, do not harvest within 37 days of treatment

**Do not use seed from treated grasses for feed or food purposes.**

Adjuvants or other spray additives (emulsifiers, surfactants, wetting agents, drift control agents, or penetrants) may be used for wetting, penetration, or drift control. Spray additives must be agriculturally approved when used in pasture applications. If spray additives are used, read and follow all use recommendations and precautions of product label.

• • • • •

7

RON K GRAPHIC SERVICES, LTD. .  
DISC San001 — File No. 147  
04-27-88 gk/jk Fut-1, Fut Bld -3, Fut Bld-6(1)  
Job No. 880551-1079 Sandoz Weertmaster Pcookie  
Galley 3

## WEEDS CONTROLLED, RATES A

Application rates and timings of WEEDMASTER Herbi When perennial weeds are reaching maturity, mowing will enhance control.

**WEED STAGE/APPLICATION**

Wood	Wood Moisture Rate			
	4.5	6.0	7.5	
Red Oak, Spotted				poor
Redwood, Red				
Chestnut	1.5"	2" bending		low
Red Oak Sp.				
Red Oak				fair
Red Oak Sp.		poor		
Red Oak Sp.		poor		poor
Chickadee, Common		rotting	1.5"	poor
Cherry				mod
Green Oak			poor to bending	mod
Cottonwood, Common	1.5"	4.5"	6.5"	12.5"

**\*The more you eat, the more you eat.**

\*Downloaded from [www.sagepub.com](http://www.sagepub.com) by guest on June 13, 2015

1

**WILD STAGE/APPLICATION ST**[illegible]

\*The above are estimates

\* Government rate not payable by South American cities

1

RON K. GRAPHIC SERVICES, LTD.  
DISC San001 — File No. 148  
04-27-88 gk/yk Fut.-1,Fut Bld.-3,Fut Bld.-6(1)  
Job No. 880551-1079 Sandoz Weedmaster Book  
Gallery 4

**WILD STAGE/APPLICATION STAGE**

**Woodwater Rate (pints/acre)**

\*Buy more open applications  
\*Downloaded are all paid by bank transfers only

**Windometer Rate (girths/acre)**

- May require repeat applications
- Recommended only on roads by local government only

**Woodmaster Rate (mths.)**

- \* They were never captured.
- \* "We would not be able to find human remains."

## IMPORTANT

Before using this product, read and carefully observe all application restrictions, and precautionary statements on the EPA-registered label and the WEEDMASTER Herbicide Directions For Use booklet.

## WEEDS CONTROLLED

**WEEDMASTER** Herbicide, when applied at the recommended rate, will control the **ANNUAL** and **BIENNIAL** weeds and suppress the 1 listed below:

## ANNUALS

- Buckwheat, Wild
- Cockle, Cow
- Cocklebur, Common
- Knotweed
- Kochia
- Lambquarters, Common
- Mallow, Common
- Mustards
- Nightshade, Black
- Pigweed, Redroot (*careless weed*)
- Pigweed, Rough
- P. sp. - Common

**ANNUALS (cont.)**

Ragweed, Common  
Smartweed, Pennsylv  
Sowthistle, Annual  
Sunflower  
Tansymustard  
Thistle, Russian  
Velvetleaf



### BIENNIALS

Carrot, Wild  
Ragwort, Tansy  
Thistle, Bull  
Thistle, Musk  
Thistle, Plumless  
Star Thistle, Yellow

### PERENNIALS

Bindweed, Field  
Dock, Curly  
Thistle, Canada

### RATES AND TIMINGS

Application may be made to fallow land, wheat stubble or land to be rotated to wheat. Application should be made to emerged and actively growing weeds. Use higher rate when treating dense vegetative growth. Avoid disturbing treated areas for seven days following application.

Wheat injury may occur if the interval between application and planting is less than 10 days for each pint per acre of WEEDMASTER Herbicide used. Exclude days when ground is frozen.

#### Weed Type & Stage

##### Annual

small, actively growing  
(less than 4 inches)  
established weed growth  
(greater than 4 inches)

#### Broadleaf Rate per Treated Acre

1 - 1½ pints

1½ - 3 pints

14

##### Biennial

rosette diameter  
3 inches or less  
3 inches or more  
greater than 4 inches, flowering,  
balled or flowering

1½ - 2 pints

2 - 4 pints


4 pints

##### Perennial

suppression or top  
growth control  
seasonal control

2 - 4 pints

4 - 8 pints



15

Add 0.5% v/v of an agriculturally-approved surfactant bicide when used alone or in a tank mix. The add enhance spray coverage and the herbicide's penetrat

Retreatments may be made as needed; however, do not pints of WEEDMASTER Herbicide per treated acre.

### TANK MIX TREATMENTS

WEEDMASTER Herbicide may be tank mixed with one herbicides for control of grasses or additional broad follow the label of each tank mix product used for pre directions for use, rates and timings, weeds controller restrictions. Add 0.5% v/v of an agriculturally-approved mases.

HERBICIDE	TREAT
atrazine	1/2
chlorosulfuron (Glean®)	0
cyanazine	1/2
glyphosate (Roundup®)	1/2
metribuzin	1/2
paraquat	1/2

16

### SUGARCANE IMPORTANT

Consult your local or state authorities for possible especially concerning aerial applications and advice c use situations.

### WEEDS CONTROLLED

WEEDMASTER Herbicide, when applied at recomme the ANNUAL and PERENNIAL broadleaf weeds can cone, such as:

ANNUALS	ANNUALS (cont.)	E
Burclover, California	Pigweed, Prostrate	A
Buttercup	Pigweed, Redroot	B
(Ranunculus Spp.)	(Carelessweed)	C
Chickweed, Common	Pigweed, Smooth	D
Clavels, Annual	Purslane, Common	D
Cocklebur, Common	Ragweed, Common	D
Eveningprimrose, Cutleaf	Shepherdspurse	G
Flabone, Annual	Sida, Prickly	G
Geranium, Carolina	(leafweed)	G
Henbit	Sowthistle, Annual	O
Lettuce, Prickly	Spurge, Prostrate	S
Pepperweed, Virginia		

17

RON K GRAPHIC SERVICES, LTD

DISC Son001 — File No. 151

04 27-88 gk/jk Fut -1, Fut Bld -3, Fut Bld -6(1);

Job No 880551-1079 Sandoz Weedmaster Booklet

Griffey 6

## APPLICATION DIRECTIONS

DO NOT USE AERIAL APPLICATION EQUIPMENT IF SENSITIVE CROPS ARE GROWING IN THE VICINITY OF THE AREA TO BE TREATED. When band treatments are desired, apply an equivalent rate and volume per acre by using the formula below:

Band width inches	×	Broadcast RATE per treated acre	=	Band RATE per treated acre
Row width inches				
Band width inches	×	Broadcast VOLUME per treated acre	=	Band VOLUME per treated acre
Row width inches				

Application of WEEDMASTER Herbicide can be made any time after the weeds have emerged and are actively growing but prior to the close-in stage of sugarcane. When possible, direct the spray beneath the sugarcane canopy in order to minimize the likelihood of crop injury. The use of directed sprays will also aid in maximizing spray coverage of weed foliage. Application rates and timing are given below. Use the higher level of listed rate ranges when treating dense vegetative growth.

- For control of listed ANNUAL broadleaf weeds, apply 1 quart of WEEDMASTER Herbicide per treated acre.
- For suppression of listed PERENNIALS, apply 1-2 quarts of WEEDMASTER Herbicide per treated acre.

18

- For control of listed PERENNIAL broadleaf weeds, apply 2-3 quarts of WEEDMASTER Herbicide per treated acre.

Retreatments may be made as needed, however, do not exceed 8 quarts of WEEDMASTER Herbicide per treated acre during a growing season.

## TANK MIX TREATMENTS

WEEDMASTER Herbicide may be tank mixed with one or more of the following herbicides for control of grasses or additional broadleaf weeds. Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, weeds controlled, geographic and other restrictions.

HERBICIDE	RATES PER TREATED ACRE (lbs. a.i.)
ametryn (Evik <sup>®</sup> )	3 1/2 to 8
asulam (Asulam <sup>®</sup> )	2 to 3 1/2
atrazine	3 1/2 to 4
dalapon (Dalapon <sup>®</sup> )	3 1/2 to 8 1/2
lenac (Fenatra <sup>®</sup> Sugarcane)	2.2 to 6.0
metribuzin (Sencor <sup>®</sup> , Leson <sup>®</sup> )	0.75 to 3.0
terbacil (Sinbar <sup>®</sup> )	1.6 to 3.2



RON K. GRAPHIC SERVICES, LTD.  
DISC San001 — File No. 151  
04-28-88 gk/jk Fut.-1, Fut. Bld.-3, Fut. Bld.-6(1)  
Job No. 880551-1079 Sandoz Weedmaster Booklet  
Galley 7

RON K. GRAPHIC SERVICES, LTD.  
DISC San001 — File No. 152  
04-28-88 gk/jk Fut.-1, Fut. Bld.-3, Fut. Bld.-6(1)

## REGISTERED TRADE

Glean<sup>®</sup>, Lenac<sup>®</sup>, Karmex<sup>®</sup>, and Sinbar<sup>®</sup> are r  
DuPont de Nemours & Co., Inc.

Roundup<sup>®</sup> is a registered trademark of Monsanto  
pany.

Sencor<sup>®</sup> is a registered trademark of Mobay C  
Evik<sup>®</sup> and Igran<sup>®</sup> are registered trademarks of C  
ricultural Division.

Asulam<sup>®</sup> is a registered trademark of Rhone-Pou

Dalapon<sup>®</sup> is a registered trademark of SDS Bio

Fenatra<sup>®</sup> Sugarcane is a registered trademark a  
Products Co., Inc.

20

RON K. GRAPHIC SERVICES, LTD.  
DISC San001 — File No. 153  
04-28-88 gk/jk Fut.-1, Fut. Bld.-3, Fut. Bld.-6(1)  
Job No. 880551-1079 Sandoz Weedmaster  
Galley 9

# TANK MIX TREATMENTS

For control of grasses or additional broadleaf weeds, WEEDMASTER herbicide may be tank mixed with the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled, geographic and other restrictions.

## BROADCAST RATE PER TREATED ACRE:

~~Apply 1/2 - 1 pint WEEDMASTER herbicide with:~~

Herbicide	amount of product*	lb. a.i.
2,4-D Amine or Ester	1/2	1/4
MCPA	1/2 - 3/4 pint	1/4 - 3/8
bromoxynil	Consult Product Labels for Rate	
metribuzin	Recommendations	
(Sencor®)	"	"
sulfonyleureas	"	"
(such as: Ally®, Express®, Finesse®, Glean®, Harmony® Extra)	"	"
diuron (Karmex® 80WP)	"	"

\*based on 4 pounds per gallon formulations of MCPA and 2,4-D.

In spring seeded wheat, 1/2 pint of WEEDMASTER may be tankmixed with the 1/3 pint rate of 2,4-D only. Do not tankmix with 2,4-D if the 1.0 pint rate of WEEDMASTER is used.

**SMALL GRAINS  
(NOT UNDERSEED TO LEGUMES)  
IMPORTANT**

If small grains are grown for pasture only, refer to the PASTURE, RANGELAND and NONCROPLAND section.

Do not graze or harvest for livestock feed prior to crop maturity.

**WEEDS CONTROLLED**

WEEDMASTER herbicide, when applied at recommended rates, will control ANNUAL and BIENNIAL broadleaf weeds commonly found in small grains, such as:

buckwheat, wild	pennycress, field
chickweed, common	pigweed, redroot (carelessweed)
*cockle, cow	pigweed, rough
*cocklebur, common	*purslane, common
eveningprimrose, common	ragweed, common
*henbit	shepherdspurse
*knotweed	*smartweed, Pennsylvania
kochia	*sowthistle, annual
lambsquarters, common	sunflower
*mallow, common	tansymustard
mustards	thistle, Russian
*nightshade, black	*velvetleaf

\*Use higher rate or tankmixes for control of noted (\*) annuals.

**RATES AND TIMINGS**

Application of WEEDMASTER herbicide may be made after wheat is fully tillered to emerged and actively growing weeds. See specific crop for timing restrictions. For best performance, make application when weeds are in the 2-3 leaf stage and rosettes are less than 2 inches across. Use the higher level of listed rate ranges when treating more mature weeds or dense vegetative growth.

12 August is the date the 13 of 15  
check the labeling.  
with amendment A ✓

**SMALL GRAINS** - ?  
(NOT UNDERSEED TO LEGUMES)  
**IMPORTANT**

Small grains

If small grains are grown for pasture only, refer to the PASTURE,  
RANGELAND and NONCROPLAND section.

Do not graze or harvest for livestock feed prior to crop maturity.

#### WEEDS CONTROLLED

Weedmaster herbicide, when applied at recommended rates, will control  
ANNUAL broadleaf weeds commonly found in small grains, such as:

buckwheat, wild	pigweed, rough
*cockle, cow	*purslane, common
*cocklebur, common	ragweed, common
*knotweed	*smartweed, Pennsylvania
kochia	*sowthistle, annual
lambsquarters, common	sunflower
*mallow, common	tansymustard
mustards	thistle, Russian
*nightshade, black	*velvetleaf
pigweed, redroot (carelessweed)	

What  
small  
grains

\*Use higher rate for control of noted (\*) annuals.

#### RATES AND TIMINGS

Application of BANVEL herbicide may be made before, during or after  
planting to emerged and actively growing weeds. See specific crop for  
timing restrictions. For best performance, make application when weeds  
are in the 2-3 leaf stage and rosettes are less than 2 inches across. Use  
the higher level of listed rate ranges when treating more mature weeds or  
dense vegetative growth.

SMALL GRAINS SECTION

FROM

Sept. 24, 1985 approved  
label

14 of 15  
✓

# FALL SEEDED WHEAT

WEEDMASTER HERBICIDE MUST BE APPLIED TO FALL SEEDED WHEAT PRIOR TO THE JOINTING STAGE.

BROADCAST RATE PER TREATED ACRE:  
3/4 - 1 pint

## TANK MIX TREATMENTS

For control of grasses or additional broadleaf weeds, Weedmaster herbicide may be tank mixed with the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled, geographic and other restrictions.

### BROADCAST RATE PER TREATED ACRE:

Apply 1/2 - 1 pint WEEDMASTER herbicide with:

Herbicide	amount product*	lb. a.i.
MCPA	1/2 - 3/4 pint	1/4 - 3/8
bromoxynil	1 - 1 1/2 pint	1/4 - 3/8
bromoxynil + MCPA	1 + 1/2 pint	1/4 + 1/4
metribuzin		
(Sencor® 75DF,		
Lexone® DF™)	1/3 lb.	1/4
(Sencor® 4, Lexone® 4L)	1/2 pint	1/4
(Lexone® 50WP)	1/2 lb.	1/4
chlorsulfuron		
(Glean® 75 DF)	1/6 - 1/2 wt. oz.	0.008-0.024
diuron (Karlex® 80WP)	1/2 - 2 lbs.	2/5 - 1 3/5
terbutryn (Igran® 80W)	1 1/2 - 2 3/4 lbs.	1 1/5 - 2 1/5

\*based on 4 pounds per gallon formulations of MCPA and 2 pounds per gallon formulation bromoxynil.

SMALL GRAINS SECTION

FROM Sept. 24, 1985

approved label

15 9 15  
✓

SPRING SEEDED WHEAT ✓

WEEDMASTER HERBICIDE MUST BE APPLIED BEFORE SPRING SEEDED WHEAT EXCEEDS THE 5 LEAF STAGE.

BROADCAST RATE PER TREATED ACRE:  
3/4 - 1 pint

TANK MIX TREATMENTS

For control of additional broadleaf weeds, Weedmaster herbicide may be tankmixed with the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled, geographic and other restrictions.

BROADCAST RATE PER TREATED ACRE:

Apply 1/2 - 1 pint WEEDMASTER herbicide with:

Herbicide	amount product*	lb. a.i.
MCPA	1/2 - 3/4 pint	1/4 - 3/8
bromoxynil	1 - 1 1/2 pint	1/4 - 3/8
chlorsulfuron (Glean® 75DF)	1/6 - 1/2 wt. oz.	0.008 - 0.024

\*based on 4 pounds per gallon formulations of MCPA and 2 pounds per gallon formulation of bromoxynil.

SMALL GRAINS SECTION  
FROM SEPT. 24, 1985  
approved label.

...