

MAR 15 1995

Jonathan E. Bryant, PH.D.
Sandoz Agro, Inc.
1300 E. Touhy Ave.
Des Plaines, IL 60018

Dear Dr. Bryant:

Subject: WEEDMASTER® Herbicide
EPA Registration No. 55947-24
Application Dated February 24, 1995, Request To
Add Use-Patterns for Use on Sorghum, Supplemental
Labeling and Amended Confidential Statement of
Formula, First Requested January 17, 1995

The revised confidential statement of formula submitted on EPA Form 8570-4 and dated February 24, 1995 has been reviewed and was found to be acceptable as an amendment to the subject EPA pesticide product registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended. ~~A copy of the confidential statement of formula has been~~ added to this Agency's records for this product.

The proposed amendment to add use-patterns for the use of WEEDMASTER Herbicide for use in the culture of sorghum has been reviewed and found acceptable for registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended provided that you:

- c Submit one (1) printed copy of your final printed labeling before you release the product of shipment under the subject labeling.

If this condition is not complied with, the registration will be subject to cancellation in accordance with FIFRA,

BEST COPY AVAILABLE

copy

-2-

section 6(e). Your release for shipment of the product constitutes acceptance of this condition.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

Joanne I. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

E.Wilson:Diskette ABC17: 03-15-95

BEST COPY AVAILABLE

2

WEEDMASTER HERBICIDE^(R)

SUPPLEMENTAL LABELING For Application of WEEDMASTER Herbicide in Sorghum EPA Reg. No. 55947-24

**All label restrictions and Directions
for Use of WEEDMASTER Herbicide
apply.**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Important

Before using this product, read carefully all applicable directions, restrictions and precautionary statements on the EPA-registered container label and Directions for Use Booklet attached to the WEEDMASTER Herbicide container.

**THIS LABELING MUST BE IN THE
POSSESSION OF THE USER AT THE
TIME OF APPLICATION.**

SORGHUM (Milo)

Applications of WEEDMASTER Herbicide to sorghum during periods of rapid growth may result in temporary leaning of plants or rolling of leaves. These effects are usually outgrown within 10 to 14 days. Sorghum growing under conditions of stress such as high moisture, low fertility, and abnormal temperature may be more sensitive to applications of WEEDMASTER Herbicide. Do not use WEEDMASTER Herbicide if the potential for sorghum injury is not acceptable.

Do not graze or feed treated sorghum forage or silage prior to mature grain stage. If sorghum is grown for pasture or hay, refer to the pasture use section on the WEEDMASTER Herbicide container label. Do not apply WEEDMASTER Herbicide to sorghum grown for seed production.

Make no more than one application per growing season.

WEEDS CONTROLLED

WEEDMASTER Herbicide, when applied at the recommended rate for grain sorghum, will control many actively growing ANNUAL broadleaf weeds and will reduce competition from established PERENNIAL broadleaf weeds as well as controlling their seedlings.

ANNUALS

Pigweed, Prostrate
Pigweed, Redroot
Pigweed, Smooth
Pigweed, Tumble
Pigweed, Rough
Cocklebur, Common
Sunflower, Annual
Morningglory, Tall
Morningglory, Ivyleaf
Russian, Thistle
Kochia

PERENNIALS (Suppression)

Bindweed, Field
Carolina horsenettle
Silverleaf nightshade

Consult your local or state authorities for possible application restrictions, especially concerning aerial applications and advice concerning special local use situations.

RATES AND TIMINGS

Postemergence applications of WEEDMASTER Herbicide can be made to sorghum in the 3 to 5 leaf stage (4 to 8" tall). For best performance make application when weeds are small (less than 3" tall). Use the higher level of listed rate range when treating broadleaf weeds that are more mature, under moisture stress, or with dense vegetative growth. Generally, the lower

ACCEPTED
with COMMENTS
In EPA Letter Dated

MAR 15 1996

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
55947-24

4/28/41

dosages recommended on this label will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species and under conditions where control is more difficult, the higher dosages will be needed. Do not use surfactants or oils. The use of surfactants or oils may increase the risk of crop injury.

WEEDMASTER Herbicide® is a
Registered Trademark of Sandoz Ltd.
Jan 1995

BROADCAST RATE PER TREATED ACRE:
0.5 to 1.5 pints per acre

TANK MIX TREATMENT

WEEDMASTER plus Atrazine:

For improved control of emerged, actively growing broadleaf weeds including triazine resistant species and added suppression of perennial broadleaf weeds, tank mix 0.5 to 1.5 pints of WEEDMASTER Herbicide with 0.5 to 1.25 lbs. a.i. atrazine per treated acre. For control of grasses (less than 1.5" tall), tank mix 0.5 to 1.5 pints of WEEDMASTER Herbicide with 2.0 lbs a.i. atrazine per treated acre. For best performance and minimal crop injury, make application when the sorghum is in the 3 to 5 leaf stage (4 to 8" tall) and when broadleaf weeds are small (less than 3" tall). The atrazine rate will depend upon soil texture and length of residual weed control desired. Do not apply atrazine on coarse soils or medium textured soil with less than 1.5% organic matter. Do not apply postemergence in OK, TX, NM, CO, and Western KS before sorghum is 6" tall. Follow all atrazine label restrictions and State and Federal restrictions pertaining to atrazine application.

Registrant:
Sandoz Agro, Inc.
1300 East Touhy Avenue
Des Plaines, Illinois 60018

