


BRUSH BUSTER® Herbicide

**FOR NONSELECTIVE BRUSH AND BROADLEAF WEED CONTROL ALONG
FENCE ROWS, DRAINAGE DITCH BANKS, ~~HIGHWAY RIGHTS-OF-WAY,~~
WASTELAND AND SIMILAR NONCROPLAND**



Active Ingredients:

Dimethylamine salt of dicamba
(3,6-dichloro-o-anisic acid)* 12.82%

Dimethylamine salt of 2,4-dichloro-
phenoxyacetic acid**..... 24.58%

Inert Ingredients: 62.60%

TOTAL.... 100.00%

*This product contains 10.6% 3,6-dichloro-o-anisic acid
(dicamba) or 1 pound per gallon (120 g/L).

**This product contains 20.4% 2,4-dichlorophenoxyacetic
acid (2,4-D) or 1.9 lbs. per gallon (228 g/L).

**Isomer specific by AOAC method 6.275-6.279, 13th edition

SHAKE WELL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panel for additional precautionary statements

EPA Reg. No. 55947-25

EPA Est. No. 55947-TX-1

Net Contents: _____

Sandoz Agro, Inc.

Des Plaines, IL 60018

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes, and clothing. When handling this product, wear chemical resistant gloves. In case of contact, wash skin with soap and water; for eyes, flush with water for 15 minutes and get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply this product only as directed on label.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

IMPORTANT

Do not apply (except as recommended) on or near desirable plants, or in locations where the chemical may be washed or moved into contact with their roots. Applications by ground rigs and hand dispensers should be carried out only when there is no hazard from drift of product to susceptible plants such as beans, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers and tomatoes. during both the growing and dormant periods. (Coarse sprays are less likely to drift). Do not contaminate irrigation ditches or water used for domestic purposes. Do not allow this product to come in direct contact with seeds, insecticides or fungicides. To avoid injury to desirable plants, equipment used for **BRUSH BUSTER** should be thoroughly cleaned before reuse to handle or apply any other agricultural chemicals. Do not graze meat animals within 3 days of slaughter. Treated grasses may be harvested for dry hay, however, do not harvest within 30 days of treatment. When using 1 gallon of **BRUSH BUSTER** per acre, do not graze dairy animals on treated areas within 7 days after application. When using 3 gallons of **BRUSH BUSTER** per acre, do not graze dairy animals on treated areas within 60 days after application.

BRUSH OR WEED	AMOUNTS FOR PREPARATION OF SPRAY MIX	APPLICATION DIRECTIONS
Perennial Broadleaf Weeds and Vines Canada thistle Leafy spurge Redvine Curly dock Milkweed Russian knapweed Darnel or toothwort Perennial ragweed Tansy ragwort Field bindweed Puncturevine Tawha (morningglory) and many other perennial broadleaf weeds	Add 1/2 gallon of Brush Buster Herbicide (equivalent to 1/2 lb dicamba plus 1 lb 2,4-D acid) plus 4 1/2 gallons of water to make 50 gallons of spray mix	For effective broad spectrum control of annual and perennial broadleaf weeds and vines apply Brush Buster Herbicide when weeds are actively growing. Apply at the rate of 50 gallons of spray mix per 1/2 acre (about 22 000 square feet or equivalent to a strip 5,500 feet long x 4 feet wide) If less than 50 gallons of spray mix is applied per half acre increase the concentration of Brush Buster Herbicide to maintain the same rate of active ingredients per half-acre
Annual Broadleaf Weeds Carpetweed Herb Ragweed Chickweed Knotweed Sheep sorrel Clover Lambquarters (red sorrel) Cocklebur Mustard Smartweed Dogfennel Pigweed Velvetleaf English daisy Purslane Wild buckwheat and many other annual broadleaf weeds	Add 1 quart of Brush Buster Herbicide (equivalent to 1/2 lb dicamba plus 1/2 lb 2,4-D acid) plus 4 1/2 gallons of water to make 50 gallons of spray mix	FOR SPOT TREATMENTS Mix 1 to 2 ounces of Brush Buster Herbicide with 2 to 3 gallons of water to treat 700 square feet
STEM FOLIAGE APPLICATION—HIGH WATER VOLUME		
Woody Brush Alder Hornbeam Ash Hornbeam Aspen Locust Basswood Maple Cedar Mesquite Cherry Oak Chinquapin Pecan Cucumber tree Pine Gum Poplar Guava Sassafras Dogwood Sycamore Elm (Christmasberry) and similar woody plant species	Add 1/2 gallon of Brush Buster Herbicide (equivalent to 1/2 lb dicamba plus 1 lb 2,4-D acid) plus 4 1/2 gallons of water to make 50 gallons of spray mix	Apply Brush Buster Herbicide after leaves are fully developed and until 3 weeks before frost Treat all stems and foliage to runoff with special emphasis on covering the root crown. Apply at the rate of 50-75 gallons of spray mix per 1/2 acre (about 11 000 square feet) depending on the height and density of the brush
AERIAL APPLICATION*—15 GALLONS OF SPRAY MIX/ACRE		
	Add 3 gallons of Brush Buster Herbicide (equivalent to 3 lbs dicamba plus 6 lbs 2,4-D acid) plus 12 gallons of water to make 15 gallons of spray mix	Apply Brush Buster Herbicide after leaves are fully developed and until 3 weeks before frost Apply at the rate of 15 gallons of spray mix per acre of brush
AERIAL APPLICATION*—30 GALLONS OF SPRAY MIX/ACRE		
	Add 3 gallons of Brush Buster Herbicide (equivalent to 3 lbs dicamba plus 6 lbs 2,4-D acid) to 27 gallons of water to make 30 gallons of spray mix	Apply Brush Buster Herbicide after leaves are fully developed and until 3 weeks before frost Apply at the rate of 30 gallons of spray mix per acre of brush

*Before making aerial applications consult your State or Local Authorities for possible aerial application restrictions. To reduce the hazards of spray drift when applying this product by ground or air equipment apply with a drift reducing agent such as foam spray adjuvant, inert system or with other systems designed to reduce spray drift.

BRUSH BUSTER IS A TRADEMARK OF SANDOZ CROP PROTECTION CORP
 AGRO, Inc

©SANDOZ CROP PROTECTION CORP
 Sandoz Agro, Inc.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage or disposal.

STORAGE

Store in original container in a well-ventilated area separately from fertilizer, feed and foodstuffs. Avoid cross-contamination with other pesticides. Spillage or leakage should be contained and absorbed with clay granules, sawdust, or equivalent material for disposal.

PESTICIDE DISPOSAL

Pesticide wastes are toxic. Triple rinse pesticide from containers and use rinsates in the pesticide application. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Plastic or Metal: After triple rinsing (or equivalent), offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities, such as burning of plastic containers. If burned, stay out of smoke.

WARRANTY AND CONDITIONS OF SALE

Sandoz Crop Protection Corporation warrants that the chemical composition of this product conforms to the chemical description on the label and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Sandoz makes no other warranty, express or implied, concerning the use of this product other than as indicated on the label. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given on the label.

Proposed Label

4-13-92