

PM 25

55947-46

page 1 of 16

PCLARITA.95B
CLARITY™ Herbicide

May 25, 1995

FOR WEED CONTROL IN CORN.

Active Ingredient:

Diglycolamine® salt of 3,6-dichloro-o-anisic acid*
.....56.8%

Inert Ingredients:43.2%

TOTAL100.0%



*This product contains 38.5% 3,6-dichloro-o-anisic acid or 4 pounds per gallon (480 g/L)

KEEP OUT OF REACH OF CHILDREN
CAUTION

See side panel for additional precautionary statements

EPA Reg. No. 55947-46
Net Contents: _____

EPA Est. No.
Sandoz Agro Inc.
Des Plaines, IL 60018



PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Avoid contact with skin, eyes or clothing. Harmful if swallowed. Avoid breathing spray mist. Wash thoroughly after handling. In case of contact, wash skin with soap and water; for eyes, flush with water for 15 minutes and get medical attention.

Personal Protective Equipment:

Applicators and other handlers must wear:

- o Long-sleeved shirt and long pants
- o Waterproof gloves
- o Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements:


When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- o Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- o Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- o Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS



PCLARITA.95B

May 25, 1995

Keep out of lakes, streams or ponds. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Apply this product only as directed on label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- o Coveralls
- o Waterproof gloves
- o Shoes plus socks

Do not apply this product through any type of irrigation system. Do not treat irrigation ditches or water used for crop irrigation or domestic uses. CLARITY is registered and used for broadleaf weed control in field, seed, popcorn and silage corn.

CLARITY is not registered for use in sweetcorn.

Before applying **CLARITY**, read all directions and precautions appearing on the container label and in this booklet. Failure to follow all directions and precautions may result in unsatisfactory weed control, crop injury, or illegal residues.

GENERAL INFORMATION

CLARITY is registered for the control of many annual, and control or suppression of many perennial broadleaf weeds found in corn (see **GENERAL WEED LIST**). **CLARITY** is absorbed by leaves, roots and shoots to give both contact and soil residual broadleaf weed control.

Do not apply **CLARITY** to seed corn or popcorn without first verifying with your local seed corn company (supplier) the **CLARITY** selectivity on your inbred line or variety of popcorn. This precaution will help avoid potential injury of sensitive varieties.

Direct contact of **CLARITY** with corn seed must be avoided. If corn seeds are less than 1 1/2 inches below the soil surface, delay application until corn has emerged.

Corn may be harvested or grazed for feed once the crop has reached the ensilage (milk) stage or later in maturity.

ENVIRONMENTAL CONDITIONS

Applications of **CLARITY** to corn during periods of rapid growth may result in temporary leaning. Corn will usually become erect within 3-7 days with no subsequent yield reduction. Cultivation should be delayed until after corn is growing normally to avoid breakage.

Corn growing under stress conditions such as drought, poor fertility, or foliar damage due to hail, wind or insects, can exhibit various injury symptoms that may be more pronounced if herbicides are applied.

For best performance, rainfall or irrigation should not occur for 4 hours after treatment if application is made to emerged weeds greater than 3 inches tall.

SENSITIVE CROP PRECAUTIONS

CLARITY may cause injury to desirable broadleaf plants such as tobacco and soybean.

DO NOT allow spray from application equipment to drift onto sensitive broadleaf crops.

Always select nozzles designed to produce coarse sprays without producing fine spray particles. Fine spray particles can be blown by wind onto sensitive broadleaf crops. Avoid making applications when spray particles can be carried by wind to areas where sensitive broadleaf crops or plants are growing, or when temperature inversions exist. Do not spray near sensitive plants if wind is in excess of 5 miles per hour and moving in the direction of sensitive crops.

PCLARITA.95B

May 25, 1995

Agriculturally approved drift reducing additives may be used.

Thoroughly clean spray equipment after use to avoid spray tank contamination which can result in injury when subsequent applications are made to broadleaf crops. (see PROCEDURE for CLEANING SPRAY EQUIPMENT)

Use in corn is intended for a normal growing interval between planting and harvest. No crop rotation restrictions exist if normal harvest of treated crop has occurred. If this interval is shortened, such as in catastrophic crop failure from hail, flood etc., injury may occur if a sensitive broadleaf crop is used for plantback.

BAND TREATMENTS

CLARITY may be applied as a band treatment. Use the formulas below to determine the appropriate rate and volume per treated acre.

$$\begin{array}{l} \text{Band width} \\ \text{in inches} \\ \text{Row width} \end{array} \times \begin{array}{l} \text{Broadcast} \\ \text{RATE} \\ \text{per treated acre} \end{array} = \begin{array}{l} \text{Band RATE} \\ \text{per treated acre} \end{array}$$

$$\begin{array}{l} \text{Band width} \\ \text{in inches} \\ \text{Row width} \end{array} \times \begin{array}{l} \text{Broadcast} \\ \text{VOLUME} \\ \text{per treated acre} \end{array} = \begin{array}{l} \text{Band VOLUME} \\ \text{per treated acre} \end{array}$$


COMPATIBILITY TEST

Before mixing in the spray tank, it is advisable to test compatibility by mixing all components in a small container in proportionate quantities (see following table).

Amount of Herbicide to Add to One Pint
of Spray Carrier
(Assuming Volume is 25 Gallons per Acre)

FORMULATIONS	HERBICIDE RATE PER ACRE	LEVEL TEASPOONS
Dry	1 lb	1 1/2
Liquid	1 pt	1/2

If herbicide(s) do not ball-up or form flakes, sludge, gels, oily films or layers, or other precipitates, then the tested spray mix is compatible. Usually, incompatibility in any of the above described forms will occur within 5 minutes after mixing.



PCLARITA.95B

May 25, 1995

If components are incompatible, the use of a compatibility agent is recommended. Rerun the above COMPATIBILITY TEST with a suitable compatibility agent (1/4 teaspoon is equivalent to 2 pints per 100 gallons of fluid fertilizer).


GENERAL WEED LIST

CLARITY, when applied at recommended rates, will control many ANNUAL broadleaf weeds and give growth suppression of many PERENNIAL broadleaf weeds commonly found in corn.

For best performance, make applications when weeds are emerged and actively growing.

ANNUALS

- Buckwheat, Wild Burclover, California
- Burcucumber
- Carpetweed
- Chickweed, Common
- Clovers (Annual)
- Cocklebur, Common
- Jimsonweed
- Knotweed
- Kochia
- Kochia (triazine resistant)
- Ladysthumb
- Lambsquarters, Common
- Lambsquarters
(triazine resistant)
- Mallow, Common
- Mallow, Venice
- Morningglory, Ivyleaf
- Morningglory, Tall
- Mustard, Tansy
- Mustard, Wild
- Mustard (Yellowtops)
- Nightshade, Black
- Pigweed, Prostrate
- Pigweed, Redroot
(Carlessweed)
- Pigweed, Rough
- Pigweed, Smooth
- Pigweed (triazine resistant)
- Pigweed, Tumble
- Puncturevine
- Purslane, Common
- Ragweed, Common
- Ragweed, Giant (Buffaloweed)
- Ragweed Lance-Leaf
- Sicklepod



PCLARITA.95B


May 25, 1995

Sida, Prickly (Teaweed)
 Smartweed, Green
 Smartweed, Pennsylvania
 Spikeweed, Common
 Spanish needles
 Spurge, Prostrate
 Sunflower, Common (Wild)
 Sunflower, volunteer
 Thistle, Russian
 Velvetleaf
 Waterhemp

PERENNIALS

Alfalfa
 Artichoke, Jerusalem
 Bindweed, Field
 Clover, Hop
 Dandelion, Common
 Dock Broadleaf (Bitterdock)
 Dock, Curly
 Dogbane, Hemp
 Horsenettle, Carolina
 Mare's Tail (Horseweed)
 Milkweed, Climbing
 Milkweed, Common
 Milkweed, Honeyvine
 Nightshade, Silverleaf
 (White Horsenettle)
 Pokeweed
 Smartweed, Swamp
 Sowthistle
 Thistle, Canada
 Vetch

MIXING AND APPLICATION

CLARITY is a water-soluble formulation that can be applied using water or sprayable fluid fertilizer as the carrier. If a fluid fertilizer is to be used, a compatibility test (see **COMPATIBILITY TEST** on page ) should be made prior to tank mixing. The use of fluid fertilizer as a carrier may damage emerged corn.

Ground or aerial application equipment which will give good spray coverage of weed foliage should be used. **HOWEVER, DO NOT USE AERIAL APPLICATION EQUIPMENT IF SENSITIVE CROPS ARE GROWING IN THE VICINITY OF THE AREA TO BE TREATED, OR WHEN TEMPERATURE INVERSIONS EXISTS.**

Apply 3 to 50 gallons of diluted spray per treated acre when using ground application equipment, or 2 to 10 gallons of diluted spray per treated acre when using aerial

PC-LAR:TA.95B

May 25, 1995

application equipment. Use the higher level of the listed spray volumes when treating dense or tall vegetation.

CLARITY should not be applied during periods of gusty wind or when wind is in excess of 15 mph as uneven spray coverage may occur. If spraying in vicinity of sensitive crops refer to "SENSITIVE CROP PRECAUTIONS".

Avoid disturbing (e.g. tillage or cultivating) treated areas for at least 7 days following application.

Up to 2 applications of **CLARITY** may be made during a growing season. Do not exceed a total of 24 fluid ounces of **CLARITY** per treated acre per crop year. Allow two weeks or more between applications of **CLARITY**. See appropriate section for rate information. For combination options or sequential treatments, refer to appropriate section.

SPRAY ADDITIVES

Sprayable fluid fertilizers (1/2-1 gallon per acre of 28%, 30% or 32% urea ammonium nitrate) or ammonium sulfate (2.5 lbs per acre) may be added to the spray mixture to improve postemergence velvetleaf control.

Agriculturally approved surfactants may also be added to the mixture to improve postemergence weed control, particularly in dry growing conditions.

The use of adjuvants containing penetrants, such as petroleum based oils, may be added to the mixture to improve postemergence weed control in dry growing conditions. Do not use penetrants after the corn exceeds five inches tall or crop injury may result.

For spray additive recommendations with tank mixtures refer to TANK MIX section of this label.


PREPLANT/PREEMERGENCE IN NO-TILLAGE CORN

Applications of **CLARITY** may be made before, during, or after planting to emerged and actively growing broadleaf weeds. Apply **CLARITY** at 16 fluid ounces per treated acre on medium or fine textured soils containing 2% or greater organic matter. Use 8 fluid ounces per treated acre on coarse textured soils (sand, sandy loam, and loamy sand) or medium and fine textured soils with less than 2% organic matter. The addition of a nonionic surfactant is recommended for improved spray coverage of emerged weeds.

When planting into a legume sod (e.g., alfalfa or clover), apply **CLARITY** after 4-6 inches of regrowth has occurred.

For added control of dandelion or plantain, 2,4-D at 1/4 to 1/2 pounds per acre may be tank mixed with **CLARITY**

PREEMERGENCE IN CONVENTIONAL OR REDUCED TILLAGE CORN



CLARITY may be applied after planting and prior to corn emergence. Application at 16 fluid ounces per treated acre may be made to medium or fine textured soils which contain 2% or greater organic matter. For best performance delay application 5-7 days following planting. **DO NOT** apply to coarse textured soils (sand, sandy loam, and loamy sand) until after crop emergence (see Early Postemergence uses below).

Preemergence application of **CLARITY** does not require mechanical incorporation to become active. A shallow mechanical incorporation is recommended if application is not followed by adequate rainfall or sprinkler irrigation. Avoid tillage equipment (e.g., drags, harrows) which concentrate treated soil over seed furrow.

**EARLY POSTEMERGENCE
(ALL TILLAGE SYSTEMS)
(Spike through 8 inch tall corn)¹**


CLARITY at 16 fluid ounces per treated acre may be applied during the period from corn emergence through 8 inch tall corn. Reduce the rate to 8 fluid ounces per treated acre if corn is growing on coarse textured soils (sand, sandy loam, loamy sand).

¹ When **CLARITY** is tank mixed with Accent or Beacon, applications may be made up through 12 inch tall corn. See TANK MIX section for more information.

OVERLAY (SEQUENTIAL) TREATMENTS

CLARITY may be applied to ground before or after treatment with one or more of the following herbicides:

- acetochlor (Surpass[®], Harness[®] Plus)
- alachlor (Lasso[®], Lasso[®] Micro Tech or Partner[®])
- Atrazine
- Bicep[®]
- Broadstrike[®] plus
- Broadstrike[®] + Dual[®]
- Bronco[®]
- Bullet[®]
- butylate (Sutan + [®])
- cyanazine (Bladex[®])
- dicamba (Banvel [®])
- dimethenamid (Frontier [®])
- EPTC (Eradicane[®])
- Extrazine[®]
- glyphosate (Roundup[®])
- Guardsman[®]



10 7 16

PCLARITA.95B

May 25, 1995

halosulfuron (Battalion[®], Permit[®])

Lariat[®]

Marksman[®]

metolachlor (Dual[®])

nicosulfuron (Accent[®])

paraquat (Gramoxone[®])

pendimethalin (Prowl[®])

primisulfuron (Beacon[®])

propachlor (Ramrod[®])

simazine (Princep[®])

Surpass[®] 100

Apply CLARITY at 8 ounces per treated acre on ground previously treated with full rates of Banvel or Marksman.


Apply Banvel at 1/2 pint per treated acre or Marksman at 2 pints per treated acre to ground previously treated with a full rate of CLARITY. Applications must be separated by two weeks or more.

READ AND FOLLOW LABEL DIRECTIONS FOR EACH OF THE ABOVE PRODUCTS.

TANK MIX TREATMENTS FOR CORN

CLARITY may be tank mixed with one or more of the following herbicides for control of grasses or additional broadleaf weeds. Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, and other restrictions. Tank mix recommendations are for use only in states where the tank mix product and application site are registered.

CLARITY + Accent[®]: Accent may be tank mixed with CLARITY for added postemergence grass control. Apply Accent at 1/2 - 1 ounce ai per acre to emerged weeds but before corn exceeds 12 inches tall. When corn is 8 - 12 inches tall use drop nozzles and spray under the corn canopy. This will avoid spray pattern interference and overdosing the corn plant. Refer to diagram below when post directing spray. Use non-ionic surfactant at 0.25% (v/v) with this tank mix. In dry growing conditions also use 28% UAN at 4% (v/v).



10

[AFFIX DIAGRAM]

Proper setup to broadcast when corn is 8 to 12 inches tall.

CLARITY + Atrazine : Apply Atrazine at 1 1/4 - 2 pounds ai per acre up to 8 inch tall corn. For added grass control crop oil concentrate may be added to this mixture if corn is less than 5 inches tall and grass weeds are emerged but less than 1 1/2 inches tall. Follow all application restrictions on the Atrazine label.

CLARITY + Beacon®: Beacon may be tank mixed with CLARITY for added postemergence grass control. Apply Beacon at 0.28 - 0.57 ounce ai per acre to emerged weeds but before corn exceeds 12 inches tall. When corn is 8 - 12 inches tall use drop nozzles and spray under the corn canopy. This will avoid spray pattern interference and overdosing the corn plant. Use non-ionic surfactant at 0.25% (v/v) with this tank mix. In dry growing conditions also use 28% UAN at 4 % (v/v). For proper boom positioning refer to diagram in the CLARITY + Accent tank mix section.

CLARITY + Bladex® : Apply Bladex at 1 1/4 - 4 pounds ai per acre prior to corn emergence. After corn emergence use only the 90DF formulation at 1 1/4 - 2 pounds ai per acre. Applications should be made before grasses are 1 1/2 inches and before the corn is beyond the 4 leaf stage.

CLARITY + Dual® : Apply Dual at 1 1/2 - 3 pounds ai per acre for added grass control. Applications can be made after planting but before corn exceeds 3 inches tall and before grasses exceed the 2 leaf stage. Applications prior to crop emergence may only be made on medium or fine textured soils containing 2 1/2 % or greater organic matter.

CLARITY + Frontier® : Apply Frontier at 13 - 25 fluid ounces per acre for preemergence grass control. Applications can be made after planting before corn exceeds 8 inches in height. This treatment must be combined with a herbicide that provides post emergent control of grass weeds if grasses are greater than 1 inch tall at the time of application.



PCLARITA.95B

May 25, 1995

Applications prior to crop emergence may only be made on medium or fine textured soils containing 2 1/2% or greater organic matter.

CLARITY + Gramoxone® : Apply Gramoxone at 1/4 - 1 pound ai per acre. Application may be made to emerged weeds but prior to corn emergence.

Clarity + Harness® Plus or Surpass® : Apply Harness® Plus or Surpass® at 1 1/2 - 3 pounds ai per acre. Applications may be made after planting and before corn emergence. Applications may only be made on medium or fine textured soils containing 2 1/2 % or greater organic matter.

CLARITY + Lasso®, Lasso® Micro Tech or Partner®: Apply Lasso, Lasso Micro Tech or Partner at 2 1/2 - 3 pounds ai per acre. Applications may be made after planting and before grass emergence. Applications may only be made on medium or fine textured soils containing 2 1/2 % organic matter or greater.

CLARITY + Princep® : Apply Princep at 1 - 3 pounds ai per acre prior to corn or weed emergence.

CLARITY + Prowl® : Apply Prowl at 3/4 - 1 1/2 pounds ai per acre. Applications can be made after planting before corn exceeds the 2 leaf stage of growth and before grass weed emergence. Application prior to crop emergence should only be made on medium or fine textured soils with 2 1/2 % or greater organic matter.

CLARITY + Roundup® : Apply Roundup at 1 - 3 pounds ai per acre to emerged weeds prior to crop emergence.


CLARITY + 2,4-D : Apply 2,4-D at 1/4 - 1/2 pounds ai per acre prior to crop emergence for added burndown of existing weeds.

CLARITY + Accent® + Dual® : Apply Accent at 1/4 - 1/2 ounces ai per acre and Dual at 3/4 - 1 1/2 pounds ai per acre. Application should be made after grass emergence and before grass is 3 inches tall. Use non-ionic surfactant at 0.25% (v/v) and 1/2 - 1 gallon liquid fertilizer with this tank mix.

CLARITY + Accent® + Frontier® : Apply Accent 75WDG at 1/4 - 2/3 ounces per acre and Frontier at 8 - 16 fluid ounces per acre. Application should be made from emergence up to 8 inch tall corn when weeds are emerged and actively growing. Use non-ionic surfactant at 0.25% (v/v) with this tank mix.

PROCEDURE FOR CLEANING SPRAY EQUIPMENT

The steps listed below are suggested for thorough cleaning of spray equipment following applications of CLARITY.



PCLARITA.95B

May 25, 1995

- 1) Hose down thoroughly the inside as well as outside surfaces of equipment while filling the spray tank half full of water. Flush by operating sprayer until the system is purged of the rinse water.
- 2) Fill tank with water while adding 1 quart of household ammonia for every 25 gallons of water. Operate the pump to circulate the ammonia solution through the sprayer system for 15 to 20 minutes and discharge a small amount of the ammonia solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
- 3) Flush the solution out of the spray tank through the boom.
- 4) Remove the nozzles and screens and flush the system with two full tanks of water.

The steps listed below are suggested for thorough cleaning of spray equipment used to apply **CLARITY** as a tank mix with wettable powders (WP) emulsifiable concentrates (EC), or other types of water-dispersible formulations. **CLARITY** tank mixes with water-dispersible formulations require the use of a water/detergent rinse.

- 5) Complete step 1.
- 6) Fill tank with water while adding 2 lbs. of detergent for every 40 gallons of water. Operate the pump to circulate the detergent solution through the sprayer system for 5 to 10 minutes and discharge a small amount of the solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
- 7) Flush the detergent solution out of the spray tank through the boom.
- 8) Repeat step 1, and follow with steps 2, 3, and 4.

STORAGE AND DISPOSAL


PROHIBITIONS

Do not contaminate water, food or feed by storage or disposal.

STORAGE

Store in original container in a well-ventilated area separately from fertilizer, feed and foodstuffs. Avoid cross-contamination with other pesticides. Spillage or leakage should be contained and absorbed with clay granules, sawdust, or equivalent material for disposal.

PESTICIDE DISPOSAL



PCLARITA.95B

May 25, 1995

Triple rinse pesticide from containers and use rinsates in the pesticide application. Wastes which cannot be used according to label instructions may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Plastic or Metal: After triple rinsing (or equivalent), offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities, such as burning of plastic containers. If burned, stay out of smoke.

BULK STORAGE AND DISPOSAL: To be printed on labeling for bulk use only

AGITATE BEFORE USE

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. This product may not be mixed, loaded, or used with 50 feet of all wells including abandoned wells, drainage wells, and sinkholes.

STORAGE

Ground water contamination may be reduced by diking and flooring of permanent liquid bulk storage sites with an impermeable material.

PESTICIDE DISPOSAL


Pesticide spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal and local procedures under Subtitle C or the Resource Conservation and Recovery Act.

BULK TANK MAINTENANCE

Follow clean-out directions in Dealer Bulk Handling Guide for CLARITY Herbicide listed under Bulk Storage Tank Requirements.

GENERAL


Consult Federal, State, or local disposal authorities for approved alternative procedures, such as limited burning.



15 of 16

LIMITATION OF WARRANTY AND LIMITATION OF LIABILITY

NOTICE: Read this Limitation of Warranty and Limitation of Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Sandoz or seller. All such risks shall be assumed by buyer or user. Sandoz warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, under normal use conditions, subject to the risks described above. **Sandoz makes no other express or implied warranty of fitness or of merchantability or any other express or implied warranty.** In no event shall Sandoz or seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **The exclusive remedy of the user or buyer, and the exclusive liability of Sandoz or seller for any and all claims, losses, injuries or damages (including claims based on breach of warranty, contract, negligence, tort, strict liability or otherwise) resulting from the use or handling of this product, shall be the return of the purchase price of the product or, at the election of Sandoz or seller, the replacement of the product.** Sandoz and seller offer this product, and buyer and user accept it, subject to the foregoing limitations of warranty and limitation of liability, which may not be modified by any oral or written agreement.



16 of 16

REGISTERED TRADEMARKS

Prowl® is a registered trademark of American Cyanamid Company.

Beacon® Bicep®, Dual®, and Princep® are registered trademarks of Ciba-Geigy Corporation, Agricultural Division.


Broadstrike ® plus and Broadstrike ® are registered trademarks of DowElanco, Inc.

Accent®, Bladex®, and Extrazine®, are registered trademarks of E. I. duPont de Nemours & Co., Inc.

Battalion®, Bronco®, Bullet®, Lariat®, Lasso® Micro Tech, Partner™, Permit®, Ramrod®, Harness® Plus and Roundup® are registered trademarks of Monsanto Agricultural Products Company.

Eradicane®, Gramoxone® and Surpass® and Surpass® 100, Sutan +® are registered trademarks of Zeneca Ag Products

Banvel® Herbicide, Frontier® Herbicide, Guardsman® Herbicide and Marksman® Herbicide are registered trademarks of Sandoz Ltd.



16