SANDOZ

IPA SALT OF DICAMBA

Active Ingredient: Isopropylamine salt of dicamba (2,6-dichloro-o-anisic acid)*	40.32
Isopropylamine salts of related acids	10.09
Inert Ingredient:	49.59
TOTAL 1	00.00%

*This product contains 31.81% 3,6-dichloro-o-anisic acid (dicamba) or 3 lbs. dicamba per gallon and 7.95% acids related to dicamba.

EPA Reg. No. 55947-

EPA Est. No. 55947-TX-1

ACCEPTED
7 JAN 1988
Under the Federal Inserticide.
Fungitation and Polymeric in the calcinomic of the control o

Keep Out of the Reach of Children WARNING
See Side Panel for Addtional Precautionary Statements

Net Contents:

Sandoz Crop Protection Corporation 341 East Ohio Street Chicago, Illinois 60611

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DUMESTIC ANIMALS

WARNING

Causes eye irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid breathing spray mist. Wash thoroughly after handling. In case of contact, wash skin with soap and water, for eyes flush with water for 15 minutes and get medical attention.

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams and ponds. Do not contaminate water by the cleaning of equipment or disposal of wastes. Apply this product only as directed on the label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Refer to USE DIRECTIONS Booklet attached to this container for proper use directions and additional precautionary statements. Apply this product only as directed on the label.

For additional copies of the USE DIRECTIONS Booklet, write to the manufacturer.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage or disposal.

STORAGE

Store in original container in a well-ventilated area separately from fertilizer, feed and foodstuffs. Avoid cross-contamination with other pesticides. Spillage or leakage should be contained and absorbed with clay granules, saw dust, or equivalent material for disposal.

PESTICIDE DISPOSAL

Triple rinse pesticide from containers and use these rinsates in the pesticide application. Other wastes which cannot be used according to label instructions may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Plastic or Metal: After triple rinsing (or equivalent), offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities, such as burning of plastic containers. If burned, stay out of smoke.

12

LIMITED WARRANTY AND LIABILITY

NOTICE: Read this Limited Warranty and Liability before buying or using this product. If the terms are not acceptable, return it at once unopened.

It is critical that this product be used only as specified on the label. The laws of a state may make some or all of this paragraph inapplicable or may give you rights in addition to your rights hereunder. Except to the extent prohibited by applicable law, the exclusive remedy of the user or buyer and the limit of liability of this Company or any other seller for any and all losses, personal injuries or damages resulting from the use of this product, shall be the purchase price paid by the user or buyer for the quantity of product involved. Except to the extent prohibited by state law, there is no warranty, and this company and other sellers disclaim all liability for losses, personal injury or damages, whether indirect, special or consequential, when such claim is not reported to this company within one year of discovery and: (i) arises form any use of this product in a manner or for a purpose not recommended in its label directions or (ii) arises from handling or storage in violation of label instructions. THERE ARE NO IMPLIED WARRANTIES OR AND NO WARRANTIES OF MERCHANTABILITY OR FITNESS.

IPA SALT OF DICAMBA

FOR USE IN REDUCED TILLAGE FALLOW SYSTEMS

DIRECTIONS FOR USE: IT IS VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION

ACCEPTED 7-JAN 1988

Under the Federal Insecticide, Fungicide, and Rolenteide fiel, as amended, for the pesticide registered under EPA Reg. No. EPA Reg. No. 55947-

SANDOZ CROP PROTECTION CORP. 341 EAST OHIO STREET CHICAGO, ILLINIOS 60611

CONTENTS

IMPORTANT
WEEDS CONTROLLED
RATES AND TIMINGS
TANK MIX TREATMENTS
PLANTBACK CROPS
REGISTERED TRADEMARKS

IPA SALT OF DICAMBA

For annual broadleaf weed control in Reduced Tillage Fallow (RTF) Systems.

DIRECTIONS FOR USE

BEFORE USING IPA SALT OF DICAMBA READ AND FOLLOW THE PRECAUTIONS APPEARING ON THE CONTAINER.

* * *

IMPORTANT

Do not contaminate irrigation ditches or water used for domestic purposes.

Do not graze or harvest for livestock feed prior to crop maturity.

Do not apply to sorghum grown for seed production.

Broadleaf crops are sensitive to IPA Salt of Dicamba and contact may result in crop injury. Do not apply when weather conditions favor drift from target areas.

Consult your local or state authorities for possible application restrictions and advice concerning these and other special local use situations.

REFER TO THE CONTAINER LABEL FOR INSTRUCTIONS CONCERNING DISPOSAL OF WASTE AND CLEANING RINSES.

WEEDS CONTROLLED

IPA Salt of Dicamba, when applied at recommended rates, will control the following ANNUAL broadleaf weeds or give growth suppression of the following PERENNIAL broadleaf weeds.

ANNUALS

buckwheat, wild chickweed, common clover cockle, corn cocklebur cowcockle gromwell henbit knotweed spp. kochia lambaquarters, common lettuce, prickly ma"weed mustard, tumble mustard, wild nightshade, cutleaf (<2") pennycress, field

pigweed, prostrate
pigweed, redroot
pigweed, smooth
pigweed, tumble
puncturevine
purslane, common
ragweed, common
shepherdspurse
spurry, corn
sunflower, common
sunflower, volunteer
tansymustard
thistle, musk
thistle, Russian
velvetleaf

PERENNIALS

alfalfa bindweed, field bursage dogbane, hemp sowthistle, perennial thistle, Canada whitetop

RATES AND TIMINGS

Apply IPA Salt of Dicamba as a broadcast or apot treatment to emerged and actively growing weeds during the fallow

period.

Ground or merial application equipment which will give good apray coverage of weed foliage should be used. Apply 5 to 40 gallons of diluted apray per treated acre when using ground application equipment or 3 to 5 gallons of diluted apray per treated acre when using merial application. Spray additives such as surfactants or oils may be used to enhance apray coverage and herbicide's penetration of weed foliage. Use the higher rate when treating dense weed growth.

- o For control of actively growing kochia, redroot pigweed, lambsquarters, wild buckwheat, and Russian thistle that are less than 4 inches in height, apply 8 ounces per treated acre.
- o For control of listed annual broadleaf weeds that are less than 6 inches in height, apply 13 ounces per treated acre.
- o For suppression of listed perennial broadleaf weeds, apply 13 ounces per treated acre.

Avoid disturbing treated areas for at least 7 days following application. These treatments may not kill weeds which develop from seed or underground plant parts, such as rhizomes or bulblets. A follow-up program for seedling control or other cultural practices should be instituted.

When sorghum is to be planted do not exceed 1 pint per growing season. See PLANTBACK CROPS section for recommended interval between application and planting to prevent crop injury.

Maximum application rate of the IPA salt of dicamba per growing season is 13 ounces (0.3 lbs. dicamba) per acre.

TANK MIX TREATMENTS

IPA Salt of Dicamba may be tank mixed with one or more of the following herbicides for control of grasses or additional annual broadleaf weeds. Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, weeds controlled and geographic or other restrictions.

Rate per treated acre
(lbs. a.i.)

atrazine
chlorsulfuron (Glean®)*

Rate per treated acre
(lbs. a.i.)

1//2 to 3
0.016 to 0.024

chlorsulfuron (Glean®)*

(1/3-1/2 wt. oz. prod.)

cyanazine (Bladex®)

glyphosate (Roundup®)

metribuzin (Sencor®, Lexone®)

paraquat

promamide (Kerb®)

2,4-D

0.016 to 0.024

(1/3-1/2 wt. oz. prod.)

1 3/5 to 3 1/5

1/4 to 1/2

1/3 to 3/4

1/2 to 1

1/4 to 3/8

1/4 to 1/2

* must use surfactant in tank mix

PLANTBACK CROPS

Wheat, sorghum, barley and oats may be planted into treated areas. Spot applications may be made anytime prior to crop emergence if crop injury can be tolerated in treated areas. Injury may occur if the interval between application and planting is less than specified. Delay planting 15 days per 1/2 pint IPA Salt of Dicamba per treated acre. Exclude days when ground is frozen.

REGISTERED TRADEMARKS

Glean and Lexone are registered trademarks of E.I. duPont de Nemours and Company Inc.

Sencor[€] is a registered trademark of Mobay Chemical Corporation

 $\operatorname{Roundup}^{\otimes}$ is a registered trademark of Monsanto Agricultural Products Company

Bladex® is a registered trademark of Shell Chemical Company

Kerb® is a registered trademark of Rohn and Haas Company

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION

USE PESTICIDES PROPERLY

PROSPER WITH PESTICIDES BY USING THEM PROPERLY! READ AND FOLLOW LABEL DIRECTIONS WARRANTY AND LIABILITY"
ON THE CONTAINER BEFORE
BUYING OR USING. IF
TERMS ARE NOT ACCEPTABLE,
RETURN AT ONCE UNOPENED

SANDOZ CROP PROTECTION CORP. 341 EAST OHIO STREET CHICAGO, ILLINOIS 60611