

PM 25 55947-47
Pg 13 12

(Container Label)

ISOPROPYLAMINE SALT OF DICAMBA

FOR USE IN REDUCED TILLAGE FALLOW SYSTEMS

Active Ingredient:

Isopropylamine salt of dicamba
(3,6-dichloro-*o*-anistic acid)* 40.32%

Inert Ingredients: 59.68%

TOTAL 100.0%

*This product contains 31.81% 3,6-dichloro-*o*-anistic acid (dicamba) or 3 pounds per gallon (360 g/L).

KEEP OUT OF REACH OF CHILDREN

WARNING

AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See side panel for additional precautionary statements

EPA Reg. No. 55947-47
Net Contents:
EPA Est. No. 55947-TX-1

SANDOZ AGRO, INC.
1300 EAST TOUHY AVENUE
DES PLAINES, IL 60018

~~CONFIDENTIAL~~
NOV 2 - 1994

HIPSADIA.94A

MAY 1994

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes eye irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid breathing spray mist. In case of contact, wash skin with soap and water; for eyes, flush with water for 15 minutes and get medical attention.

Personal Protective Equipment:

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams or ponds. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply this product only as directed on label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Do not apply this product through any type of irrigation system.

Refer to the DIRECTIONS FOR USE booklet attached to this container for proper use directions and additional precautionary statements.



STORAGE AND DISPOSAL

STORAGE

Do not contaminate water, food or feed by storage or disposal.

Store in original container in a well-ventilated area separately from fertilizer, feed and foodstuffs. Avoid cross-contamination with other pesticides. Spillage or leakage should be contained and absorbed with clay granules, sawdust, or equivalent material for disposal.

PESTICIDE DISPOSAL

Triple rinse pesticide from containers and use rinsates in the pesticide application. Wastes which cannot be used according to label instructions may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Plastic or Metal: After triple rinsing (or equivalent), offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities, such as burning of plastic containers. If burned, stay out of smoke.

LIMITED WARRANTY AND LIABILITY

NOTICE: Read this Limited Warranty and Liability before buying or using this product. If the terms are not acceptable, return it at once, unopened.

It is critical that this product be used and mixed only as specified on the label. The laws of a State may make some or all of this paragraph inapplicable or may give you rights in addition to your rights hereunder. Except to the extent prohibited by applicable law, the exclusive remedy of the user or buyer and the limit of liability of this Company or any other Seller for any and all losses, personal injuries or damages resulting from the use of this product shall be the purchase price paid by the user or buyer for the quantity of product involved. Except to the extent prohibited by State Law, there is no warranty, and this Company and other Sellers disclaim all liability for losses, personal injury or damages: (i) arising from any use of this product in a manner or for a purpose not recommended in its label directions or from mixing this product before use with any substance except as recommended by the product's label; (ii) arising from handling or storage in violation of label instructions; (iii) for all indirect, special or consequential damages; and (iv) when not reported to this Company within one year of discovery. **THERE ARE NO IMPLIED WARRANTIES AND NO WARRANTIES OF MERCHANTABILITY OR FITNESS.**

(DIRECTIONS FOR USE)

IPA SALT OF DICAMBA

DIRECTIONS FOR USE: IT IS VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

FOR USE IN REDUCED TILLAGE FALLOW SYSTEMS.

THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

EPA REG. NO.: 55947-47

**SANDOZ AGRO, INC.
1300 EAST TOUHY AVENUE
DES PLAINES, IL 60018**

CONTENTS

IMPORTANT

WEEDS CONTROLLED

RATES AND TIMINGS

TANK MIX TREATMENTS

PLANTBACK CROPS

REGISTERED TRADEMARKS

7-12

IPA SALT OF DICAMBA

For annual broadleaf weed control in Reduced Tillage Fallow (RTF) Systems.

DIRECTIONS FOR USE

BEFORE USING IPA SALT OF DICAMBA READ AND FOLLOW THE PRECAUTIONS APPEARING ON THE CONTAINER.

IMPORTANT

Do not contaminate irrigation ditches or water used for domestic purposes.

Do not graze or harvest for livestock feed prior to crop maturity.

Do not apply to sorghum grown for seed production.

Broadleaf crops are sensitive to IPA Salt of Dicamba and contact may result in crop injury. Do not apply when weather conditions favor drift from target areas.

Consult your local or state authorities for possible application restrictions and advice concerning these and other special local use situations.

REFER TO THE CONTAINER LABEL FOR INSTRUCTIONS CONCERNING DISPOSAL OF WASTE AND CLEANING RINSES.

WEEDS CONTROLLED

IPA Salt of Dicamba, when applied at recommended rates, will control the following ANNUAL broadleaf weeds or give growth suppression of the following PERENNIAL broadleaf weeds.

ANNUALS

buckwheat, wild
chickweed, common
clover
cockle, corn
cocklebur
cowcockle
gromwell
henbit
knotweed spp.
kochia
lambquarters, common
lettuce, prickly
mayweed
mustard, tumble
mustard, wild
nightshade, cutleaf (< 2")
pennycress, field

pigweed, prostrate
pigweed, redroot
pigweed, smooth
pigweed, tumble
puncturevine
purslane, common
ragweed, common
shepherdspurse
spurry, corn
sunflower, common
sunflower, volunteer
tansymustard
thistle, musk
thistle, Russian
velvetleaf

PERENNIALS

alfalfa
bindweed, field
bursage
dogbane, hemp

sowthistle, perennial
thistle, Canada
whitetop

RATES AND TIMINGS

Apply IPA Salt of Dicamba as a broadcast or spot treatment to emerged and actively growing weeds during the fallow period.

Ground or aerial application equipment which will give good spray coverage of weed foliage should be used. Apply 5 to 40 gallons of diluted spray per treated acre when using ground application equipment or 3 to 5 gallons of diluted spray per treated acre when using aerial application. Spray additives such as surfactants or oils may be used to enhance spray coverage and herbicide's penetration of weed foliage. Use the higher rate when treating dense weed growth.

- For control of actively growing kochia, redroot pigweed, lambsquarters, wild buckwheat, and Russian thistle that are less than 4 inches in height, apply 8 ounces per treated acre.
- For control of listed annual broadleaf weeds that are less than 6 inches in height, apply 13 ounces per treated acre.
- For suppression of listed perennial broadleaf weeds, apply 13 ounces per treated acre.

Avoid disturbing treated areas for at least 7 days following application. These treatments may not kill weeds which develop from seed or underground plant parts, such as rhizomes or bulblets. A follow-up program for seedling control or other cultural practices should be instituted.

When sorghum is to be planted do not exceed 1 pint per growing season. See PLANTBACK CROPS section for recommended interval between application and planting to prevent crop injury.

Maximum application rate of the IPA salt of dicamba per growing season is 13 ounces (0.3 lbs. dicamba) per acre.

TANK MIX TREATMENTS

IPA Salt of Dicamba may be tank mixed with one or more of the following herbicides for control of grasses or additional annual broadleaf weeds. Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, weeds controlled and geographic or other restrictions.

Herbicide	Rate per treated acre (lbs. a.i.)
atrazine	1/2 to 3
chlorsulfuron (Glean®)*	0.016 to 0.024 (1/3-1/2 wt. oz. prod.)
cyanazine (Bladex®)	1 3/5 to 3 1/5
glyphosate (Roundup®)	1/4 to 1/2
metribuzin (Sencor®, Lexone®)	1/3 to 3/4
paraquat	1/2 to 1
promamide (Kerb®)	1/4 to 3/8
2,4-D	1/4 to 1/2

*must use surfactant in tank mix

PLANTBACK CROPS

Wheat, sorghum, barley and oats may be planted into treated areas. Spot applications may be made anytime prior to crop emergence if crop injury can be tolerated in treated areas. Injury may occur if the interval between application and planting is less than specified. Delay planting 15 days per 1/2 pint IPA Salt of Dicamba per treated acre. Exclude days when ground is frozen.

BEST AVAILABLE COPY

REGISTERED TRADEMARKS

Glean® and Lexone® are registered trademarks of E.I. duPont de Nemours and Company, Inc.

Sencor® is a registered trademark of Mobay Chemical Corporation

Roundup® is a registered trademark of Monsanto Agricultural Products Company

Bladex® is a registered trademark of Shell Chemical Company

Kerb® is a registered trademark of Rohn and Haas Company

123 12

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION

USE PESTICIDES PROPERLY

PROSPER WITH PESTICIDES BY USING THEM PROPERLY! READ AND FOLLOW LABEL DIRECTIONS

NOTICE: READ "LIMITED WARRANTY AND LIABILITY" ON THE CONTAINER BEFORE BUYING OR USING. IF TERMS ARE NOT ACCEPTABLE, RETURN AT ONCE UNOPENED

SANDOZ AGRO, INC.
1300 EAST TOUHY AVENUE
DES PLAINES, IL 60018