


10/3/80

10f 3  
Pr. 1  
55947-48

10/3/80

**ACCEPTED**  
**OCT 30 1980**  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, by the pesticide registered under EPA Reg. No. 11273-1

**BEST AVAILABLE COPY**



# thio

## MICRONIZED W

ACTIVE INGREDIENT	BY WT
Sulfur	80%
INERT INGREDIENTS	20%
TOTAL	100%

### CAUTION! KEEP OUT OF REACH OF CHILDREN

Sulfur powder is a fire hazard. Do not use in a confined space. Breathing dusts or fumes may irritate the respiratory tract. Avoid contact with eyes.

Sulfur may irritate the skin. Avoid contact with the eyes. Do not use in a confined space. Breathing dusts or fumes may irritate the respiratory tract. Avoid contact with eyes.

#### DIRECTIONS FOR DILUTION

**Dilute Application:** Ground Application: Apply specified rate in 100 to 200 gallons of water per acre. (Do not apply to less than 100 gallons of water per acre.)

**Concentrate Application:** Ground Application: Apply specified rate in not less than 5 gallons of water per acre. Do not apply to less than 20 to 100 gallons of water per acre. See instructions for further application details.

**Air Application:** Apply specified rate in 30 to 200 gallons of water per acre. Do not apply to less than 30 gallons of water per acre.

#### DIRECTIONS FOR USE

In compliance with Federal law to control sprays, this product must be used with the following:

Properly calibrated spray tank or other application equipment. If necessary, add a few drops of oil to the tank.

#### Crops

**Almonds, Cherries, Plums, Prunes, Peaches, Nectarines:** Brown mite, Powdery mildew, Leaf spot, Silver mite, Flat mite, Red spider mite. Apply 10 to 30 lbs. per acre. Apply at bloom or early petal fall. Repeat as necessary, usually 10 to 14 days or after a period of wet weather.

**Apples, Pears:** Powdery mildew, Scab, Blister mite, Red spider mite. Apply 10 to 30 lbs. per acre. Do not apply to Anjou pears. Apply pre-bloom and petal fall. Apply with cover sprays throughout the season.

**Asparagus:** Rust. Apply 10 to 30 lbs. per acre. Use after cutting stops. Irrigate and cultivate before applying the sulfur. Repeat at 7 to 10 day intervals throughout the season.

**Avocados:** Brown mite. Apply 10 to 30 lbs. per acre as needed.

**Beans, Peas, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kale, Mustard Greens, Rutabagas, Turnips, Peppers, Potatoes:** Powdery mildew, Septoria leaf spot, Red spider mite. Apply 3 to 5 lbs. per acre. Start in early leaf stage. Repeat application every 10 to 14 days with or without pesticides immediately after a rain.

**Blackberries, Boysenberries, Dewberries, Loganberries:** Powdery mildew. Apply 6 to 15 lbs. per acre. Begin when new shoots are 6 inches long and before blossoms open. Repeat at 10 day intervals or as necessary.

**Blueberries, Gooseberries, Huckleberries, Currants:** Powdery mildew. Apply 6 to 15 lbs. per acre. Begin when disease first appears. Repeat at 7 to 10 day intervals or as necessary.

**Carrots, Onions, Garlic:** Petiole mite and Powdery mildew. Apply 3 to 10 lbs. per acre when the disease first appears and repeat as needed.

NET WT.  
LOT



**BEST AVAILABLE COPY**

# olux<sup>®</sup>

## ETTABLE SULFUR

**Citrus:** Flat mite, Silver mite, Rust mite, Red spider mite. Apply 8 to 20 lbs per acre. Apply in any spray—NOT CONTAINING OIL—or following oil by 14 days

**Corn, Small Grains, Sorghum, Soybeans:** For suppression of the Red spider mite, Two-spotted mite, Grass banks mite, and Pacific mite. For control of the Atlantic mite (small grain only) and for control of Powdery mildew. Apply 6 to 15 lbs per acre. Apply when disease first appears and repeat as necessary or with the regularly scheduled insect program

**Cotton, Alfalfa:** Red spider mite. Apply 3 to 5 lbs per acre as needed

**Grapes:** Powdery mildew, Bud mite, Blister mite, Red spider mite. Apply 2 to 5 lbs per acre. Apply in late dormant or with cutworm spray, early shoot period and through season on 14 day schedule or with pesticides

**Mangoes:** Powdery mildew. Apply 10 to 30 lbs per acre. Begin before bloom. Repeat in bloom, after fruit set and 3 weeks later or as necessary

**Melons:** Powdery mildew. Use 20 to 25 lbs per acre. Treat when the disease first appears and repeat as necessary. Sulfur can injure plants, especially when the temperature reaches 95°F. Do not use on sulfur sensitive varieties.

**Peanuts:** Powdery mildew, Red spider mite, Leaf spot. Apply 3 to 5 lbs per acre. Start in early leaf stage. Repeat applications as necessary, usually 10 to 24 days

**Raspberries:** Powdery mildew (*Sphaerotheca humuli*). Apply 6 to 15 lbs per acre. Spray each week starting at first bloom color. Continue until all fruit is set, then repeat as weather requires

**Strawberries:** Powdery mildew, Red spider mite. Apply 5 to 10 lbs per acre. Start in early leaf stage, repeat applications with or without pesticides. For Powdery mildew, begin at first signs of disease and repeat as necessary. Do not use on sulfur sensitive varieties

**Sugar Beets, Table Beets:** Powdery mildew, Red spider mite. Apply 3 to 5 lbs per acre as needed with or without other pesticides

**Tomatoes:** Russet Mite. Apply 3 to 5 lbs per acre. Thorough coverage is required.

**Walnuts:** Pacific mite, Almond mite, Two-spotted mite, Red spider mite. Apply 10 to 20 lbs per acre. Apply with mite sprays during season to increase the effectiveness. Caution. Certain varieties of walnuts are sensitive to sulfur sprays under certain conditions. Do not apply unless the varieties are known to be tolerant of sulfur.

### Ornamentals

Apply 3 to 5 lbs per 100 gallons of aqueous spray, ensuring complete plant coverage. Begin when disease first appears and repeat at 5 to 10 day intervals

**Asters, Carnations:** For control of Powdery mildew.

**Calendula, Cherry Laurels, Chrysanthemums, Hollyhock, Smilax:** For control of Leaf spot and Powdery mildew

**Clematis, Columbine, Ivy (English), Laurel, Petunia, Sage:** For control of Leaf spot.

**Cosmos, Crepe Myrtle, Dahlia, Delphinium, Gladiolus, Ligustrum, Sunflowers, Sweet Peas, Violets, Zinnias:** For control of Powdery mildew.

**Roses:** For control of Powdery mildew and Black spot

### STORAGE AND DISPOSAL

**Storage:** Do not store near food or feed.

**Disposal:** Dispose of empty bag by burning (keep out of smoke).

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. The manufacturer makes no other warranties of any kind, express or implied. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Inc.  
Protection  
Camino Del Rio So.  
Diego, California 92108

**EPA EST. NO. 36841-FR-01**  
**REG. NO. 11273-1**