

## LASSIE INDOOR/OUTDOOR DOG & CAT REPELLENT

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with skin. Wash hands with soap and water after spraying. Do not use in commercial food processing or preparation areas. In the home, all food processing surfaces and utensils should be covered during treatment. Cover exposed food. Do not apply to food crops.

#### STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water; get medical attention if irritation persists.

#### PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Never throw container into fire or incinerator.

#### ACTIVE INGREDIENTS:

Methyl nonyl Ketone

1.9%

Related compound

0.1%

#### INERT INGREDIENTS

98.0%

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**INDOORS:** To break your pet of undesirable habits (such as sleeping on the furniture, rugs and other forbidden areas in your home), spray the article of furniture, rugs, etc. lightly with this product so that the area or object is slightly moistened. Before spraying extensive areas of rugs, fabrics or plastics, test the effect of this product by first treating a small inconspicuous area. This product may stain or soften some fabrics or plastics. An alternate method to avoid possible damage is to spray a piece of cloth lightly and place it on the object or area in your home to be protected. Repeat daily or as necessary until undesirable habit is broken.


**OUTDOORS:** To prevent your pet or stray animals from soiling flower beds, areas of lawn, trees, shrubbery or evergreen, spray the edges of flower beds, trunks of trees, the bases of shrubbery and evergreens, patches of lawn, etc. lightly every day or every other day with this product. Stray animals are frequently attracted to areas where urine or feces have been deposited. First remove any droppings, then spray the spot with this masking odor. Reapply as often as necessary until undesirable habits of your pet or stray animals are changed. Do not apply directly to foliage of shrubs or to ornamental soft-bodied plants.

**GENERAL:** For both indoor and outdoor situations the most effective control for a pet is the sharply spoken word backed up with disciplinary action. This repellent product should be used in conjunction with training.

#### STORAGE AND DISPOSAL

**STORAGE:** Store in a cool area away from heat or open flame.

**DISPOSAL:** Do not reuse empty container. Wrap container and put in trash collection.



Distributed by Carter-Wallace, Inc., Cranbury, NJ 08512  
EPA Reg. No. 8220-17 EPA Est. 7168 CA 3 Product No. 46310 CH 46310 01  
LASSIE is a registered trademark of LASSIE Television, Inc.

# LASSIE®

## INDOOR/OUTDOOR DOG AND CAT REPELLENT



Helps Keep Pets and Strays Away From  
Shrubs, Furniture and "Forbidden" Areas

KEEP OUT OF  
REACH OF CHILDREN  
**CAUTION**  
See back for additional  
precautions and ingredients

MADE IN U.S.A.

NET WT. 14 OZ.

NOTE  
LABEL  
PER PRODUCT

DATE

BEST AVAILABLE COPY

NOV 02 1990

Ms. Ada M. Breaux  
Sandoz Crop Protection Corporation  
1300 East Touhy Avenue  
Des Plaines, Illinois 60018

Dear Ms. Breaux:

Subject: ThuricideR 32 LV  
EPA Registration No. 55947-69  
Your submission of September 28, 1990

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

NOTE: This approval is for the following:

1. Increase application rates and add BIU/acre equivalent.
2. Deletion of directions for single applications for gypsy moth.

Sincerely yours,

Phil Hutton  
Product Manager (17)  
Insecticide-Rodenticide Branch  
Registration Division (H7505C)

BEST AVAILABLE COPY

# THURICIDE® 32 LV

AQUEOUS CONCENTRATE FOR AERIAL OR GROUND APPLICATION  
FOR CONTROL OF ORNAMENTAL, SHADE TREE, AND FOREST PESTS

## ACTIVE INGREDIENT:

Bacillus thuringiensis, subspecies kurstaki, potency of 8,000  
International Units (at least 12 million viable spores) per milligram\* .. 1.6%

INERT INGREDIENTS: ..... 98.4%

TOTAL ..... 100.0%

\*Equivalent to 8.0 billion International Units per quart.

KEEP OUT OF REACH OF CHILDREN

CAUTION

EPA Reg. No. 55947-69  
EPA Est. No. 55947-CA-2

Net Contents: 53 Gallons (200 L)

THURICIDE® is a Registered Trademark of Sandoz Ltd.

SANDOZ CROP PROTECTION CORPORATION  
Des Plaines, Illinois 60018

September 1990

## **PRECAUTIONARY STATEMENTS**

### **HAZARDS TO HUMANS**

Avoid inhalation or contact with eyes or open wounds.

### **ENVIRONMENTAL HAZARDS**

Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Aerial application is permissible only above tree canopy. Do not contaminate water when disposing of equipment washwaters.

### **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Suspension must be shaken or stirred before use. Pour the recommended amount of THURICIDE® 32LV into the required volume of water in the spray tank. Agitate as necessary to maintain suspension. Do not allow diluted sprays to remain in the tank for more than 72 hours. THURICIDE® 32LV is formulated to provide desirable coverage and stickability on leaf surfaces. Additional adjuvants, spreaders, or stickers may be added but are not essential.

Do not apply this product through any type of irrigation.

#### **GROUND APPLICATION**

Use adequate water to obtain good foliar coverage. Wet foliage but do not allow excessive run-off. Apply the recommended per acre rates of THURICIDE® 32LV with the following suggested amounts of water:

100 gallons with high-gallonage hydraulic sprayers  
10 gallons with mist blower

#### **AERIAL APPLICATION**

Apply THURICIDE® 32LV at recommended rates as a spray mix diluted with water. Use not less than one half gallon spray mix per acre. Best results are obtained when THURICIDE® 32LV is applied to dry foliage with a calibrated aircraft capable of obtaining droplet sizes below 300 microns and preferably in the range of 50-150 microns.

## RECOMMENDATIONS FOR SHADE TREES, ORNAMENTALS, AND FOREST

Thorough coverage is essential when using THURICIDE® 32LV. Use the lower rate for light to moderate infestations. Use the higher rates for heavier infestations.

---

Pest	Ounces Per Acre	Dosage BIU Per Acre	Directions For Use
Spring Cankerworm	16- 64	4-16	Apply when leaf expansion reaches 40-50% as infestation warrants. If eggs hatch over a long period of time or if reinfestation occurs, respray about 14 days after the first application.
Fall Cankerworm	16- 64	4-16	
Elm Spanworm	16- 64	4-16	
Tent Caterpillar	16- 64	4-16	
Gypsy Moth	32-160	8-40	
Spruce Budworm	24-160	6-40	Apply when most larvae are 3rd-4th instar. Also consider the opening of the bud cap to ensure foliage exposure.
Douglas Fir Tussock Moth	16-128	4-32	Apply after eggs have hatched and early instar larvae are feeding on exposed foliage.
Jack Pine Budworm	16- 64	4-16	
Bagworm	16- 64	4-16	
California Oak Moth	16- 64	4-16	
Western Tussock Moth	16- 64	4-16	
Fruit Tree Leafroller	16- 64	4-16	
Mimosa Webworm	16- 64	4-16	
Redhumped Caterpillar	8- 48	2-12	
Fall Webworm	8- 32	2- 8	
Pine Butterfly	24- 96	6-28	

---