

Evaluation:

If a uniform mixture cannot be made, the mixture should not be used.

If either mixture remains uniform for 30 minutes, the combination may be used.

Should either mixture separate after 30 minutes, but readily remixes uniformly with ten jar inversions, the mixture can be used if adequate agitation is maintained in the tank.

If the mixture with adjuvant is satisfactory, but the one without adjuvant is not, be sure to use the adjuvant in the spray tank.

If adjuvant is needed, add it first at a rate of 3 pint per 100 gallons of fluid fertilizer or water.

Foaming can be minimized by using moderate agitation.

If nondispersible oil, sludge or clumps of solids form in the mixture, the combination should not be used.

Rate Table for ERADICANE 6.7-E in Compatibility Tests

Table 7.

Gallons of Fluid Fertilizer to be Applied per Acre	Amount of Adjuvant to Add to the 1 Pint of Liquid		mL or Teaspoon to be Added to 1 Pint of Liquid for the ERADICANE 6.7-E rate listed below							
			3.75 Pints		4.75 Pints		6 Pints		7.33 Pints	
	mL	Tsp	mL	Tsp	mL	Tsp	ml	Tsp	mL	Tsp
10	2	2/5	22.0	4-2/5	28.0	5-3/5	35.0	7	42.0	8-2/5
15	2	2/5	16.0	3-1/5	20.0	4	25.0	5	30.0	6
20	2	2/5	13.0	2-3/5	16.0	3-1/5	20.0	4	24.0	4-4/5
25	2	2/5	9.0	1-4/5	12.0	2-2/5	15.0	3	18.0	3-3/5
30	2	2/5	6.0	1-1/5	8.0	1-3/5	10.0	2	12.0	2-2/5
40	2	2/5	6.0	1-1/5	8.0	1-3/5	10.0	2	12.0	2-2/5

APPENDIX II

Impregnation on Dry Bulk Fertilizers

Consult your local dealer for more details.

ERADICANE 6.7-E alone and tankmix combinations with atrazine and/or Bladex may be impregnated on dry bulk fertilizer.

CAUTION: ERADICANE 6.7-E alone and in combination tankmixes must not be impregnated on ammonium nitrate, potassium nitrate or sodium nitrate fertilizers. Such mixtures may cause explosion and fire.

All individual state regulations relating to dry bulk fertilizer blending, registration, labeling and application are the responsibility of the individual and/or company selling the herbicide and fertilizer mixtures.

Approved Dry Bulk Fertilizer Ingredients for Use with ERADICANE 6.7-E alone

Table 8.

Fertilizer Ingredient	N	P	K
Ammonium sulfate	21	0	0
Diammonium phosphate	18	46	0
Potassium chloride	0	0	60
Potassium sulfate	0	0	52
Single Super-phosphate	0	20	0
Treble Super-phosphate	0	46	0
Urea*	45	0	0
Ammonium phosphate-sulfate	16	20	0
K-Mag/Sul-Po-Mag	0	0	21
Monoammonium phosphate	11	48	0

*Some ureas may be phytotoxic when high rates are applied to corn.
 Use only urea rates known to be safe for corn application.

Approved Dry Bulk Fertilizer Ingredients for Use with ERADICANE 6.7-E in tankmix combinations with atrazine and/or Bladex

Table 9.

Fertilizer Ingredient	N	P	K
Ammonium sulfate	21	0	0
Diammonium phosphate	18	46	0
Potassium chloride	0	0	60
Potassium sulfate	0	0	52
Single Super-phosphate	0	20	0
Treble Super-phosphate	0	46	0
Urea*	45	0	0
Ammonium phosphate-sulfate	16	20	0
K-Mag/Sul-Po-Mag	0	0	21
Monoammonium phosphate	11	48	0

*Some ureas may be phytotoxic when high rates are applied to corn.
 Use only urea rates known to be safe for corn application.

Do not combine ERADICANE 6.7-E plus atrazine 4L or 80W or BLADEX 4L or 80W with 0-20-0 or 0-46-0.

Uniform impregnation of the herbicides on dry bulk fertilizer particles and uniform application in the field are necessary to assure good results.

A minimum of 200 and a maximum of 700 pounds of approved impregnated ingredients from Tables 8 and 9 must be applied per acre.

Use a closed rotary drum mixer or similar type of closed blender equipped with suitable spray equipment.

The spray nozzle should be positioned inside of the mixer to provide uniform spray coverage of the tumbling fertilizer and provide a uniform fine spray pattern.

Tankmix combinations may be added separately or mixed in the proposed use ratio in a uniform slurry for joint spray impregnation.

Physical properties of fertilizers vary in liquid absorptive capacity. When absorptivity is sufficient, simple spray impregnation of the fertilizer with the herbicides provides a satisfactory, dry mixture.

When the absorptive capacity is not adequate, use of a drying agent is required to provide dry, free-flowing mixtures.

Drying agent for spinning-disc applicators are:

Micro-Cel™ E calcium silicate powder

Drying agents for pneumatic applicators are:

Micro-Cel E calcium silicate powder
Agsorb™ 16/30 R /M-MS granular clay
Celatom™ MP-79

Drying agents should be added separately and uniformly to the previously impregnated herbicide-fertilizer mixture to insure that the mixture is free-flowing. Generally the following amounts are sufficient:

Micro-Cel E calcium silicate powder	less than 2% by weight
Agsorb 16/30 RVM-MS granular clay	less than 5% by weight
Celatom MP-79	less than 5% by weight

The amount of ERADICANE 6.7-E, atrazine and/or Bladex actually required in the manufacture of individual fertilizer mixtures should be determined carefully for each production operation (see Tables 10 and 11). This is necessary to ensure that the amount of herbicide actually contained in the mixture applied to the soil represents the correct use rate.

Physical Data:

Specific Gravity: 68°F/20°C: 0.972 (typical)
Pounds/Gallon: 8.09 (typical)
Flashpoint: 186°F/86°C (Tagliabue closed cup)
Viscosity: Sprayable down to minus 20°F/-29°C

RATE CHART FOR IMPREGNATION OF DRY BULK FERTILIZER WITH ERADICANE 6.7-E AND ATRAZINE AND BLADEx PR

Table 10.

Fertilizer Rate Pounds Per Acre	ERADICANE 6.7E Rate per Acre			Atrazine Alone in Tankmix				Bladex PR	
	3.75	4.75	7.33	Rate of 4L per ace		Rate of 80W per acre		Rate of 4L per	
				2	3	1½	2	3	
200	18-3/4 qts	23-3/4 qts	36-2/3 qts	10 qts	15 qts	12-1/2 lbs	20 lbs	15 qts	
250	15 qts	19 qts	29-1/3 qts	8 qts	12 qts	10 lbs	16 lbs	12 qts	
300	12-1/2 qts	15-4/5 qts	24-2/5 qts	6-2/3 qts	10 qts	8-1/3 lbs	13-1/3 lbs	10 qts	13
350	10-3/4 qts	13-3/5 qts	21 qts	5-3/4 qts	8-3/5 qts	7-1/5 lbs	11-2/5 lbs	8-3/5 qts	11
400	9-2/5 qts	11-7/8 qts	18-1/3 qts	5 qts	7-1/2 qts	6-1/4 lbs	10 lbs	7-1/2 qts	
450	8-1/3 qts	10-1/2 qts	16-1/3 qts	4-1/2 qts	6-2/3 qts	5-2/3 lbs	8-7/8 lbs	6-2/3 qts	6
500	7-1/2 qts	9-1/2 qts	14-2/3 qts	4 qts	6 qts	5 lbs	8 lbs	6 qts	
550	6-4/5 qts	8-2/3 qts	13-1/3 qts	3-2/3 qts	5-1/2 qts	4-2/3 lbs	7-1/3 lbs	5-1/2 qts	7
600	6-1/4 qts	8 qts	12-1/5 qts	3-1/3 qts	5 qts	4-1/5 lbs	6-2/3 lbs	5 qts	6
650	5-3/4 qts	7-1/3 qts	11-1/4 qts	3-1/5 qts	4-3/5 qts	3-7/8 lbs	6-1/5 lbs	4-3/5 qts	6
700	5-1/3 qts	6-4/5 qts	10-1/2 qts	2-7/8 qts	4-1/4 qts	3-2/3 lbs	5-3/5 lbs	4-1/4 qts	5

RATE CHART FOR IMPREGNATION OF DRY BULK FERTILIZER WITH ERADICANE 6.7-E AND ATRAZINE AND BLADAX PER TON OF FERTILIZER.

Table 11.

Fertilizer Rate Pounds Per Acre	ERADICANE 6.7-E Rate per Acre			Atrazine Rate for a 3-Way Tankmix				Bladax Rate for a 3-Way Tankmix			
				Rate of 4L per acre		Rate of 80W per acre		Rate of 4L per acre		Rate of 80W per acre	
	3.75	4.75	7.33	1	2	3/4	1 1/4	2	4	1 1/4	2 1/2
200	18-3/4 qts	23-3/4 qts	36-2/3 qts	5 qts	10 qts	7-1/2 lbs	12-1/2 lbs	10 qts	20 qts	12-1/2 lbs	25 lbs
250	15 qts	19 qts	29-1/3 qts	4 qts	8 qts	6 lbs	10 lbs	8 qts	16 qts	10 lbs	20 lbs
300	12-1/2 qts	15-4/5 qts	24-2/5 qts	3-1/3 qts	6-3/5 qts	5 lbs	8-1/3 lbs	6-3/4 qts	13-1/3 qts	8-1/3 lbs	16-2/3 lbs
350	10-3/4 qts	13-3/5 qts	21 qts	2-7/8 qts	5-4/5 qts	4-3/4 lbs	7-1/8 lbs	5-4/5 qts	11-2/5 qts	7-1/8 lbs	14-1/4 lbs
400	9-2/5 qts	11-7/8 qts	18-1/3 qts	2-1/2 qts	5 qts	3-3/4 lbs	6-1/4 lbs	5 qts	10 qts	6-1/4 lbs	12-1/2 lbs
450	8-1/3 qts	10-1/2 qts	16-1/3 qts	2-1/4 qts	4-2/5 qts	3-1/3 lbs	5-1/2 lbs	4-2/5 qts	8-7/8 qts	5-1/2 lbs	11-1/8 lbs
500	7-1/2 qts	9-1/2 qts	14-2/3 qts	2 qts	4 qts	3 lbs	5 lbs	4 qts	8 qts	5 lbs	10 lbs
550	6-4/5 qts	8-2/3 qts	13-1/3 qts	1-7/8 qts	3-2/3 qts	2-3/4 lbs	4-2/3 lbs	3-2/3 qts	7-1/3 qts	4-2/3 lbs	9-1/8 lbs
600	6-1/4 qts	8 qts	12-1/5 qts	1-2/3 qts	3-1/3 qts	2-1/2 lbs	4-1/5 lbs	3-1/3 qts	6-2/3 qts	4-1/5 lbs	8-1/3 lbs
650	5-3/4 qts	7-1/3 qts	11-1/4 qts	1-3/5 qts	3-1/8 qts	2-1/3 lbs	3-7/8 lbs	3-1/8 qts	6-1/5 qts	3-7/8 lbs	7-3/4 lbs
700	5-1/3 qts	6-4/5 qts	10-1/2 qts	1-1/2 qts	2-7/8 qts	2-1/5 lbs	3-2/3 lbs	2-7/8 qts	5-3/4 qts	3-2/3 lbs	7-1/5 lbs

35
 937

PESTICIDE DISPOSAL: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Metal - Triple rinse (or equivalent), then offer for recycling or recondition, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Plastic: Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

FOR BULK/MINI-BULK CONTAINERS

CONTAINER DISPOSAL: Reseal container and offer for reconditioning, or triple rinse (or equivalent) and offer for recycling or reconditioning, or clean in accordance with manufacturer's instructions.

CONTAINER PRECAUTIONS

Before refilling, inspect thoroughly for damage such as cracks, punctures, bulges, dents, abrasions, and damaged or worn threads on closure devices.

REFILL ONLY WITH ERADICANE 6.7-E. The contents of this container cannot be completely removed by cleaning. Refilling with materials other than ERADICANE 6.7-E will result in contamination and may weaken container.

After filling and before transporting, check for leaks.

Do not refill or transport damaged or leaking container.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.

CONDITIONS OF SALE AND LIMITED WARRANTY:

The Directions for Use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended or other influencing factors in the use of the product, all of which are beyond the control of the seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label, subject to the inherent risks referred to above, when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller and Buyer and User assume the risk of any such use. SELLER DISCLAIMS ALL OTHER WARRANTIES EXPRESSED OR IMPLIED INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY.

When Buyer or User claims losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability or other legal theories), Buyer or User must promptly notify in writing Seller of any claims to be eligible to receive either of the remedies set forth below. The EXCLUSIVE REMEDY OF BUYER OR USER and the LIMIT OF LIABILITY of Seller will be, at the election of Seller, refund of the purchase price paid for product bought, or replacement of amount of product used. SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND SELLER'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE.

- ERADICANE® is a trademark of a ZENECA Group Company.
- Agsorb™ is a trademark of Oil Dri Corp. of America.
- Banvel® is a trademark of Sandoz Crop Protection Corporation.
- Bladex® is a trademark of E. I. du Pont de Nemours & Co. (Inc.).
- Caliber® and Princep® are trademarks of CIBA-GEIGY Corporation.
- Celatom™ is a trademark of Eagle-Picher Minerals, Inc.
- Compex™ is a trademark of Kalo, Inc.
- Micro-Cel™ is a trademark of Manville Sales Corporation.
- Sponto™ is a trademark of Witco Corporation.
- Unite™ is a trademark of HACO, Inc.

PM 18 55947-82
United States Environmental Protection Agency
Washington, D.C. 20460

File
Page 1910

OCT 12 1993

Office of
Prevention, Pesticides and
Toxic Substances

Jonathan E. Bryant
SANDOZ AGRO, INC.
1300 EAST TOUHY AVENUE
DES PLAINES IL 60018

Subject: Label Amendment Submission of 06/10/93
in Compliance with WPS Labeling Requirements
EPA Reg No. 55947-82
ENSTAR II INSECT GROWTH REGULATOR

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is accepted subject to the comments listed below.

Based on your certification, the Agency has accepted only those changes to your labeling which are necessary to comply with PR Notices 93-7 and 93-11, which reflect the WPS labeling requirements of 40 CFR part 156, subpart K. Any other labeling changes submitted in connection with this amendment application and not directly related to compliance with PR Notice 93-7 or 93-11 have neither been reviewed nor accepted by the Agency. If you wish to make any such changes, you must submit a separate amendment application proposing them. If your product registration is currently suspended, acceptance of this labeling amendment does not affect the suspension in any way.

A copy of your proposed labeling stamped "Accepted with comments" is attached. Make any required changes described in the attached and send three copies of final labeling as soon as it is available to:

Document Processing Desk (FIN-LABEL)

2 7 10

Office of Pesticide Programs (H-7504C)
U.S. Environmental Protection Agency
401 M Street SW
Washington, DC 20460-0001


Hand or courier deliveries of final labels may be made to:

Document Processing Desk (FIN-LABEL)
Room 266A Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Please correct the typographical errors circled on the draft before printing final labeling.

In your final labeling the "Agricultural Use Requirements" text must be contained in a clearly separate box. This box may be set apart by a line, by another graphical device, by a different color background, or in any other way that clearly distinguishes it from surrounding text.

Sincerely,



James Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

3 of 10

CONTAINER LABEL

ENSTAR® 5E INSECT GROWTH REGULATOR

For Control of Whiteflies, Aphids,
Soft and Armored Scales, Mealybugs and
Fungus Gnats in Greenhouses and Interior Scapes
on Ornamental Plants and Vegetable Seed Crops

ACTIVE INGREDIENT

Kinoprene [2-propynyl (E,E)-3,7,11-trimethyl-2,4-dodecadienoate]*	65.3%
INERT INGREDIENTS	<u>34.7%</u>
Total	100.0%

Contains 0.625 lb. active ingredient per pint

ENSTAR® is a registered trademark of Sandoz Ltd., *U.S. Patent 4021461

EPA Reg. No. 55947-82
Net Contents

EPA Est. No. 55947-CA-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

ACCEPTED
with COMMENTS
In EPA Letter Dated

OCT 12 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
55947-82

See Side Panel For Additional Precautionary Statements

Sandoz Agro, Inc.
Des Plaines, IL 60018

ENSTAR®E

June 23, 1993

4 7 10

DIRECTIONS FOR USE

**ENSTAR® 5E
INSECT GROWTH REGULATOR**

**For Control of Whiteflies, Aphids,
Soft and Armored Scales, Mealybugs and
Fungus Gnats in Greenhouses and Interior Scapes
on Ornamental Plants and Vegetable Seed Crops**

ACTIVE INGREDIENT

Kinoprene [2-propynyl (E,E)-3,7,11-trimethyl-2,4-dodecadienoate]* 65.3%
INERT INGREDIENTS 34.7%

Total 100.0%

Contains 0.625 lb. active ingredient per pint

ENSTAR® is a registered trademark of Sandoz Ltd., *U.S. Patent 4021461

EPA Reg. No. 55947-82
Net Contents

EPA Est. No. 55947-CA-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side Panel For Additional Precautionary Statements

Sandoz Agro, Inc.
Des Plaines, IL 60018

ENSTAR®E

June 23, 1983

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash thoroughly after handling. Do not store near or contaminate food or feedstuffs.

Personal Protective Equipment

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for category F on an EPA chemical resistance category section chart.

Applicators and other handlers must wear:

Long-sleeved shirt and long pants

Chemical-resistant gloves, such as Barrier laminate, Butyl rubber, Nitrile rubber, or Viton

Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:
Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately and wash thoroughly. Wash your hands before eating, drinking, chewing gum, using tobacco or using the toilet.

BEST AVAILABLE COPY

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

BEST AVAILABLE COPY

ADMINISTRATIVE USE ONLY

[REDACTED]

Do not enter any area until the treated area has been cleared for a minimum (REI) of 12 hours.

PPE required for entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

Coveralls

Chemical resistant gloves, such as nitrile, neoprene, Viton

Shoes or socks

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Container Disposal:

Triple rinse (or equivalent), then dispose of in a sanitary landfill or by other approved state and local procedures.

Pesticide Disposal:

Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to federal, state or local procedures.

BEST AVAILABLE COPY

8 7 10

INSTRUCTIONS FOR USE

1. Partially fill spray tank with water, then add recommended amount of ENSTAR® 5E, agitate and complete filling.
2. Diluted spray solution should be used within twenty-four (24) hours; always agitate before using.

RECOMMENDED GREENHOUSE APPLICATIONS RATES OF APPLICATION

Preventative Program: Before insect populations reach damaging levels, mix ENSTAR® 5E at the rate of 6-10 ounces per 100 gallons of water (1 tsp./2 gal.) and apply to runoff every two (2) weeks.

Curative Program: For an existing population of adult insects, make two (2) applications seven (7) days apart of ENSTAR® 5E to runoff at a rate of 20 ounces per 100 gallons of water (1 tsp./gal.). Two (2) weeks later, begin the Preventative Program.

For Root (Ground) Mealybug and Fungus Gnat Control: When infestations of root mealybugs and fungus gnats are observed, the entire pot containing the infested plant should be drenched by submersion in a solution of ENSTAR® 5E at a rate of one (1) ounce per five (5) gallons of water. Submersion should continue until no more air bubbles are seen escaping from the fluid surface. This insures that the root system is saturated and the container is thoroughly treated. Repeat as necessary.

METHODS OF APPLICATION

Do not apply ENSTAR® 5E through any type of irrigation system.

Complete coverage of foliage is essential for control of whiteflies, aphids, mealybugs, fungus gnats and scales. For conventional spray equipment, mix the recommended amount of ENSTAR® 5E and water and spray both top and bottom surfaces of leaves until runoff. For other applications, follow the equipment manufacturer's specifications.

ENSTAR® 5E is compatible with other commonly used insecticides and miticides, such as PENTAC AQUAFLOW® and MAVRIK AQUAFLOW®. Before mixing in spray tank however, it is advisable to test physical compatibility by mixing all components in a small container in proportionate quantities. Read and follow all precautionary statements and label directions on tank mix products.

EFFECTS ON GREENHOUSE PLANTS

Slight to moderate injury has occasionally been noted on blooms under certain conditions. For insect control, it is suggested that applications be made in the pre-bloom stage. Before making large-scale application to any crop, a grower should test spray small numbers of plants to determine plant safety under his own cultural conditions.

Applications of ENSTAR® 5E have been made to greenhouse plants (azalea, chrysanthemum, dianthus, fuchsia, geranium, hydrangea, marigold, pelargonium, petunia, snapdragon) and vegetable seed crops in greenhouses (fava bean, green bean, spinach, sugar beet, tomato) without adverse effects.

Application of ENSTAR® 5E to poinsettia has produced bract damage of most varieties and foliar damage of some varieties. Some varieties of roses, such as yellow blooded roses, show delayed damage. Seeds from treated vegetable crops must be used only for subsequent planting. Portions of treated plants other than seeds must be disposed of according to local regulations and must not be used for food or feedstuffs.

10 of 10

LIMITATION OF WARRANTY AND LIMITATION OF LIABILITY

NOTICE: Read this Limitation of Warranty and Limitation of Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Sandoz or seller. All such risks shall be assumed by buyer or user.

Sandoz warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, under normal use conditions, subject to the risks described above. **SANDOZ MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

In no event shall Sandoz or seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SANDOZ OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SANDOZ OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sandoz and seller offer this product, and buyer and user accept it, subject to the foregoing limitations of warranty and limitation of liability, which may not be modified by any oral or written agreement.