U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (WH-567)	TERM OF ISSUANCE
WASHINGTON, D.C. 20460	
NOTICE OF PESTICIDE: REGISTRATION REREGISTRATION	NAME OF PESTICIDE PRODUCT
(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)	Thurlolne (10) hebballe power
NAME AND ADDRESS OF REGISTRANT (Include ZIP code)	
Sardez, Inc., Cro. Protection 4.0 Camino Iol Rio Scuti Buite 206	16/ 129 69
L 12360	
submitted to and accepted by the Registration Division price product always refer to the above U.S. EPA registration number of the product always refer to the above U.S.	
On the basis of information furnished by the registrant, the the Federal Insecticide, Fungicide, and Rodenticide Act.	above named pesticide is hereby Registered/Reregistered und
A copy of the labeling accepted in connection with this Re	gistration/Reregistration is returned herewith.
Act is not to be construed as giving the registrant a right t by others.	o exclusive use of the name or to its use if it has been covere
	्राक्ट (क्षा क्षा क्षा क्षा क्षा क्षा क्षा क्षा
	en e
a. I will take to desire the work of a constant of the department labeling.	is a second of the second of t
or the concernation is corotrol of the concernation in the concernation of the concern	and taken constituted. I are referred for a
A observed outy of the 1 Mar and	olosee for your receife.
[†] · · · · · · · · · · · · · · · · · · ·	Milm D. H. C.o. Rt Banayer (17)
	oticidesto enticado Brench Etratico Division (MS-7670)
aetosures	CORNEL CONTRACTOR CONT
ATTACHMENT IS APPLICABLE	
SIGNATURE OF APPROVING OFFICIAL	DATE
No constant of the constant of	

BEST COPY AVAILABLE



COMMERCIAL STRENGTH

A NATURAL MICROBIAL INSECTICIDE
TO CONTROL VARIOUS LISTED INSECTS
ON SHADE TREES, ORNAMENTAL FRUIT TREES, SHRUBS,
AND LISTED FRUIT AND VEGETABLES

ACTIVE INGREDIENT: Bacillus thuringiensis Berliner, potency of 16,000 International Units (at least 30 million viable spores) per milligram*

100.0%

*Equivalent to 7.2 billion International Units per pound.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARUS TO HUMANS: Avoid contact with eyes or open wounds, Avoid breathing spray mist.

ENVIRONMENTAL HAZARDS: Keep out of lakes, points or streams. Do not contaminate water by cleaning of equipment or dispusel of waste.

NOTICE: Saidul, Inc. makes no warranty, express or implied, including the warrinter of merchantability end/or triness for any particular purpose, concerning this material except those which are contained on Sandoa' label.

NET___POUNDS

EPA Eu. No. 36299-CA-1

LOT

EPA Reg. No. 11273





Sandaz, Inc., Crop Protection 480 Camina Del Rio South San Diego, CA 92108

DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation to use this product in a manner inconsistent with its labelling.

Mix 1/3 tablespoon per gallon of water. Shake sprayer occasionally during spraying. Apply surey thoroughly to cover both upper and under sides of folings as macts must sat THURICIDE to be controlled. Use of a spreader-sticker in the spray helps keep THURICIDE on the folings to maintain affectiveness where heavy dew or sight rains occur. Reapply after heavy rains, Apply at first sign of infestation and repeat at weekly intervals as necessary.

*The following table indicates the number of plants which can be adequately covered with one sation of spray mixture. For entire crops, the number of plants indicated is based on young sendlings. As they grow and mature, the number of plants which can be adequately covered will have to be reduced from 2 to 6 times.

NUMBER OF PLANTS COVERED/GALLON OF SPRAY MIXTURE

PLANT TYPE

200	 Clustered Cote Crops: Broccoli,
	Brussels Sprouts, Cabbage,
	Cauliflawer
300	 Lealy Cole Cropt: Collards, Kale,
	Mustard Greens, Turnip Greens
	+ Celary
500	- Beans or Spinach
100	Cucumbers or Melons
160	= Tomato Plants
30	- Grapevines
1 to 4	a Average size trees (ornamental frui
	or shada)
16	Dwarf ornamental trees
36	• Strub•
· ·	

HOW THURICIDE WORKS

THURICIDE causes paralysis of the digestive tract of the insects listed on this laber when they set it, literally causing them to starve to death, They stop feeding, but may otherwise appear unaffected for several days.

STORAGE AND DISPOSAL

CONTAINER DISPOSAL; Do not reuse container, Wrap and place in tresh cultection, STORAGE; Store in a cool place.

BEST COPY AVAILABLE

THURICIDE

PESTS CON'

Cabbage loog moth on: Bri Collards, Kale

Cabbage loo (Southeast on

Rindwarm ca Grape leaffai

hatching and feaves thorou

Fruit tree le Catifornia)

PESTS CONT

[Western U.S Webwerm, 6 humped Cate PESTS CON

Only)*, Fall Only), Tent (

PESTS CON

twestern U.S Onlyl*, Tent *For Gypsy

tion occurs, application.

SOX. If soos