

PM 23 55947-143
10/19
UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

AUG 29 1994

James S. Fickle, Ph.D.
Sandoz Agro, Inc.
1300 East Touhy Avenue
Des Plaines, IL 60018

Dear Dr. Fickle:

Subject: Proposed Label Amendment
Barricade 65WG Herbicide (Water Soluble Packs)
EPA Registration Number 55947-143
Your Submission Dated August 25, 1994

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable with the following provisions:

1) Per our telephone conversation of August 29, 1994, delete the following sentence from footnote #1 under the application rate chart for established turf:

"Consult your distributor or Sandoz representative for specific rate recommendations."

2) For the Tolerant Ornamental Species list, place the asterisk footnote, "Ornamental species only. Do not use on food producing trees or vines", at the end of the list, on each page, that the listing covers. If the list occupies only one page of the final printed label, then the footnote need only be placed at the list's end on that page.

A stamped copy is enclosed for your records. Please submit one (1) copy of final printed labeling, incorporating the above changes, before releasing the product for shipment.

Sincerely yours,

Joanne I. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

CONCURRENCES

SYMBOL	7505C						
SURNAME	J. Miller						
DATE	8/29/94						

201/18

BARRICADE 65WG Herbicide in Water Soluble Packs, 55947-143
August 24, 1994, pbarwsa.94b

1

**BARRICADE® 65 WG Herbicide
in Water Soluble Packs**

ACTIVE INGREDIENT:

Prodiamine [N³,N³-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine] 65.0%

INERT INGREDIENTS: 35.0%
Total 100.0%

**For selective preemergence control of grass and broadleaf weeds in established turf
grasses (excluding golf course putting greens), and landscape ornamentals.**

ACCEPTED
with COMMENTS
In EPA Letter Dated

KEEP OUT OF REACH OF CHILDREN

CAUTION

AUG 29 1994
Under the Federal Insecticide,
Fungicide and Rodenticide Act
as amended for the pesticide
registered under EPA Reg. No.
55947-143

See next page for additional precautionary statements and
statement of practical treatment.

EPA Reg. No. 55947-143
EPA Est. No.

Net Contents _____

Sandoz Agro, Inc.
1300 E. Touhy Avenue
Des Plaines, IL 60018

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact, while mixing or handling the concentrated material, may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE):

WPS USES:

Applicators and other handlers (other than mixers and loaders) who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) -- in general, agricultural-plant uses are covered -- must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Mixers and Loaders must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

NON-WPS USES:

Mixers and loaders who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) -- in general, only agricultural-plant uses are covered by the WPS -- must wear:

- Waterproof gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

BEST AVAILABLE COPY

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

STATEMENT OF PRACTICAL TREATMENT

If on skin: Wash with soap and water. Rinse thoroughly.

If inhaled: Remove to clear air.

If in eyes: Flush thoroughly with water for several minutes. Contact a physician if irritation persists.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. For terrestrial use, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

507-1

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

BARRICADE® 65WG Herbicide (BARRICADE) in this package is in 0.5 lb water soluble packs. These packs are intended to be added to the spray tank without opening (except for compatibility testing).

Do not remove water soluble packs from container except for immediate use. Reseal the outer container after use.

BARRICADE is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in:

BEST AVAILABLE COPY

- established turf grasses (excluding golf course putting greens) and lawns
- landscape ornamentals including perennials and wildflower plantings

BARRICADE controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when BARRICADE is activated by at least 0.5 in of rainfall or irrigation, or shallow (1 to 2 in) mechanical incorporation within 14 days after application and prior to weed seed germination.

Do not graze or feed livestock forage cut from areas treated with BARRICADE.

All applicable directions, restrictions, and precautions on the labels of EPA registered tank mix partners are to be followed.

Do not blend BARRICADE onto dry fertilizer or any other granular material.

Do not apply this product through any type of irrigation system.

Do not apply aerially.

Do not apply to golf course putting greens.

Mixing and Application

Apply Barricade in a minimum of 20 gal/A (0.5 gal/1000 ft²) of carrier (water and/or fluid fertilizer) using a calibrated, low pressure sprayer with 50 mesh or coarser screens. A broadcast boom or hand held wand designed for herbicide or insecticide application will provide the best results. Select nozzle pressure and gallonage to provide complete coverage.

A spray colorant may be used to modify the color of BARRICADE to improve application accuracy by minimizing swath skips and overlaps.

Thorough mixing of BARRICADE in the spray tank is essential for uniform application. Fill the spray tank half full with clean water or fluid fertilizer. Start agitation and check to ensure it is working properly, then add the desired number of water soluble packs of BARRICADE directly into the tank. Do not remove BARRICADE from the water soluble pack as it will dissolve in the tank.

Add the rest of the carrier to obtain the final spray volume. Maintain vigorous agitation to completely dissolve the bags and suspend the contents before and during the application. This will ensure a well mixed spray suspension.

Thoroughly clean the sprayer after use by flushing the system with water containing a detergent. Refer to Pesticide Disposal Section of this label for waste disposal. Do not allow spray suspension to dry in the tank.

Mixing Order for Tank Mixtures

When mixing BARRICADE with other components (carrier and partner pesticide products), add

BEST AVAILABLE COPY

70418

the products to the spray tank in the following order:

1. Water dispersible granules (WG formulations) and wettable powders (WP formulations). Wettable powders should be premixed with a small amount of water to form a slurry before addition; water dispersible granules can be added directly during filling. Allow the product to disperse completely before other products are added.
2. Flowable liquids (FL) or suspension concentrates (SC).
3. Emulsifiable concentrates (EC).
4. Surfactants approved for application to turf or ornamentals listed on this label. Check surfactant label before use.

Compatibility Test

Before mixing BARRICADE with other pesticides in the spray tank, test the compatibility by mixing all components (carrier and pesticide products) in a small container in proportionate quantities (see following table).

Amount of Component to Add to One Pint of Spray Carrier
(Assuming Volume is 25 Gallons per Acre)

COMPONENT FORMULATIONS	—RATE PER—		LEVEL TEASPOONS*
	ACRE	1000 ft ²	
Dry	1.0 lb	0.4 oz	1.5
Liquid	1.0 pt	0.4 oz	0.5

* Remove the required amount of BARRICADE from a water soluble pack for the compatibility test, saving the remainder in the pack for spray mixture preparation. The amount used will not decrease weed control if the pack is used according to the Maximum Rate table under Rates of Application.

If components do not ball-up or form flakes, sludge, gels, oily films or layers, or other precipitates, then the tested spray mix is compatible. Incompatibility in any of the above forms will usually occur within 5 minutes after mixing.

If components are incompatible, the use of a compatibility agent is recommended. Rerun the above compatibility test with commercially available compatibility agents until a suitable compatibility agent is identified.

ESTABLISHED TURF

BARRICADE is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of BARRICADE that may be applied per year is given for each turf grass species in the Maximum Annual Rates section of this label.

Most effective weed control in turf grasses will be obtained when BARRICADE is activated by at

BEST AVAILABLE COPY

20719

BARRICADE 65WG Herbicide in Water Soluble Packs, 55947-143
August 24, 1994, pbarwspa.94b

7

least 0.5 in of rainfall or irrigation within 14 days after application and prior to weed seed germination. See the map below for approximate crabgrass seed germination dates.

CRABGRASS SEED GERMINATION DATE

Approximate Date

)
)

)

70417

USE PRECAUTIONS

The following use precautions apply to the use of BARRICADE in turf grasses and lawns:

Application of BARRICADE may thin emerged annual bluegrass and newly overseeded grasses.

Do not apply to overseeded turf for 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if BARRICADE is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

Do not apply to newly set sod until the following year.

Application of BARRICADE to turf stressed by drought, low fertility, or pest damage may result in turf injury.

Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.

Do not apply BARRICADE to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (Poa annua) are desirable species.

10-2-17

RATES OF APPLICATION

BARRICADE may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. BARRICADE will not control established weeds.

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by BARRICADE is related to the rate applied.

Maximum Annual Rates

BARRICADE is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Maximum Rate / Calendar Year of BARRICADE by Turf Grass Species ¹			
Turf Species	ft ² /pack	lb/A	oz/1000 ft ²
Bermudagrass ² Bahigrass Centipedegrass Seashore Paspalum Tall Fescue (including turf-type) Zoysia	9638	2.30	0.83
Buffalograss Kentucky Bluegrass Perennial Ryegrass	14,545	1.50	0.55
Creeping Red Fescue St. Augustinegrass	20,000	1.15	0.40
Creeping Bentgrass	21,622	1.00	0.37

¹ These are the maximum rate per calendar year by species limitations. Consult your distributor or Sandoz representative for specific rate recommendations.

² May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 0.80 lb/A or 0.30 oz/1000 ft². Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded.

BEST AVAILABLE COPY

110418

Weeds Controlled

When used in accordance with this label BARRICADE will provide control of the following weeds:

Crabgrass (large, smooth)	Carpetweed
Annual Bluegrass (<u>Poa annua</u>)	Common Chickweed
Goosegrass ¹	Common Lambsquarter
Annual Foxtails	Common Purslane
Barnyardgrass	Henbit
Broadleaf Signalgrass	Knotweed
Johnsongrass (from seed)	Pigweed
Rescuegrass ³	Prostrate Spurge
	Shepherd's purse ²
	Yellow woodsorrel (from seed)

-
1. In many areas a single application of 1 to 2.3 lb/A will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, make an initial application of 1 to 1.5 lb/A followed after 60 to 90 days by a second application of 1.3 to 0.8 lb/A, respectively, for season long goosegrass control. Do not exceed the maximum rate for turf grass species listed in the Maximum Annual Rate Table above.
 2. Applications for this weed should be made in late summer, fall or winter prior to germination.
 3. Suppression only.

BEST AVAILABLE COPY

10/18

Overseeding / Reseeding

BARRICADE will inhibit the germination of turf species if overseeded too soon after application. Wait 4 to 6 months per 0.75 lb/A of BARRICADE applied when grass seed is to be broadcasted over the turf surface. This overseeding interval can be shortened to 3 to 4 months per 0.75 lb/A when grass seed is drilled directly into the soil. The interval between application and overseeding / reseeding is dependent on the application rate, soil temperature, and soil moisture. A shorter period is required if growing conditions are warm and wet, while the longer interval is needed if conditions are cool and dry.

Turf Tank Mixes

BARRICADE may be tank mixed with other registered turf herbicides for broader spectrum weed control. Tank mixes may be used only in states where the tank mix partner (s) are registered for the application site and the turf species listed above.

Consult the label(s) of the individual tank mix partner (s) for use, rate recommendations, application timing, weeds controlled, turf grass safety and specific precautions and/or restrictions.

Before mixing tank mix partner (s) in the spray tank it is advisable to test compatibility by mixing the products in a small container. See the **Compatibility Test** section under **Mixing and Application**.

130A18

LANDSCAPE ORNAMENTALS

BARRICADE may be applied for residual preemergence weed control in landscape ornamentals, including perennials and wildflower plantings..

Use Rates

Apply BARRICADE at 1 to 2.3 lb/A equivalent to 0.36 to 0.83 oz/1000 ft² in fall and/or spring. Use the higher rate of application for longer control periods. Sequential or single applications are allowed so long as the total amount of product applied does not exceed the maximum annual application rate of 2.3 lb/A.

Application Timing and Use Information

BARRICADE may be applied to newly transplanted and established landscape ornamentals over the top or as a directed spray. Irrigation or rainfall soon after application will wash residues off plant foliage and activate BARRICADE in the soil.

BARRICADE may be applied at any time to established plants. Delay application to allow soil to settle around new transplants and water thoroughly before applying BARRICADE.

BARRICADE is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when BARRICADE is activated by at least 0.5 in of rainfall or irrigation, or with shallow (1 to 2 in) mechanical incorporation, below weed seeds germinate. Acceptable control will be obtained when BARRICADE is activated within 14 days after application.

Best weed control is obtained when BARRICADE is applied to soil free of clods, weeds, and debris such as leaves and mulch. Prior to application of BARRICADE, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

Tank Mixtures for Use on Ornamentals

BARRICADE may be tank mixed with other registered landscape herbicides for a broader spectrum of preemergence control or control of emerged weeds. Tank mixes are for use only in states where the tank mix partner (s), application site and intended use pattern are registered.

Follow the label (s) of potential tank mix partner (s) for use, rate recommendations, application timing, weeds controlled, phytotoxicity and specific precautions and / or restrictions.

Before combining tank mix partners in the spray tank it is advisable to test compatibility by mixing the products in a small container. See the **Compatibility Test** section under **Mixing and Application**.

Tolerant Ornamental Species

BARRICADE will not harm most established ornamental trees, shrubs and flowers. The species listed below are tolerant to BARRICADE. BARRICADE may be applied over the top of these species without injury.

BEST AVAILABLE COPY

11/07/18

Scientific name

Common name

<hr/>	<hr/>
Acer palmatum	Japanese Maple
A. platanoides	Norway Maple
Actinidia chinensis	Kiwi*
Agapanthus africanus	Lily-of-the-Nile
Arctostaphylos densiflora	Vine Hill Manzanita
Arctotheca calendula	Cape Weed
Aucuba japonica	Japanese Aucuba
Barberis gladwynensis	Barberry
B. julianae	Wintergreen Barberry
B. mentorensis	Mentor Barberry
B. thunbergii	Japanese Barberry
B. verruculosa	Warty Barberry
Buxus microphylla	Japanese Boxwood
Callistemon viminalis	Weeping Bottlebrush
Calluna vulgaris	Scotch Heather
Carpobrotus edulis	Hottentot Fig (Ice Plant)
Cassia artemisoides	Feathery Cassia
Ceanothus rigidus	Wild Lilac
Chamaecyparis pisifera	False Cypress
Cleyera japonica	Cleyera
Citrus spp*	Citrus species
Cornus florida	Flowering Dogwood
C. stolonifera	American Dogwood
Cortaderia selloana	Pampas Grass
Cotoneaster apiculatus	Cranberry Cotoneaster
C. buxifolius	Cotoneaster
C. Jammeri	Bearberry Cotoneaster
C. microphyllus	Rockspray Cotoneaster
Cretaeagus spp.	Hawthorne
Cupressus sempervirens	Italian Cypress
Delasperma alba	White Trailing Ice Plant
Dodonea viscosa	Hop Bush
Elaeagnus pungens	Silverberry
Euonymus fortunei	Wintercreeper
E. japonica	Evergreen Euonymus
E. kiautschovica	Spreading Euonymus
Fatsia japonica	Japanese Aralia
Forsythia intermedia	Border Forsythia
F. viridissima	Greenstem Forsythia

Scientific name

Common name

<hr/>	<hr/>
Gardenia jasminoides	Gardenia, Cape-Jasmine
Hedera helix	English Ivy
Hibiscus rosa-sinensis	Chinese Hibiscus

BEST AVAILABLE COPY

150718

Ilex cornuta	Chinese Holly
I. crenata	Japanese Holly
I. opaca	American Holly
I. pemyi	Holly
Jasminium nudiflorum	Winter Jasmine
Juniperus chinensis	Chinese Juniper
J. conferta	Shore Juniper
J. horizontalis	Creeping Juniper
Juglans sp.	Walnut*
Justicia brandegeana	Shrimp Plant
Lagerstromia indica	Crepe Myrtle
Ligustrum amurense	Amur Privet
L. japonicum	Japanese Privet
L. lucidum	Glossy Privet
Liriope muscari	Big Blue Lily Turf
Lonicera japonica	Japanese Honeysuckle
L. tatarica	Tatarian Honeysuckle
Maleophora luteola	Ice Plant
Malus sp.	Crabapple*
Nandina domestica	Heavenly Bamboo
Olea europa	Olive*
Ophiopogon japonicus	Mondo Grass
Osteospermum fruticosum	Trailing African Daisy
Oxydendrum arboreum	Sourwood
Persea americana	Avocado*
Photinia fraseri	Frasier's Photinia
Picea abies	Norway Spruce
Pieris japonica	Lily-of-the-Valley Shrub
Pinus brutia	Calabrian Pine
P. canariensis	Canary Island Pine
P. halepensis	Aleppo Pine
P. nigra	Austrian Black Pine
P. radiata	Monterey Pine
P. thunbergiana	Japanese Black Pine
P. strobus	Eastern White Pine
P. sylvestris	Scotch Pine
Pistachio sp.	Pistachio*
Pittosporum rhombifolium	Queensland Pittosporum
P. tobira	Mock Orange
Podocarpus macrophyllus	Japanese Yew
Prunus laurocerasus	English Laurel

Scientific name

Common name

Prunus spp.

Almond, Apricot, Nectarine,
Peach, Plum and Prune*

Pyracantha coccinea
P. fortuneana

Firethorn
Firethorn

BEST AVAILABLE COPY

160418

- | | |
|---------------------------|------------------------|
| P. koidzumii | Firethorn |
| Pyrus sp. | Bradford Pear sp. |
| Quercus rubra | Red Oak |
| Raphiolepis indica | Indian Hawthorne |
| Rhododendron | 'Coral Bells' |
| (including Azalea) | 'Formosa' |
| | 'Hino-crimson' |
| | "PJM" |
| | 'Roseum Elegans' |
| Rosa banksiae | Lady Bank's Rose |
| Rosmarinus officinalis | Rosemary |
| Rumohra adiantiformis | Leatherleaf Fern |
| Santolina virens | |
| Sedum album | Stonecrop |
| Syzygium paniculatum | Japanese Boxcherry |
| Taxus cuspidata | Japanese Yew |
| T. media | Yew |
| Thuja occidentalis | American Arborvitae |
| Trachelospermum asiaticum | Star Jasmine |
| Tsuga canadensis | Canada Hemlock |
| Viburnum japonicum | Japanese Viburnum |
| V. odoratissimum | Sweet Viburnum |
| V. plicatum | Japanese Snowball |
| V. rigidum | Canary Island Viburnum |
| V. tinus | Laurustinus |
| V. trilobium | Cranberry Bush |
| V. wrightii | |
| Vinca minor | Dwarf Periwinkle |
| Vitis sp. | Grape* |
| Weigela florida | Old Fashioned Weigela |
| Yucca aloifolia | Spanish Bayonet |
| Y. filamentosa | Yucca, Adam's Needle |

* Ornamental species only. Do not use on food producing trees or vines.

BEST AVAILABLE COPY

1/24/18

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE

Store in original container away from fertilizer, feed, or food stuffs and separated from other pesticides.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Paper bags and boxes: Completely empty container into application equipment. Then dispose of empty bag or box in a sanitary landfill or at an incineration facility; or, if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

BEST AVAILABLE COPY

189718

LIMITATION OF WARRANTY AND LIMITATION OF LIABILITY

NOTICE: Read this Limitation of Warranty and Limitation of Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Sandoz or seller. All such risks shall be assumed by buyer or user.

Sandoz warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, under normal use conditions, subject to the risks described above. **SANDOZ MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

In no event shall Sandoz or seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SANDOZ OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE, RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SANDOZ OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sandoz and seller offer this product, and buyer and user accept it, subject to the foregoing limitations of warranty and limitation of liability, that may not be modified by any oral or written agreement.

Barricade® is a registered trademark of Sandoz Ltd. ©

BEST AVAILABLE COPY