

BANVEL® 10G GRANULAR HERBICIDE

FOR NONCROPLAND AREAS
SUCH AS FENCE ROWS, ROADWAYS, WASTELAND, AND
SIMILAR NONCROPLAND

ACTIVE INGREDIENT:

Dicamba (3,6-dichloro-*p*-anisic acid) 10.0%


INERT INGREDIENTS. 90.0

TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN

WARNING

See side panel for additional precautionary statements



EPA Reg. No. 55947-164

Net Contents:

EPA Est. No.

Sandoz Agro, Inc.
Des Plaines, IL 60018

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING**

Harmful if swallowed. Avoid breathing vapors. Dust may cause eye irritation. Avoid contact with skin, eyes and clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

Do not store near feed or food stuffs.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by the cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT

Do not contaminate irrigation ditches or water used for domestic purposes.

Banvel 10G Herbicide may cause injury to desirable trees or plants, particularly beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes, and other broadleaf plants when contacting their roots, stems, or foliage.

Do not treat areas where the possible downward movement into the soil or surface washing may cause contact of herbicide with the roots of desirable plants such as trees and shrubs.

To avoid injury to desirable plants, equipment used for BANVEL 10G Herbicide should be thoroughly cleaned before reuse to handle or apply any other chemicals.

BANVEL 10G Herbicide may temporarily stunt or eliminate the growth of grass in the treated areas.

WEEDS AND BRUSH CONTROLLED

BANVEL 10G Herbicide, when applied in accordance with this label, will give control or growth suppression of many BIENNIAL and PERENNIAL broadleaf weeds as well as many WOODY brush and vine species including:

BIENNIAL and PERENNIAL BROADLEAVES

Bindweed, Field
Blueweed, Texas
Bursage, Skeletonleaf
(Bur Ragweed, Povertyweed)
Bursage, Woollyleaf
(Lakeweed)
*Fern, Bracken
Knapweed, Russian
Nightshade, Silverleaf
Ragwort, Tansy
Sowthistle, Perennial
*Spurge, Leafy
Starthistle, Yellow
Thistle, Artichoke
Thistle, Bull
Thistle, Canada

- * Noted weeds require higher rates of BANVEL 10G than other listed weeds.
See APPLICATION RATES AND TIMING section of this label.

WOODY BRUSH and VINES

Creosotebush
Elm, Siberian
Kudzu
Olive, Autumn
Persimmon, Eastern
Plum, Sand
Saltcedar
Sumac
Tarbush

APPLICATION RATES AND TIMING

BANVEL 10G Herbicide may be applied using standard ground or air granular application equipment for broadcast or spot treatments. BANVEL 10G Herbicide is activated when there is sufficient moisture to move the herbicide into the root zone of plants to be controlled. Application may be made any time when soil is not frozen. However, best performance usually occurs when treatments are made in the spring just before or during active growth of the weeds and brush. CONSULT YOUR LOCAL OR STATE AUTHORITIES FOR THE BEST TIMING OF APPLICATION.

) Application rates are given below. Use the higher level of listed rate range when treating dense vegetative growth.

- Apply ~~20 to 40 pounds (2-4 lb ai) of BANVEL 10G Herbicide per treated~~ acre to control listed BIENNIAL and PERENNIAL broadleaf weeds. For spot treatments, use 1/2 to 1 pound of product per 1000 square feet for control of listed BIENNIAL and PERENNIAL weeds.
- Apply ~~40 to 80 pounds (4-8 lb ai) of BANVEL 10G Herbicide per treated~~ acre to control noted (*) BIENNIAL and PERENNIAL broadleaf weeds. For spot treatments, use 1 to 2 pounds of product per 1000 square feet for noted (*) BIENNIAL and PERENNIAL weeds.
- Apply ~~80 to 100 pounds (8-10 lb ai) of BANVEL 10G Herbicide per~~ treated acre to control WOODY brush and vines. For spot treatments, use ~~2 to 2 1/2~~ 2 to 2 1/2 pounds of product per 1000 square feet for control of listed WOODY BRUSH and VINES. Uniformly spread granules over the area where the plant roots are growing

In most cases these treatments will not control weed seedlings which germinate one to two years after treatment. Once the effect of the herbicide has been lost, a follow-up program for seedling control or other cultural practices should be instituted.

Do not apply more than 20 pounds of BANVEL 10G per treated acre per year.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage or disposal.

STORAGE

Store in original container in a well-ventilated area away from fertilizer, feed or foodstuffs and separated from other pesticides.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Triple rinse (or equivalent) plastic pails. For paper or plastic bags, or for fiber drums with liners, completely empty into application equipment by shaking and tapping container. Offer pails or drums for recycling or reconditioning or puncture before disposal. Dispose of bags, liners, or punctured containers in sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

LIMITATION OF WARRANTY AND LIMITATION OF LIABILITY

NOTICE: Read this Limitation of Warranty and Limitation of Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Sandoz or seller. All such risks shall be assumed by buyer or user. Sandoz warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, under normal use conditions, subject to the risks described above. **Sandoz makes no other express or implied warranty of fitness or of merchantability or any other express or implied warranty.** In no event shall Sandoz or seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **The exclusive remedy of the user or buyer, and the exclusive liability of Sandoz or seller for any and all claims, losses injuries or damages (including claims based on breach of warranty, contract, negligence, tort, strict liability or otherwise) resulting from the use or handling of this product, shall be the return of the purchase price of the product or, at the election of Sandoz or seller, the replacement of the product.** Sandoz and seller offer this product, and buyer and user accept it, subject to the foregoing limitations of warranty and limitation of liability, which may not be modified by any oral or written agreement.

REGISTERED TRADEMARKS

BANVEL® is a registered trademark of Sandoz Ltd.