

PM 23 56077-31

477-96 10/4

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number
225575

Application for Pesticide - Section I

1. Company/Product Number 56077-31	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Butoxone Herbicide	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Cedar Chemical Corp. 5100 Poplar Ave., Suite 2414 Memphis, TN 38137 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION
APR 24 1970

Explanation: Use additional page(s) if necessary. (For section I and Section II.)
Addition of an alternate name - Butoxone 175 Herbicide

NOTIFICATION
APR 24 1996

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Metal Plastic Glass Paper Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Jeanine H. Betsher	Title Mgr., Regulatory Affairs	Telephone No. (Include Area Code) 901-685-5348
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>J.H. Betsher</i>	3. Title Mgr., Regulatory Affairs	
4. Typed Name Jeanine H. Betsher	5. Date 4/16/96	



Butoxone 175

Herbicide

For the control of seedling broadleaf weeds in peanuts, soybeans, and seedling stands of alfalfa, and for use on established stands of alfalfa.

ACTIVE INGREDIENT:

Dimethylamine salt of 4-(2,4-dichlorophenoxy)

butyric acid

23.00%

INERT INGREDIENTS:

77.00%

TOTAL

100.00%

*Equivalent to 19.5% by weight of 4-(2,4-dichlorophenoxy) butyric acid, or not less than 1.75 lbs. 4-(2,4-DB) acid per gallon at 68°F.

EPA Registration No. 56077-31

EPA Establishment No.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Drink 1 to 2 glasses of water and induce vomiting by touching back of throat with finger. Do not give anything by mouth to an unconscious person. Get medical attention.

If On Skin: Wash contaminated areas thoroughly with soap and water. Do not reuse contaminated clothing until washed. Get medical attention if irritation persists.

If In Eyes: Flush with copious amounts of clean water for 10 - 15 minutes. Get medical attention.

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin, eyes, and clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- long-sleeved shirt and long pants
- waterproof gloves
- shoes plus socks
- chemical-resistant headgear for overhead exposure

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FOR CHEMICAL SPILL, LEAK, FIRE OR EXPOSURE CALL TOLL FREE

1-800-424-9300

AGRICULTURAL CHEMICAL

DO NOT SHIP OR STORE WITH FOODS, FEEDS, DRUGS, OR CLOTHING

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not contaminate water intended for irrigation or domestic purposes. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls
- waterproof gloves
- shoes plus socks
- chemical-resistant headgear for overhead exposure

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not store this product near fertilizers, seeds, insecticides, or fungicides. Store at temperatures above 32°F. If product is allowed to freeze, warm to 50°F and agitate thoroughly before using. Containers should not be stacked more than six (6) high. Reclose all partially used containers by thoroughly tightening screw cap. Damaged or leaking containers which cannot be used immediately should be transferred to suitable sound containers and properly marked. Absorb any spill with a suitable clay absorbent and dispose of as indicated under "Pesticide Disposal".

For safety and prevention of unauthorized use, all pesticides should be stored in locked facilities.

To prevent accidental misuse, different pesticides should be stored in separate areas with enough distance between to provide clear identification.

Opened, partially used pesticides should be stored in original labeled containers when possible. When transfer to another container is necessary because of leakage or damage, carefully mark and identify contents of the new container.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency or the hazardous waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Apply this product only as specified on this label.

NOTE: Avoid spray drift to susceptible plants and crops: cotton, tobacco, tomatoes, ornamentals, etc. Coarse sprays are less likely to drift.

Spray equipment previously used to apply another phenoxy must be thoroughly cleaned with alkali and water prior to use with Butoxone. Similarly, after using Butoxone wash sprayer thoroughly before using again to spray susceptible crops. Do not

use near greenhouses. Observe all caution and limitations on labeling of all products used in mixtures.

Use agitation to keep solution well mixed, especially if it has been allowed to stand unagitated.

Calibrate sprayer accurately.

DO NOT APPLY WHEN CROP IS STRESSED FROM LACK OF MOISTURE.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

GROUNDWATER CONTAMINATION: Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-DB have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-DB pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will prevent groundwater contamination.

ALFALFA

APPLICATION:

For use in seedling alfalfa, spray when the crop has reached the 1 to 2 trifoliate leaf stage and growing conditions are good. In established alfalfa certain weeds will emerge in the fall and over winter in a rosette stage. Best control of these weeds will result from application in late fall or early winter rather than in the spring. Do not apply after flowering.

WHEN TO APPLY:

Weeds must be in the young seedling stage and actively growing to achieve satisfactory results. For best results spray weeds in the 2 to 5 leaf stage of growth.

AMOUNT TO USE:

Apply Butoxone as an overall spray by ground sprayer or airplane. Apply at rates listed below according to weed problems. Use the higher rate if weeds are past the seedling stage. Aerial applications should be made in at least 5 gallons of water per acre, while ground-sprayer applications should be in a minimum of 10 gallons of water per acre. Higher spray gallonage per acre will give better coverage and weed control. Spray gallonage should give adequate coverage of the weeds without run-off. The use of a non-ionic or non-ionic/anionic surfactant or crop oil concentrate approved for agricultural uses at their label rates will usually result in better weed control, especially if weeds are beyond the seedling stage.

BROADLEAF WEEDS	Rate of Butoxone per Acre
Annual Morningglory (Ipomoea spp.)	4 1/2 pints
Cocklebur (Xanthium spp.)	
Common Lambsquarters (Chenopodium album)	
Jimsonweed (Datura stramonium)	
Kochia or Mexican Fireweed (Kochia scoparia)	
Pigweed (Amaranthus spp.)	
Velvetleaf (Abutilon theophrasti)	
Wild Turnip (Brassica campestris)	
Black Mustard (Brassica nigra)	6 1/2 pints
Buckhorn Plantain (Plantago lanceolata)	
Common Ragweed (Ambrosia artemisiifolia)	
Curly Dock (Rumex crispus)	
Field Pennycress (Fanweed or Stinkweed) (Thlaspi arvense)	
Hedge Smartweed (Polygonum scandens)	
Ladysthumb (Polygonum persicaria)	
Prickly Lettuce (Lactuca serriola)	
Shepherdspurse (Capsella bursa-pastoris)	
Sweetclover (Volunteer plants) (Melilotus spp.)	
Wild Beet (Beta maritima)	
Wild Mustard (Brassica kaber)	

Do not graze established alfalfa, nor feed straw or hay from established alfalfa to livestock within 30 days after application.

Do not graze or feed SEEDLING alfalfa within 60 days after application. Irrigation, in particular overhead sprinkler irrigation, should be delayed as long as possible (10 days or more) following application of Butoxone to avoid washing the chemical into the root zone. Do not apply when crop is stressed from lack of moisture. DO NOT use on peas.

Apply as a postemergent spray. When properly timed there is little or no effect on the crop. In established alfalfa there may be some twisting of stems and malformation of leaves. This condition is usually outgrown.

Do not spray when daytime temperatures are expected to exceed 90°F within the next 2 or 3 days. Do not apply if temperatures are likely to fall below 40°F during or shortly after treatment.

Use of a surfactant or crop oil concentrate in the desert areas of California and Arizona may cause some crop injury under certain climatic and crop stress conditions. Do not use a surfactant unless possible crop injury is acceptable.

PEANUTS

APPLICATION:

Apply Butoxone postemergent as an overall spray by ground sprayer or airplane. With ground applications use a boom sprayer with flat fan-type nozzle. Adjust the height of the boom above the crop so the growing terminals of weeds are sprayed. Use low pressure (about 20 to 40 lbs. per square inch). Do not use boom jet nozzles. With airplane applications Butoxone has been successfully applied using 5 gallons of total solution per acre. The height of the application should insure THOROUGH COVERAGE of the tops of the weeds. Best control has resulted from airplane application with coarse spray when cocklebur is at least crop high and actively growing. Other weed species should be in the seedling stage and actively growing. Airplane application is not recommended in the vicinity of susceptible crops.

Do not apply if peanuts are suffering from lack of moisture.

Amount to Use: Apply the recommended rates as indicated in the table below.

BROADLEAF WEEDS	Rate of Butoxone per Acre
Cocklebur (Xanthium spp.)	1 pint
Annual Morningglory (Ipomoea spp.)	1 1/2 pints
Common Lambsquarters (Chenopodium album)	
Common Ragweed (Ambrosia artemisiifolia)	
Jimsonweed (Datura stramonium)	
Sicklepod (Coffeeweed) (Cassia obtusifolia)	
Velvetleaf (Abutilon theophrasti)	

Do not apply more than two (2) applications per season; second application should be made no later than the late bloom stage of peanuts (about 90 to 100 days after planting). Do not feed treated peanut vines or peanut hay to livestock. Do not apply to peanuts within 30 days of harvest.

SOYBEANS

APPLICATION: Apply Butoxone postemergent as an overall spray by ground sprayer or airplane. With ground applications use a boom sprayer with flat fan-type nozzles. Adjust the height of the boom above the crop so the growing terminals of all weeds are sprayed. Use low pressure (about 20 to 40 lbs. per square inch). Do not use boom jet nozzles. With airplane applications Butoxone has been successfully applied using 5 gallons of total spray solution per acre. The height of the application should insure thorough coverage of the tops of the weeds. Best control has resulted when cocklebur is at least crop high and actively growing. Airplane application is not recommended in the vicinity of susceptible crops. Do not apply if soybeans are suffering from lack of moisture.

TOPICAL (OVERHEAD) APPLICATIONS

Apply from 7 to 10 days before soybeans bloom through mid-bloom. A good indication that the beans are about to bloom is when the soybean plants have turned a dark green color. If

3044